

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**



B6

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
WO 03/014375 A2

- (51) International Patent Classification<sup>7</sup>: C12Q
- (21) International Application Number: PCT/US02/25319
- (22) International Filing Date: 9 August 2002 (09.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- |            |                                |    |
|------------|--------------------------------|----|
| 60/311,378 | 9 August 2001 (09.08.2001)     | US |
| 60/313,932 | 21 August 2001 (21.08.2001)    | US |
| 09/952,680 | 13 September 2001 (13.09.2001) | US |
| 60/338,186 | 13 November 2001 (13.11.2001)  | US |
| 60/349,959 | 18 January 2002 (18.01.2002)   | US |
| 60/364,486 | 13 March 2002 (13.03.2002)     | US |
| 60/367,991 | 25 March 2002 (25.03.2002)     | US |
| 60/369,887 | 4 April 2002 (04.04.2002)      | US |
| 60/376,744 | 1 May 2002 (01.05.2002)        | US |
| 60/385,097 | 31 May 2002 (31.05.2002)       | US |
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:
- |          |                                |
|----------|--------------------------------|
| US       | 60/311,378 (CIP)               |
| Filed on | 9 August 2001 (09.08.2001)     |
| US       | 60/313,932 (CIP)               |
| Filed on | 21 August 2001 (21.08.2001)    |
| US       | 60/338,186 (CIP)               |
| Filed on | 13 November 2001 (13.11.2001)  |
| US       | 60/349,959 (CIP)               |
| Filed on | 18 January 2002 (18.01.2002)   |
| US       | 60/364,486 (CIP)               |
| Filed on | 13 March 2002 (13.03.2002)     |
| US       | 60/367,991 (CIP)               |
| Filed on | 25 March 2002 (25.03.2002)     |
| US       | 60/369,887 (CIP)               |
| Filed on | 4 April 2002 (04.04.2002)      |
| US       | 60/376,744 (CIP)               |
| Filed on | 1 May 2002 (01.05.2002)        |
| US       | 60/385,097 (CIP)               |
| Filed on | 31 May 2002 (31.05.2002)       |
| US       | 09/952,680 (CIP)               |
| Filed on | 13 September 2001 (13.09.2001) |
- (71) Applicant (for all designated States except US): ARCH-  
CHEMIX CORPORATION [US/US]; 20 Hampden  
Street, Boston, MA 02111 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): STANTON, Martin  
[US/US]; 64 Harvard Road, Stow, MA 01775 (US).  
EPSTEIN, David [US/US]; 56 Pine Street, Belmont,  
MA 02478 (US). HAMAGUCHI, Nobuko [JP/US]; 7  
Dartmouth Drive, Framingham, MA 01701 (US). KURZ,  
Markus [DE/US]; 44 Pine Street, Newton, MA 02465  
(US). KEEFE, Tony [GB/US]; 97 1/2 Thorndike Street,  
#1, Cambridge, MA 02139 (US). WILSON, Charles  
[US/US]; 229 Lexington Road, Concord, MA 01742  
(US). GRATE, Dilara [US/US]; 16 Wheelock Terrace,  
Waltham, MA 02453 (US). MARSHALL, Kristin, A.  
[US/US]; 15 College Avenue, Arlington, MA 02472 (US).  
MCCAULEY, Thomas [US/US]; 7 Spencer Avenue,  
Somerville, MA 02144 (US). KURZ, Jeffrey [US/US];  
431 Broadway, Apartment 23, Somerville, MA 02145  
(US).
- (74) Agent: ELRIFL, Ivor, R.; Mintz, Levin, Cohn, Ferris,  
Glovsky and Popeo, P.C., One Financial Center, Boston,  
MA 02111 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished  
upon receipt of that report

[Continued on next page]

(54) Title: NUCLEIC ACID SENSOR MOLECULES AND METHODS OF USING SAME

(57) Abstract: Methods for engineering a nucleic acid sensor molecule are provided. Biosensors comprise a plurality of nucleic acid sensor molecules labeled with a first signaling moiety and a second signaling moiety. The nucleic acid sensor molecules recognizes target molecules which do not naturally bind to DNA. Binding of a target molecule to the sensor molecules triggers a change in the proximity of the signaling moieties which leads to a change in the optical properties of the nucleic acid sensor molecules on the biosensor. Reagents and systems for performing the method are also provided. The method is useful in diagnostic applications and drug optimization.

WO 03/014375 A2

WO 03/014375 A2



*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## NUCLEIC ACID SENSOR MOLECULES AND METHODS OF USING SAME

5

### Field of the Invention

The invention relates generally to nucleic acids and more particularly to nucleic acid sensor molecules containing a catalytically active domain that is modulated upon recognition of a target by a target modulation domain of the nucleic acid sensor molecule.

10

### Background of the Invention

In addition to carrying genetic information, nucleic acids can adopt complex three-dimensional structures. These three-dimensional structures are capable of specific  
15 recognition of target molecules and, furthermore, of catalyzing chemical reactions. Nucleic acids will thus provide candidate detection molecules for diverse target molecules, including those which that do not naturally recognize or bind to DNA or RNA.

A nucleic acid which binds to a non-nucleic acid target molecule through non-Watson-Crick base pairing is termed an aptamer. In aptamer selection, combinatorial  
20 libraries of oligonucleotides are screened *in vitro* to identify oligonucleotides, or aptamers, which bind with high affinity to pre-selected targets. Both small biomolecules (*e.g.*, amino acids, nucleotides, NAD, S-adenosyl methionine, chloramphenicol), and large biomolecules (thrombin, Ku, DNA polymerases) are effective targets for aptamers. Aptamer biosensors have been used to detect specific analyte molecules. For example, fluorescently labeled  
25 anti-thrombin aptamers attached to a glass surface have been used to directly detect the presence of thrombin proteins in a sample by detecting changes in the optical properties of the aptamers (Potyrailo, et al., 1998). In this method continuous binding of thrombin to the labeled aptamer is requisite for detection to occur, since the concentration of thrombin in a test sample is monitored by directly detecting fluorescent emission of the aptamer-ligand  
30 complex. Another method of detecting binding of a ligand to an aptamer has also been described which relies on the use of fluorescence-quenching pairs whose fluorescence is sensitive to changes in secondary structure of the aptamer upon ligand binding (Stanton, et al. 2000) to form a fluorescent aptamer-ligand complex, and again, continuous binding of the ligand to the aptamer is required for signal generation and, hence, for detection to occur.  
35 A limitation with this type of aptamer-derived biosensor is that ligand-mediated changes in

secondary structure were engineered into the aptamer molecule via a laborious engineering process in which four to six nucleotides were added to the 5' end of the aptamer that was complementary to the bases at the 3' end of the thrombin binding region. In the absence of thrombin, this structure forms a stem loop structure, while it forms a G-quartet structure in the presence of thrombin. Fluorescent and quenching groups attached to the 5' and 3' end signal this change. In aptamer-based detection without the use of amplification steps, assay sensitivity and hence the limit of detection is set empirically by the affinity of the aptamer-ligand complex, the  $K_D$  value. Using aptamer-based sensor molecules one can detect analyte binding in both solution (homogeneous) and on solid supports (heterogeneous).

Other ligand detection methods known in the art are based upon antibody binding. Similar to the aptamer-based methods, antibody-based detection requires continuous ligand binding and ligand-antibody complex formation for the generation of a detectable signal. In addition, antibody methods such as ELISA or competitive RIA, while robust, are restricted in utility because these methods require that heterogeneous assay conditions be employed:

1] detection is done on a solid surface; 2] in most applications both a capture antibody and detection antibody are required; 3] for ELISA-based protein detection methods, the antibodies must recognize the folded, native structure of the protein that is present in cell or tissue isolates and; 4] antibody and protein based detection methods have not been described for intracellular or *in vivo* based analyte detection. That antibodies have not been employed for intracellular and *in vivo* based detection of proteins, drugs or metabolites is due to several technological factors. First monoclonal antibody fragments are unstable and do not fold properly when expressed as intracellular protein molecules. Second, intracellular detection requires homogeneous assay formats and these solution-based detection methods require the sensor to have a ligand sensing or modulation domain coupled directly to a catalytic or signal generating region of an enzyme or catalytic biomolecule. A fundamental and important consequence of the limitations of antibody-based detection methods is that they can not function as a universal reagent for all assays and tests that can be employed in drug discovery and development. These assays include 1] the initial discovery of a drug target through protein or metabolite profiling, 2] the subsequent use of that same drug target in the discovery of drug leads through high throughput screening and, 3] the optimization of drug leads against that same drug target through an evaluation of lead efficacy in mechanistic cellular and *in vivo* animal assays.

To streamline the drug discovery and development process and improve the efficiency of evaluating drug targets and drug leads, detection reagents are needed that can

function in a context-independent manner. Needed then is a molecular sensor that can function in multiple assay environments and in multiple assay formats. The present invention is generally drawn to nucleic acid molecular sensors that can function in environments and formats that includes but are not limited to solution-based detection  
5 (homogeneous *in vitro* biochemical assays or *in vivo* cellular and animal assays), chip-based (heterogeneous *in vitro* assays on solid surfaces), and assays in complex biological isolates from blood plasma, cell lysates or tissue extracts.

Nucleic acid-based detection schemes have exploited the ligand-sensitive catalytic properties of some nucleic acids, *e.g.*, such as ribozymes. Ribozyme-based prototype  
10 nucleic acid sensor molecules have been designed both by engineering and by *in vitro* selection methods. Engineering methods exploit the apparently modular nature of RNA structures; these sensors couple molecular recognition to signaling by simply joining individual target-modulation and catalytic RNA domains through a double-stranded or partially double-stranded RNA linker. ATP sensors, for example, were created by  
15 appending the previously-selected, ATP-binding aptamer-derived sequences (Sassanfar and Szostak, 1993) to either the self-cleaving hammerhead ribozyme (Tang and Breaker, 1997) or the L1 self-ligating ribozyme (Robertson and Ellington 2000). Robertson and Ellington (2000) have demonstrated that the enzymatic activity of a ligase ribozyme (derived from the L1 ligase described in Robertson and Ellington (1999)) can be modulated by a small  
20 molecule ligand, or small molecule target recognition. In this case, the ligase ribozyme can be employed as a nucleic acid sensor molecule and used to detect the presence and level of its cognate ligand by monitoring the ligation of a small, labeled second oligonucleotide substrate on to the ribozyme. A distinct feature of this detection method is that the actual detection event, *e.g.*, monitoring oligonucleotide substrate ligation to the ribozyme, occurs  
25 after the ligand interacts with the nucleic sensor molecule. Hence, unlike antibody, or aptamer based detection methods, the ribozyme-based ligand detection method of Robertson and Ellington does not require continuous binding of the ligand to the sensor molecule in order to generate a detectable signal. In a complementary approach, radiolabeled hammerhead ribozymes which undergo cleavage upon binding to a ligand, have been used  
30 to detect ligand by monitoring the release of the label from the ribozyme (Soukup, et al., 2000, and Breaker, 1998). Limitations of the use of ligand modulated hammerhead ribozymes described by Soukup, et al., 2000, and Breaker, 1998 include: 1] the need for a two-step detection method for determining the enzymatic activity of the surface-immobilized hammerhead-derived sensors; 2] the need for radiometric determination of

hammerhead activity in both solution and solid-surface based assay formats; 3] the need for significant chemical and structural modification of the hammerhead-based biosensor to render them suitable for optical based detection methods.

Another limitation of the engineering method to sensor generation is that it has been  
5 generally thought not to be applicable to the development of protein-dependent ribozymes. Robertson and Ellington (2001) describe their own efforts to extend this methodology to the identification of protein and peptide-dependent ribozymes, but state that simply appending aptamer-derived sequences to the catalytic domain of the L1 ligase at stem C, yields little or no target dependent modulation. Furthermore, the authors state that the “principles required  
10 for engineering protein-dependent ribozymes must be fundamentally different from those for identifying ribozymes dependent on small-molecules.” Hence, in order to identify protein and peptide dependent ribozymes, Ellington and Robertson undertook a laborious *in vitro* selection process which involved randomization of the catalytic core of the L1 ligase coupled with multiple rounds of positive and negative selection. Ellington and Roberts  
15 (2000) describe several limitations of the ligase-derived sensors that they developed. First, the nucleoprotein enzymes developed by Ellington and Robertson (2000) required a laborious *in vitro* selection process to identify peptide and protein dependent ligases. Secondly, Ellington and Roberts (1999) describe a region of the L1 ligase that is required for allosteric ribozyme function, termed the effector oligonucleotide binding domain. It was  
20 postulated that the effector oligonucleotide binding domain of the ligase formed complementary base pairing interactions with the oligonucleotide substrate binding site, driving the ribozyme into an inactive conformation. The effector oligonucleotide, when added to the L1 ligase activates ( $k_{act}$ ) the enzyme by over 10,000 fold over the L1 ligase reaction measure in the absence of effector ( $k_{unact}$ ). Hence, the native L1-ligase has a  
25 switch factor ( $k_{act}/k_{unact}$ ) greater than 10,000, which determines the sensitivity of a ribozyme-based detection method. When the effector oligonucleotide binding domain of the L1 ligase is deleted, the ligase activity of the deletion mutant is only 3-5 fold lower than the ligase activity of L1 ligase with the effector oligonucleotide bound to the effector oligonucleotide binding domain (Ellington and Robertson (1999). This indicates that L1  
30 ligases deleted of the effector oligonucleotide binding domain may not be not subject to further allosteric regulation. Hence, a hindrance to the development of L1 ligase-based biosensor technology is the lack of a general method for the generation of biosensors that can work in multiple assay and detection formats required of solution-based and chip-based

biosensors and, those that can work in multiplexed formats and in complex biological extracts.

### Summary of the Invention

The nucleic acid sensor molecules of the present invention are used to monitor the  
5 presence or concentration of various target molecules (proteins, post-translationally  
modified forms of proteins, peptides, nucleic acids, oligosaccharides, nucleotides,  
metabolites, drugs, toxins, biohazards, and ions) and function in solution-based  
homogeneous assays using optical or other detection methods; in solution-based  
homogeneous assays using PCR-based or other nucleotide amplification-based detection  
10 methods; in homogeneous intracellular assays using PCR-based detection or other  
nucleotide amplification-based detection methods; in heterogeneous assays (surface-  
immobilized nucleic acid sensor molecules and surface-capture nucleic acid sensor  
molecules) using optical detection methods; and in heterogeneous assays (surface-  
immobilized nucleic acid sensor molecules and surface-capture nucleic acid sensor  
15 molecules) using PCR-based detection or other nucleotide amplification-based detection  
methods. And finally the nucleic acid sensor molecules of the present invention function in  
formats where the target analytes are present in complex biological mixtures, or the assays  
are themselves performed in multiplexed formats.

The nucleic acid sensor molecules of the present invention were developed through  
20 a combination of engineering and selection methods that are now shown to be useful for  
identifying nucleic acid sensor molecules against a wide variety of target molecules  
including proteins (including specific post-translationally modified forms of proteins)  
peptides, nucleic acids, oligosaccharides, nucleotides, metabolites, drugs, toxins,  
biohazards, ions, carbohydrates, glycoproteins, hormones, receptors, antibodies, viruses,  
25 transition state analogs, cofactors, dyes growth factory nutrients, etc.

In one embodiment of the invention the nucleic acid sensor molecules are based on  
cis-cleaving hammerhead ribozymes that have been designed to work as optical signaling  
molecules in a homogeneous assay format, and utilize fluorescence and FRET based  
methods of signal generation and detection.

30 In one embodiment of the invention nucleic acid sensor molecules are based on cis-  
cleaving hammerhead ribozymes that have been designed to work as optical signaling  
molecules affixed to a solid-support, and utilize fluorescence and FRET based methods of  
signal generation and detection.

In one embodiment of the invention the nucleic acid sensor molecules are based on cis-cleaving hammerhead ribozymes that have been designed to work as optical signaling molecules affixed to a solid-support and utilize surface plasmon resonance methods of signal generation and detection.

- 5 In one embodiment of the invention the nucleic acid sensor molecules are based on a 3-piece L1 ligase ribozyme that retains the effector oligonucleotide binding domain, and has been designed to detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule, in solution using quantitative PCR-based methods.

- 10 In one embodiment of the invention the nucleic acid sensor molecules are based on a 2-piece L1 ligase ribozyme that couples the effector oligonucleotide substrate to the oligonucleotide substrate forming an oligonucleotide supersubstrate and, has been designed to detect target molecules, as a function of oligonucleotide supersubstrate ligation to the nucleic acid sensor molecule, in solution using quantitative PCR-based methods.

- 15 In one embodiment of the invention the nucleic acid sensor molecules are based on a 1-piece L1 ligase ribozyme that deletes the effector oligonucleotide binding domain. The 1-piece ligase is designed to self-ligate or circularize by joining the 3'-OH end and the PPP-5'-end of the ligase and detects target molecules, as a function of circularization of the nucleic acid sensor molecule, in solution using PCR-based methods.

- 20 In one embodiment of the invention the nucleic acid sensor molecules are based on a 3-piece L1 ligase ribozyme that retains the effector oligonucleotide binding domain, and has been designed to detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule immobilized on a solid support and where the detection uses fluorescence-based methods.

- 25 In one embodiment of the invention the 3-piece L1 ligase nucleic acid sensor molecules detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule in solution and then are subsequently captured on a solid support and detected using fluorescence-based methods.

- 30 In one embodiment of the invention the 3-piece L1 ligase nucleic acid sensor molecules detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule in solution and then are subsequently captured on a solid support and detected using radiometric-based methods.

In one embodiment of the invention the nucleic acid sensor molecules are based on a 2-piece L1 ligase ribozyme and has been designed to detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule immobilized on a



solid support and where the detection uses optical methods such as fluorescence-based methods.

In one embodiment of the invention the 2-piece L1 ligase nucleic acid sensor molecules detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule in solution and then are subsequently captured on a solid support and detected directly using fluorescence-based methods.

In one embodiment of the invention the 2-piece L1 ligase nucleic acid sensor molecules detect target molecules, as a function of oligonucleotide substrate ligation to the nucleic acid sensor molecule in solution and then are subsequently captured on a solid support and directly detected using radiometric-based methods.

In one embodiment of the invention the nucleic acid sensor molecules is based on a 1-piece L1 ligase ribozyme that is transfected into a mammalian cell. The 1-piece ligase is designed to self-ligate or circularize by joining the 3'-OH end and the PPP-5'-end of the ligase and detects intracellular target molecules, as a function of circularization of the nucleic acid sensor molecule. The circularized ligase molecules are then re-isolated from the cellular lysate and the amount of target present in the cell is quantified using solution based nucleotide amplification methods.

In one aspect, the invention includes a nucleic acid sensor molecule which includes a target modulation domain which recognizes a target molecule, a linker domain, a catalytic domain, and an optical signal generating unit.

In one embodiment, the nucleic acid sensor molecule of the invention has an optical signal generating unit that includes at least one signaling moiety. In another embodiment, the nucleic acid sensor molecule of the invention has an optical signal generating unit which includes at least a first signaling moiety and a second signaling moiety. In another embodiment, the first and second signaling moieties change proximity to each other upon recognition of a target by the target modulation domain. In another embodiment, the first and second signaling moieties include a fluorescent donor and a fluorescent quencher, and recognition of a target by the target modulation domain results in an increase in detectable fluorescence of said fluorescent donor. In another embodiment, the first signaling moiety and said second signaling moiety include fluorescent energy transfer (FRET) donor and acceptor groups, and recognition of a target by the target modulation domain results in a change in distance between said donor and acceptor groups, thereby changing optical properties of said molecule.

In another embodiment, the invention includes a nucleic acid sensor molecule where the optical signal generating unit consists essentially of a first signaling moiety which changes conformation upon recognition of a target by the target modulation domain, thereby resulting in a detectable optical signal.

5 In yet another embodiment, the nucleic acid sensor molecule includes at least one modified nucleic acid.

In still another embodiment, the catalytic domain of the nucleic acid sensor molecule includes an endonucleolytic ribozyme. The endonucleolytic ribozyme can be, for example, a *cis*-endonucleolytic ribozyme or a *trans*-endonucleolytic ribozyme. In one embodiment,  
10 the endonucleolytic ribozyme is a hammerhead ribozyme.

In another embodiment, the catalytic domain of the nucleic acid sensor includes a self-ligating ribozyme. The self-ligating ribozyme can be, for example, a *cis*-ligase ribozyme or a *trans*-ligase ribozyme. The self-ligating ribozyme can be, e.g., a 1-piece ligase, 2-piece ligase or 3-piece ligase.

15 In another embodiment, the target modulation domain recognizes a target that is selected from proteins, post-translationally modified forms of proteins, peptides, nucleic acids, oligosaccharides, nucleotides, metabolites, drugs, toxins, biohazards, ions, carbohydrates, polysaccharides, hormones, receptors, antigens, antibodies, viruses, metabolites, co-factors, drugs, dyes, nutrients, or growth factors.

20 In one embodiment, the target modulation domain recognizes a protein or a post-translationally modified protein.

In another embodiment, the target modulation domain recognizes a post-translationally modified protein, wherein the post-translational modification can be a phosphorylation, prenylation, glycosylation, methionine removal, N-acetylation, acylation, acylation of cysteines, myristoylation, alkylation, ubiquitinylation, prolyl-4-hydroxylation,  
25 carboxylation of glutaminy residues, advanced glycosylation, deamination of glutamine and asparagine, addition of glycoposphatidylinositol, disulfide bond formation, hydroxylation, or lipidation.

In some embodiments, the target is a protein kinase. In other embodiments, the  
30 target is a phosphorylated protein kinase. The phosphorylated protein kinase can be a monophosphorylated protein kinase or a diphosphorylated protein kinase. In one embodiment, the target protein is ERK, such as ERK1 or ERK2. In other embodiments, the post-translationally modified protein is ppERK, such as ppERK1 or ppERK2..

In other embodiments, the nucleic acid sensor molecule includes a target modulation domain which recognizes a component of a MAP kinase pathway, a product of a MAP kinase pathway, a MAP kinase pathway associated protein, or an extracellular component of a MAP kinase pathway. In one embodiment, the target modulation domain recognizes a component of the ERK1/2 MAP kinase pathway, the JNK MAP kinase pathway, or the P38 MAP kinase pathway.

In some embodiments, the target modulation domain recognizes an endogenous form of a MAP Kinase (MEK), an endogenous form of a MAP Kinase Kinase (MEKK), or an endogenous form of MAP Kinase Kinase Kinase, (MEKKK). In other embodiments, the target modulation domain recognizes an endogenous form of RAF kinase. In still other embodiments, the target modulation domain recognizes a Ras protein, a phosphatase, a GTP binding protein, a GPCR, a cytokine, a growth factor, a cellular metabolite, or a small molecule.

In some embodiments, the nucleic acid sensor molecule includes RNA, DNA, or both RNA and DNA.

The invention also relates to compositions containing a nucleic acid sensor molecule of the invention and a buffer. In some embodiments, the invention includes a composition containing a nucleic acid sensor molecule and a tissue extract, a cell extract or an *in vitro* cell culture. In other embodiments, a composition of the invention also includes an RNase inhibitor, such as, for example, Va-riboside, vanadyl, tRNA, polyU, RNaseIn or RNaseOut. In some embodiments, a composition of the invention is substantially RNase-free.

The invention also relates to a composition which includes at least one nucleic acid sensor molecule affixed to a substrate. In some embodiments, the substrate is glass, gold or other metal, silicon or other semiconductor material, nitrocellulose, nylon, or plastic. In particular embodiments, the nucleic acid sensor molecule is covalently attached to the substrate. In other embodiments, the nucleic acid sensor molecule is non-covalently attached to the substrate. In some embodiments, the nucleic acid sensor molecule is immobilized to the substrate *via* hybridization of a terminal portion of the nucleic acid sensor molecule to an oligonucleotide that is bound to the surface of the substrate.

In some embodiments, a composition includes a plurality of nucleic acid sensor molecules immobilized to the substrate *via* hybridization of a terminal portion of the nucleic acid sensor molecule to an array of oligonucleotides bound to the substrate at spatially discrete regions. In some embodiments, at least two members of this plurality each recognize different target molecules. The substrate can include, for example, at least 50

nucleic acid sensor molecules. In other embodiments, the substrate includes at least 250 nucleic acid sensor molecules, at least 500 nucleic acid sensor molecules, or at least 5000 nucleic acid sensor molecules.

5 In some aspects, the invention includes a system for detecting a target molecule which includes a composition according to the invention and a detector in optical communication with the composition, where the detector detects changes in the optical properties of the composition. In some embodiments, the system further includes a light source in optical communication with the composition. In some embodiments, the system also includes a processor for processing optical signals detected by the detector.

10 In one embodiment, the system for detecting a target molecule includes a plurality of nucleic acid sensor molecules where at least two of the biosensor molecules each recognize different target molecules.

In another aspect, the invention includes a method of identifying or detecting a target molecule in a sample by contacting a sample suspected of containing a target  
15 molecule with a nucleic acid sensor molecule of the invention, wherein a change in the signal generated by the optical signal generating unit indicates the presence of target in the sample. In some embodiments, the method further includes quantifying the change signal generated by the optical signal generating unit to quantify the amount of target molecule in the sample. In some embodiments, the sample is an environmental sample, biohazard  
20 materials, organic samples, drugs and toxins, flavors and fragrances, and biological samples. In particular embodiments, the sample is a biological sample such as cells, cell extracts or lysates, tissues or tissue extracts, bodily fluids, serum, blood and blood products.

In some embodiments, the method of identifying or detecting a target molecule in a sample detects or identifies proteins, post-translationally modified forms of proteins,  
25 peptides, nucleic acids, oligosaccharides, nucleotides, metabolites, drugs, toxins, biohazards, ions, carbohydrates, polysaccharides, hormones, receptors, antigens, antibodies, viruses, metabolites, co-factors, drugs, dyes, nutrients, or growth factors. In other embodiments, the method of the invention detects or identifies proteins or post-translationally modified forms of proteins. In some embodiments, the target is a post-  
30 translationally modified protein, where the post-translation modifications is phosphorylation, prenylation, glycosylation, methionine removal, N-acetylation, acylation, acylation of cysteines, myristoylation, alkylation, ubiquitinylation, prolyl-4-hydroxylation, carboxylation of glutaminyl residues, advanced glycosylation, deamination of glutamine

and asparagine, addition of glycoposphatidylinositol, disulfide bond formation, hydroxylation, or lipidation.

In some embodiments, the target is a protein kinase. In particular embodiments, the target is a phosphorylated protein kinase.

5 In another aspect, the invention includes a diagnostic system for identifying or detecting a target molecule, where the diagnostic system includes a nucleic acid sensor molecule of the invention and a detector in communication with the nucleic acid sensor molecule, wherein the detector detects changes in the signal generated by the optical signal  
10 further includes a processor for processing signals detected by the detector.

In another aspect, the invention includes a method of identifying or detecting a protein kinase in a sample by contacting a sample suspected of containing a protein kinase with a nucleic acid sensor molecule, wherein said nucleic acid sensor molecule has a target  
15 recognition domain that recognizes a protein kinase, and wherein a change in the signal generated by the optical signal generating unit indicates the presence of protein kinase in the sample. In some embodiments, the method further includes quantifying the amount of signal generated by the optical signal generating unit to quantify the amount of protein kinase in the sample.

In yet another aspect, the invention provides a method of identifying a modulator of protein kinase activity by contacting a test agent with a protein kinase and nucleic acid  
20 sensor molecule, wherein the nucleic acid sensor molecule has a target recognition domain that recognizes a protein kinase, recognition of the protein kinase by the target recognition domain of the nucleic acid sensor molecule results in a change in the signal generated by the optical signal generating unit and wherein changes in the signal generated by the optical  
25 signal generating unit in the presence and absence of the test agent indicates the test agent is a modulator of the protein kinase activity.

In some embodiments, the catalytic domain of the nucleic acid sensor molecule includes a *cis*-ligase ribozyme or a *trans*-ligase ribozyme.

In another aspect, the invention provides a nucleic acid sensor molecule which  
30 includes a target modulation domain that recognizes ERK, a catalytic domain that includes a ligase or cis-hammerhead, and a linker domain that links the target modulation domain and the catalytic domain.

In another aspect, the invention includes a nucleic acid sensor molecule which includes a target modulation domain that recognizes phosphoERK, a catalytic domain that

includes a ligase or a cis-hammerhead, and a linker domain that links the target modulation domain and the catalytic domain.

In yet another aspect, the invention provides a nucleic acid sensor molecule which includes a target modulation domain that recognizes lysozyme, a catalytic domain that  
5 includes a 1-piece cis-ligase and a linker domain that links the target modulation domain and the catalytic domain.

In still another aspect, the invention provides a nucleic acid sensor molecule which includes a target modulation domain that recognizes any one of cCMP, cAMP, or cGMP, a catalytic domain, and a linker domain that links the target modulation domain and the  
10 catalytic domain, wherein the nucleic acid sensor molecule includes an optical signal generating unit or a non-radioactive detectable label. In some embodiments, the nucleic acid sensor molecule includes an optical signal generating unit. In other embodiments, the nucleic acid sensor molecule includes a detectable label. In a particular embodiment, the label is a radioactive label, such as, for example,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{14}\text{C}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ , or  $^{125}\text{I}$ . In other  
15 embodiments, the nucleic acid sensor molecule comprises a fluorescent label, such as, for example, fluorescein, DABCYL, or a green fluorescent protein (GFP) moiety. In some embodiments, the optical signal generating unit includes a fluorescent moiety and a quenching moiety, wherein recognition by the target modulation domain causes a change in detectable fluorescence by the optical signal generating unit. In some  
20 embodiments, the nucleic acid sensor molecule includes an enzymatic label, or an affinity capture tag label.

In some embodiments, the nucleic acid sensor molecule includes a target modulation domain recognizes ERK1, ERK2 or both. In a particular embodiment, the nucleic acid sensor molecule includes a target modulation domain and catalytic domain are as shown by  
25 SEQ ID NO. 80, and the linker is randomized.

In some embodiments, the nucleic acid sensor molecule includes a target modulation domain and a catalytic domain as shown in any one of SEQ ID NOS. 47, 118 and 119, and the linker is randomized.

In another aspect, the invention provides a nucleic acid sensor molecule that  
30 recognizes ERK having the SEQ ID NO. 90-95, 108-116, 131-133, 140-295, 349, 351, or 356.

In another aspect, the invention provides a nucleic acid sensor molecule that recognizes phosphoERK having the SEQ ID NO. 5-8, 37-39, 44-45, 81-89, 96-100, 121-130, 352, or 353.

In another aspect, the invention provides a nucleic acid sensor molecule that recognizes any one of cCMP, cAMP or cGMP, having the SEQ ID NO. 40-43, 103, or 135-139.

In another aspect, the invention provides a nucleic acid sensor molecule that  
5 recognizes lysozyme, having the SEQ ID NO. 46, 47, 76, or 105-107.

In another aspect, the invention provides a 1-piece ligase ribozyme including a target modulation domain that recognizes a target, a linker domain, and a catalytic domain wherein the 5' and 3' ends of the ligase ligate to each other upon recognition of the target by the modulation domain.

10 In another aspect, the invention provides a 2-piece ligase ribozyme including a target modulation domain that recognizes a target, a linker domain, and a catalytic domain including an oligonucleotide substrate ligation site and an oligonucleotide supersubstrate binding domain, wherein upon recognition of the target by the modulation domain, the 3' end of the an oligonucleotide supersubstrate is ligated to the 5' end of the oligonucleotide  
15 substrate ligation site.

In yet another aspect, the invention provides a 3-piece ligase ribozyme including a target modulation domain that recognizes a target, a linker domain, and a catalytic domain comprising comprising an oligonucleotide substrate binding domain capable of binding an oligonucleotide substrate and an effector-oligonucleotide binding site capable of binding an  
20 effector oligonucleotide, wherein upon recognition of the target by the modulation domain, and in the presence of binding of the effector oligonucleotide to the effector-oligonucleotide binding site, then the 3' end of the oligonucleotide substrate is ligated to the 5' end of the ligase.

The invention also provides a 1-piece ligase ribozyme comprising the nucleic acid  
25 sensor molecule shown in any one of SEQ ID NOS. 47, 105-107, 119.

The invention also provides a 2-piece ligase ribozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 347, 349, and 351.

The invention further provides a 3-piece ligase ribozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 46, 75, 76, 108-116, 118, 121-130, and  
30 352.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and

materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

5 Other features and advantages of the invention will be apparent from the following detailed description and claims.

### **Brief Description of the Drawings**

10 Figure 1 depicts methodology for selecting nucleic acid sensor molecules of the invention. Figure 1A is a flow diagram showing a method for selecting nucleic acid sensor precursor molecules having a target molecule activatable ligase activity according to one embodiment. Figure 1B is a flow diagram showing a gel-based method for selecting nucleic acid sensor precursor molecules having a target molecule activatable endonuclease activity  
15 according to one embodiment.

Figures 2A and B show a nucleic acid sensor molecule (SEQ ID NO:46) according to one embodiment, in which the catalytic site includes a ligase site. Figure 2A shows the conformation of the target molecule (SEQ ID NO:46) bound form of the nucleic acid sensor molecule with an effector oligo hybridized to its 3' end (SEQ ID NO:51). Figure 2B shows  
20 the conformation of the non-target bound form of the nucleic acid (SEQ ID NO:46) sensor molecule.

Figures 3A and B show a nucleic acid sensor molecule (SEQ ID NO:47) derived from the nucleic acid sensor precursor molecule shown in Figures 2A and B in which first and second nucleotides are labeled with first and second signaling moieties (F and D,  
25 respectively).

Figure 4 is a flow diagram showing a method for selecting nucleic acid sensor molecules having a target molecule activatable self-cleavage activity according to one embodiment.

Figures 5A and B show a nucleic acid sensor molecule according to one  
30 embodiment, in which the catalytic site includes a self-cleavage site which is the catalytic core of a hammerhead ribozyme. Figure 5A shows the conformation of the target molecule bound form of the nucleic acid sensor molecule (SEQ ID NO:48). Figure 5B shows the conformation of the non-target bound form of the nucleic acid sensor molecule (SEQ ID NO:48).



Figures 6A and B show a nucleic acid sensor molecule derived from the nucleic acid sensor molecule shown in Figures 3A and B in which first and second nucleotides are labeled with first and second signaling moieties (F and D, respectively) (SEQ ID NO:49).

Figure 7 is a schematic diagram illustrating pathway target molecules according to one embodiment.

Figure 8 is a flow chart showing steps in a drug optimization method according to one embodiment, in which nucleic acid sensor molecules are used at each step in the method.

Figure 9A shows an example of a self-cleaving nucleic acid sensor bound to a solid support when used in an epi-illuminated FRET detection scheme. Figure 9B shows the same sensor in an epi-illuminated beacon configuration, with the acceptor fluorophore replaced by a quencher group. Figure 9C shows the same sensor in an TIR-illuminated beacon configuration.

Figure 10A shows an example of a self-ligating nucleic acid sensor bound to a solid support when used in a TIR-illuminated detection scheme where there is a signal increase upon target binding. Figure 10B shows the same sensor in an epi-illuminated configuration, where target binding is detected by monitoring changes of the fluorophore bound to the substrate at the surface of the array. Figure 10C shows the same epi-illuminated configuration, where target binding is detected by monitoring changes in the fluorescence polarization.

Figure 11 illustrates the use of beads in a homogeneous assay format utilizing a self-ligating nucleic acid sensor. Figure 11A shows the beads prior to target binding and ligation (no emission from acceptor). Figure 11B shows the beads after target binding and ligation (emission from acceptor detected).

Figure 12 shows a schematic of a previously constructed scanning detection system utilizing TIR laser evanescent wave excitation in either large area illumination/CCD imaging mode, or scanned spot/PMT imaging mode. The schematic shows how an array can be scanned and FP or fluorescence intensity data extracted.

Figure 13. shows a schematic of fluorescence data generated by a biosensor array using the indicated nucleic acid sensors and target molecules.

Figures 14 shows a schematic showing different catalytic platforms for detection methods for nucleic acid sensor molecules.

Figure 15 is a schematic showing a ligase nucleic acid sensor molecule sensor system. It shows SEQ ID NO:76 across the top, and SEQ ID NO:77 hybridized to a portion of SEQ ID NO:76 and bound to an insulating moiety.

Figure 16 is a schematic showing a hammerhead (endonuclease) Nucleic acid sensor molecule sensor system. It has a nucleic acid sensor molecule sensor (SEQ ID NO:78), and SEQ ID NO:79 hybridizing to a portion of the sensor and bound to an insulating moiety.

Figure 17 is a schematic of the net electron transfer to or from the electrode.

Figure 18 is a schematic of a peak in the faradaic current, centered at the redox potential of the electron donor species (specified for a given reference electrode) and superposed on top of the capacitive current baseline which is observed in the absence of surface-immobilized signaling probes.

Figure 19 is the sequence of the entire ERK2 activated allosteric ribozyme (SEQ ID NO:80). Also shown are the sequences of the Stem II connector domain for selective clones.

Figure 20A is a chart showing measurement of cis-hammerhead cleavage. Figure 20B shows a chart showing measurement of cis-hammerhead cleavage.

Figure 20C is a chart showing measurement of cis-hammerhead cleavage.

Figure 21 is a chart showing ERK2-dependence of cis-hammerhead cleavage.

Figure 22A is a chart showing the measurement of ERK2-inhibitor IC50 values by nucleic acid sensor molecule.

Figure 22B is a chart showing the fraction of construct I-14 cleaved in the presence of 100 nM ERK2, ricin, or MEK, or with no protein.

Figure 23A shows a ppERK cis-hammerhead nucleic acid sensor molecule construct (SEQ ID NO:353).

Figure 23B shows construct 6 (SEQ ID NO:81) 7 (SEQ ID NO:82), 8 (SEQ ID NO:83), 9 (SEQ ID NO:84), 10 (SEQ ID NO:85), 11 (SEQ ID NO:86) 12 (SEQ ID NO:87), 13 (SEQ ID NO:88), and 14 (SEQ ID NO:89).

Figure 24 shows the linker region, activity, and stability of constructs 6 through 14.

Figures 25A and B show a bar graph and corresponding radiograph demonstrating the relative pp ERK dependence of constructs 6, 10, and 12.

Figure 26 shows the sequences for lysozyme modulated ligase nucleic acid sensor molecules C.lys.L1.A (SEQ ID NO:105), C.lys.L1.B (SEQ ID NO:106), and C.lys.L1.C (SEQ ID NO:107).

Figure 27 is a schematic of how an L1 ligase is configured for self-circularization, and how its self-circularization can be detected using RT-PCR.

Figure 28 shows increases in amplification of circularized C.lys.L1.B (SEQ ID NO:106) in response to the addition of lysozyme. The signal is strengthened as additional  
5 cycles of PCR are performed.

Figure 29 shows that the ligase nucleic acid sensor molecule C.lys.L1.B (SEQ ID NO:106) still self ligates in response to the presence of lysozyme, even in the presence of HeLa cell extract, demonstrating the stability of this nucleic acid sensor molecule.

Figure 30 shows modulation of a 1-piece ligase nucleic acid sensor molecule *in vivo*.

10 Figure 31 is a schematic showing the construct design for an ERK dependent 3-piece ligase nucleic acid sensor molecule (SEQ ID NO:118).

Figure 32 shows the sequences for constructs 17 (SEQ ID NO:109), 18 (SEQ ID NO:110), 19 (SEQ ID NO:111), 20 (SEQ ID NO:112), and 21 (SEQ ID NO:113), 22 (SEQ ID NO:114), 23 (SEQ ID NO:115), 24 (SEQ ID NO:116), 25 (SEQ ID NO:117), and 26  
15 (SEQ ID NO:118).

Figure 33 shows the ERK dependent activity of constructs 17 (A), 18 (B), 20 (C), 21 (D), 19 (E), 22 (F), 23 (G), 25 (H), and 26 (I).

Figure 34 is a graph that shows ligase time-dependent activity assays for construct 17 (clone A) (SEQ ID NO:109) and construct 19 (clone E) (SEQ ID NO:111).

20 Figure 35 shows a graph showing the time-dependent activity of construct 19 (clone E) (SEQ ID NO:111) with varying concentrations of ERK2.

Figure 36 shows secondary structure representations of 3-piece construct 27 (SEQ ID NO:118) and 1-piece construct 28 (SEQ ID NO:119) ERK dependent ligases. 1-piece ERK dependent ligase is a slightly modified version of 3-piece system where the effector  
25 and substrate regions are replaced by a stable GNRA tetraloop.

Figure 37 shows a graph demonstrating continued ERK2 dependence of a nucleic acid sensor molecule in the 3-piece and 1-piece formats (constructs 19 and 28, respectively).

Figure 38 shows a secondary structure representation of the 2-piece ERK dependent ligase platform (SEQ ID NO:347), and its oligonucleotide substrate (SEQ ID NO:350).

30 Figure 39A shows a secondary structure representation of two 2-piece ERK-modulated nucleic acid sensor molecules: construct 29 (SEQ ID NO:349), and construct 30 (SEQ ID NO:351), with their oligonucleotide substrate (SEQ ID NO:348).

Figure 39B shows ligation assays run on constructs 29 (SEQ ID NO:349) and 30 (SEQ ID NO:351), in the absence or in the presence of 1  $\mu$ M ERK.

Figure 40 shows the ligation efficiency of ERK nucleic acid sensor molecule construct 19 (SEQ ID NO:111) detected with quantitative-PCR (Taqman<sup>®</sup>). All incubation with various concentration of ERK were performed in the presence of 10% 293 extracts with exogenously added 10 mM MgCl<sub>2</sub>.

5        Figure 41 shows a schematic of the template for ppERK dependent ligase nucleic acid sensor molecules (SEQ ID NO:352) and its oligonucleotide substrate (SEQ ID NO:350).

10        Figure 42 shows the nucleotide sequences for construct 31 (TK.16.118.K) (SEQ ID NO:121), construct 32 (TK.16.118.L) (SEQ ID NO:122), construct 33 (TK.16.118.M) (SEQ ID NO:123), construct 34 (TK.16.118.N) (SEQ ID NO:124), construct 35 (TK.16.118.O) (SEQ ID NO:125), construct 36 (TK.16.118.P) (SEQ ID NO:126), construct 37 (TK.16.118.Q) (SEQ ID NO:127), construct 38 (TK.16.118.R) (SEQ ID NO:128), construct 39 (TK.16.118.S) (SEQ ID NO:129), and construct 40 (TK.16.118.T) (SEQ ID NO:130).

15        Figure 43 shows template sequences for the creation of a ppERK (SEQ ID NO:354) or ERK (SEQ ID NO:355) library of nucleic acid sensor molecules.

20        Figure 44A shows the stem sequences of ERK dependent nucleic acid clones CW45-33-A08 (SEQ ID NO:356), CW45-33-C08 (SEQ ID NO:131), CW45-33-C09 (SEQ ID NO:132), CW45-33-D09 (SEQ ID NO:133), CW45-33-F08 (SEQ ID NO:90), CW45-33-H08 (SEQ ID NO:91), CW45-33-H09 (SEQ ID NO:92), CW45-33-A10 (SEQ ID NO:93), CW45-33-F09 (SEQ ID NO:94), and CW45-33-G08 (SEQ ID NO:95).

25        Figure 44B shows the stem sequences of pp ERK dependent nucleic acid clones CW45-33-A02 (SEQ ID NO:44), CW45-33-B04 (SEQ ID NO:45), CW45-33-C04 (SEQ ID NO:5), CW45-33-D04 (SEQ ID NO:6), CW45-33-F03 (SEQ ID NO:7), CW45-33-D01 (SEQ ID NO:8), CW45-33-D02 (SEQ ID NO:37), CW45-33-D05 (SEQ ID NO:38), CW45-33-E01 (SEQ ID NO:39), CW45-33-G02 (SEQ ID NO:96), CW45-33-G03 (SEQ ID NO:97), CW45-33-H03 (SEQ ID NO:98), CW45-33-H1 (SEQ ID NO:99), and CW45-33-B05 (SEQ ID NO:100).

30        Figure 45 nucleotide sequences of CW45-33-A02 (SEQ ID NO:44), and CW45-33-D04 (SEQ ID NO:6).

      Figure 46 shows a schematic demonstrating amplicon-dependent nucleic acid amplification (ADNA).

      Figure 47 shows a schematic describing a mechanism used by nucleic acid sensor molecules to transduce signal and the kinetic constants used to characterize NASMs.

Figure 48 shows a graph demonstrating the determination of threshold cycle versus log of target molecule concentration using amplicon-dependent nucleic acid amplification via quantitative PCR analysis.

5 Figure 49 shows a graph demonstrating the determination of threshold cycle versus log of target molecule concentration using amplicon-dependent nucleic acid amplification via SYBR-green analysis.

Figure 50 shows a radiograph demonstrating lysozyme sensitive ligase nucleic acid sensor molecule activity in reticulocyte and HeLa cell extract.

10 Figure 51 shows a radiograph showing that ligase activity is relatively unchanged in the presence of cell lysate and various RNase inhibitors.

Figure 52 shows a radiograph of a lysozyme modulated nucleic acid sensor molecule in the presence of human serum.

Figure 53 shows a schematic describing rolling circle amplification of an amplicon derived from immobilized trans-acting ligase nucleic acid sensor molecules.

15 Figure 54 shows a schematic describing exponential amplification of an amplicon.

Figure 55 shows a schematic describing FRET-based signal generation coupled to nucleic acid synthesis of nucleic acid sensor molecules.

Figure 56 shows a schematic describing cellular assays using 1-piece ligase nucleic acid sensor molecules.

20 Figure 57 shows bar graphs plotting the rate of activity of nucleic acid sensors when in the presence of different target molecules *in vitro*. Panel A shows the rate of activity in the presence of ERK and phosphorylated ERK for construct 19 (ligase E) on the left bar and construct 33 (ligase M) on the right bar. Panel B shows construct 19 and 33 rates of activity in the presence of Ras, MEK, ERK, p38, and ricin.

25 Figure 58 shows a graph describing the activity of an ERK modulated nucleic acid sensor molecule in the presence of 10% 293 cell extract in the left panel. The right panel shows the activity of an ERK modulated nucleic acid sensor molecule in the presence of increasing concentrations of staurosporine. Both panels show data determined using quantitative PCR methods (ADNA).

30 Figure 59 lists the switch factor, dissociation constants, catalytic constant and detection limit for an ERK aptamer in comparison to four ERK dependent ligase nucleic acid sensor molecules.

Figure 60A and B show RT-PCR gels, and C and D corresponding bar graphs showing ERK modulation of a nucleic acid sensor molecule *in vitro* (panels A and C) and in biological extracts (panels B and D).

Figure 61 shows a schematic describing optical detection based on the modulation of an intron-derived nucleic acid sensor molecule.

Figure 62 shows the original solution-phase cGMP-dependent hammerhead nucleic acid sensor molecule FRET construct (SEQ ID NO:101) and its effector/capture oligo (SEQ ID NO:102) from which the solid-phase FRET sensor was derived. In the solution-phase construct shown in Figure 62A, the fluorophore (F) and quencher (Q) are FAM and DABCYL, respectively. In the solid-phase construct shown in Figure 62B, the donor fluorophore (D) and acceptor fluorophore (A) are FAM and AlexaFluor 568, respectively.

Figure 63A shows the surface-immobilized FRET sensor before, and Figure 63B shows after, exposure to the activating target molecule (cGMP), followed by subsequent cleavage and dissociation of the sequence fragment containing the acceptor fluorophore (A). Figure 63C shows the expected kinetic time course signals and Figure 63D shows the actual kinetic time course signals observed from these sensors in the presence of various concentrations of target.

Figure 64 shows fitted kinetic time course signals observed from the solid-phase FRET sensor constructs in a solution-phase assay. Figure 64A shows a graph that plots the signal observed from the donor fluorophore only in the presence of 200  $\mu$ M cGMP. Figure 64B shows a graph of the parametric fit to the experimental data shown in Figure 64A, verifying that the rate constant for the solid-phase construct is in fact similar to that for the solution-phase construct under similar conditions.

Figures 65A and 65B compare the observed pseudo-first order rate constants from solution- and solid-phase FRET sensor constructs. Figure 65 C, D, and E shows experimental data and constructs for multiplexed detection using solution-phase cGMP and cAMP FRET.

Figure 66A shows an endonuclease (hammerhead ribozyme)-based nucleic acid sensor immobilized linked to a gold surface via a thiol linker. Figure 66B shows the fraction of this type of sensor cleaved and dissociated as a function of time in the presence of a fixed concentration of target. Figure 66C shows the signal (image density) from a panel of immobilized sensors prior to their exposure to a target-mixture. Figure 66D shows the signal from the uncleaved sensors after exposure to the mixture of all listed targets, while

Figure 66E represents the target-dependent cleavage signal. Specific target-dependent activity of each sensor is seen in each case for this multiplexed assay.

Figure 67 shows a schematic diagram of the integrated SPReeta SPR sensor module (Figure 67A), as well as the nucleic acid (hammerhead ribozyme) sensor molecule that is  
5 immobilized on the gold SPR layer (Figure 67B). Figures 67C and 67D show typical real-time data generated by the SPR sensor system during sensor loading and target analyte-induced cleavage, respectively.

Figure 68 gives the sequences for three cyclic nucleotide-dependent nucleic acid (hammerhead ribozyme) sensors dependent upon cGMP (SEQ ID NO:135), cCMP (SEQ ID  
10 NO:136), and cAMP (SEQ ID NO:137) in a conformation suitable for direct 5' surface attachment. The schematic shows the SPR sensor construct intended for direct 5' attachment to a native gold surface via a terminal thiol linker.

Figure 69 gives the sequences for three cyclic nucleotide-dependent nucleic acid (hammerhead ribozyme) sensors dependent upon cCMP (SEQ ID NO:138), cAMP (SEQ ID  
15 NO:103), and cGMP (SEQ ID NO:139) in a conformation suitable for direct 3' surface attachment. The figure schematic shows the SPR sensor construct intended for direct 3' attachment to a neutravidin surface which has been passively adsorbed onto the gold SPR surface via cysteine residues.

Figure 70 gives the sequences for three cyclic nucleotide-dependent nucleic acid (hammerhead ribozyme) sensors dependent upon cCMP (SEQ ID NO:40), cAMP (SEQ ID  
20 NO:41), and cGMP (SEQ ID NO:42) in a conformation suitable for indirect surface attachment via a capture oligo. The schematic shows the SPR sensor construct intended for indirect surface attachment via a capture oligo to a neutravidin surface which has been passively adsorbed onto the gold SPR surface via cysteine residues.

Figure 71 shows the surface loading (Figure 71A) and target-dependent cleavage (Figure 71B) of cGMP-dependent nucleic acid (hammerhead ribozyme) sensor molecules, as well as the physical configurations of the various nucleic acid sensor molecules (Figure  
25 71C) that give rise to the indicated portions of the kinetic data shown. The high signal to noise ratio (SNR) and dynamic range (DNR) of this SPR sensor array are clearly evident  
30 from the plots.

Figure 72 shows a plot of SPR signal (in refractive index units, RIU) vs. time for a typical SPR sensor array assay: surface cleaning (dH<sub>2</sub>O, NaOH, PBS), surface loading of the gold SPR layer with neutravidin (NA), equilibration with PBS (PBS), loading of the

surface with biotinylated sensor molecules in PBS(HH+B/PBS), requilibration of sensor surface in assay buffer (HH buff), and addition of target in assay buffer (target/HH buff).

Figure 73 shows a schematic representation of the secondary structure and components of a ligase-based nucleic acid array sensor. The sensor (SEQ ID NO:75) is shown attached to the chip surface via hybridization to a capture oligo (SEQ ID NO:104), and with an external substrate oligo bearing a fluorescent label already ligated into place. The substrate oligo can be either directly labeled (as shown), or labeled with an affinity tag (e.g., biotin) for subsequent indirect labeling or signal amplification (e.g., via tyramide signal amplification).

Figure 74 contrasts the two principal solid-phase array (chip) formats used for ligase-based nucleic acid sensors.

Figure 75 shows a multiplex in situ ligase sensor chip, with pre-immobilized radiolabeled sensors activatable by lysozyme (LYS) and FMN.

Figure 76 shows dose-response data for ERK-dependent ligase-based nucleic acid sensors using a gel-assay (panel A) and a capture chip (pane IB).

Figure 77 shows dose-response data for an in situ ligase-based nucleic acid sensor array populated with ERK-dependent unlabeled ligase sensors in Figure 77A and in Figure 77B the retained ligation signal from each spot in the concentration profile was plotted vs. its corresponding target concentration.

Figure 78 shows dose-response data for a ERK-dependent ligase-based nucleic acid sensor capture array in Figure 78A and captured and amplified fluorescent ligation signal for each spot is plotted in Figure 78B vs. its corresponding target concentration.

Figure 79 shows the components of a generalized construct for an amplifiable ligase-based nucleic acid sensor molecule.

Figure 80 shows a generalized strategy for performing a multiplexed capture chip formatted assay with ligase-based nucleic acid sensors.

Figure 81 shows a generalized strategy for performing a highly sensitive capture chip formatted assay with ligase-based nucleic acid sensors.

Figure 82 shows a schematic describing multiplexed chip assays.

### Detailed Description of the Invention

The invention is generally drawn to catalytic NASMs (also know as allosteric ribozymes, aptazymes and the like) and optical nucleic acid sensor molecules that may be used to monitor the presence or concentration of various target molecules. Target



molecules include a variety of biologically relevant molecules, such as, for example, proteins (including specific post-translationally modified forms of proteins), peptides, nucleic acids, nucleotides, natural products, metabolites, drugs, toxins, biohazards, and ions.

The invention also includes methods by which a change in the activity or  
5 conformation of a nucleic acid sensor molecule upon recognition of a specific target molecule can be coupled to a quantifiable, measurable signal. The invention also includes methods which allow one to test the inhibitory activity of one or more compounds simultaneously against one or more enzymes or biochemical targets. Assays can be carried out in a variety of formats, including *in vitro* biochemical assays, *in vitro* cellular assays, *in vivo* cellular assays, in solution, on chips or other substrates, or *in vivo* animal models.  
10 These assays have applications in all phases of drug discovery, including target validation and discovery and development, high throughput screening, biochemical assays, *in vitro* cellular models and *in vivo* animal models.

Nucleic acid sensor molecules are RNAs, DNAs, RNA/DNA hybrids, or derivatives  
15 or analogs of nucleic acids that catalyze a chemical reaction and/or undergo a conformational change upon the recognition of a specific target molecule. Nucleic acid sensor molecule - based assays can be carried out using all catalytic platforms, which include endonucleases, such as the hammerhead ribozyme, the hairpin ribozyme, the HDV ribozyme, and the VS ribozyme; ligases, such as the L1 ligase, and the class I-III ligases  
20 and; group I and group II self-splicing introns.

Catalytic NASMs can be generated or selected by a variety of methods both disclosed herein and known in the art. For examples, see WO98/27104, WO01/96559, and WO 00/26226

Also disclosed herein are optical nucleic acid sensor molecules and methods making  
25 them. In general, optical catalytic NASMs generate a detectable optical signal upon recognition of a target molecule. Optical NASMs are generated from catalytic NASMs by addition of an optical signal generating unit.

Also disclosed herein are two new classes of catalytic NASMs (referred to as one-piece or cis ligases) and two piece ligases and methods of generating them.

30 In general, catalytic NASMs can be used, *e.g.*, to detect target molecules either by generation of an optical signal or an amplicon detectable, *e.g.*, by RT-PCR, size gel purification procedures and any other means of separating variously sized or conformed nucleic acid molecules. Optical NASMs, on the other hand, can be used, *e.g.*, to detect target molecules by generation of an optical signal.

Optical signals can be generated by optical NASMs in a number of ways. In some embodiments the signal is an optical signal generated, *e.g.*, by the fluorescence of a fluorescent dye. In other embodiments, the signal is an optical signal generated by molecules in close proximity to the nucleic acid sensor molecule whose optical or electrochemical properties are affected by the presence of the target molecule bound nucleic acid sensor molecule. In some embodiments the nucleic acid sensor molecules comprise at least one signaling moiety. In other embodiments, the nucleic acid sensor molecules comprise first and second signaling moieties whose optical properties change in response to the binding of a target molecule through changes in the proximity of the first and second signaling moieties. Thus, detection can be direct or indirect.

In one embodiment, a plurality of nucleic acid sensor molecules are provided, either in solution, or immobilized on a substrate, generating a biosensor. In a further embodiment, a diagnostic system is provided which comprises at least one biosensor in optical communication with a optical signal detector. Methods of using the diagnostic system are also provided, as well as kits for performing the method.

In other embodiments, the NASMs are used to detect the presence of target molecules *in vivo*.

In order to more clearly and concisely describe and point out the subject matter of the claimed invention, the following definitions are provided for specific terms which are used in the following written description and the appended claims.

As defined herein, a “oligonucleotide” is used interchangeably with the term “nucleic acid” and includes RNA or DNA (or RNA/DNA) sequences of more than one nucleotide in either single strand or double-stranded form. A “modified oligonucleotide” includes at least one residue with any of: an altered internucleotide linkage(s), altered sugar(s), altered base(s), or combinations thereof.

As defined herein, a “target molecule” is any molecule to be detected. The term “target molecule” refers to, any molecule for which nucleic acid sensor molecule exists or can be generated and can be naturally occurring or artificially created.

As defined herein, a “signature target molecule” is a target molecule whose expression is correlatable with a trait.

As used herein, a “diagnostic signature target molecule” is a signature target molecule whose expression is, by itself or in combination with other signature target molecules, diagnostic of a trait.

As used herein, a **“pathway target molecule”** is a target molecule involved in a biological or metabolic pathway and whose accumulation and/or activity is dependent on other target molecules in the same biological or metabolic pathway, or whose accumulation and/or activity affects the accumulation and/or activity of other target molecules in the same biological or metabolic pathway.

As used herein, a **“diagnostic pathway target molecule”** is a pathway target molecule whose expression/activity and/or structural properties, by itself or in combination with other pathway target molecules, is diagnostic of a particular trait.

As used herein, a **“profiling nucleic acid sensor molecule”** is a nucleic acid sensor molecule that recognizes a signature target molecule, a diagnostic signature target molecule, a pathway target molecule, and/or a diagnostic pathway target molecule.

As defined herein, a **“biosensor”** comprises a plurality of nucleic acid sensor molecules.

As defined herein, a **“profiling biosensor”** comprises a plurality of profiling nucleic acid sensor molecules.

As used herein, a molecule which **“naturally binds to DNA or RNA”** is one which is found within a cell in an organism found in nature.

As defined herein, a **“target modulation domain”** is the portion of a nucleic acid sensor molecule which recognizes a target molecule. The target modulation domain is also sometimes referred to herein as the **“target activation site”** or **“effector modulation domain”**.

As used herein a **“catalytic domain”** is the portion of a nucleic acid sensor molecule possessing catalytic activity which is modulated in response to binding of a target molecule to the target modulation domain.

As used herein, a **“linker region”** or **“linker domain”** is a portion of a nucleic acid sensor molecule by or at which the **“target modulation domain”** and **“catalytic domain”** are joined. Linker regions include, but are not limited to oligonucleotides of varying length, basepairing phosphodiester, phosphothiolate, and other covalent bonds, chemical moieties (*e.g.*, PEG), PNA, formacetal, bismaleimide, disulfide, and other bifunctional linker reagents. The linker domain is also sometimes referred to herein as a **“connector”** or **“stem”**.

As used herein, a **“random sequence”** or a **“randomized sequence”** is a segment of a nucleic acid having one or more regions of fully or partially random sequences. A fully random sequence is a sequence in which there is an approximately equal probability of each base (A, T, C, and G) being present at each position in the sequence. In a partially random

sequence, instead of a 25% chance that an A, T, C, or G base is present at each position, there are unequal probabilities.

As defined herein, **“a fixed region”** is a nucleic acid sequence which is known.

As defined herein, **“amplifying”** means any step or process or any combination of  
5 steps or processes that increases the amount or number of copies of a molecule or class of molecules.

As defined herein, a **“catalytic nucleic acid sensor molecule”** is a nucleic acid molecule comprising a target modulation domain, a linker region, and a catalytic domain.

As defined herein, an **“optical nucleic acid sensor molecule”** is a catalytic nucleic  
10 acid sensor molecule wherein the catalytic domain has been modified to emit an optical signal as a result of and/or in lieu of catalysis by the inclusion of an optical signal generating unit.

As defined herein, a **“nucleic acid sensor molecule”** or **“NASM”** refers to either or both of a catalytic nucleic acid sensor molecule and an optical nucleic acid sensor molecule.

As used herein, a **“signal”** is a detectable physical quantity, impulse or object.  
15

As used herein, an **“optical signal”** is a signal the optical properties of which can be detected.

As defined herein, an **“optical signal generating unit”** is a portion of a nucleic acid sensor molecule comprising one or more nucleic acid sequences and/or non-nucleic acid  
20 molecular entities, which change optical or electrochemical properties or which change the optical or electrochemical properties of molecules in close proximity to them in response to a change in the conformation or the activity of the nucleic acid sensor molecule following recognition of a target molecule by the target modulation domain.

As defined herein, a nucleic acid sensor molecule which **“recognizes a target  
25 molecule”** is a nucleic acid molecule whose activity is modulated upon binding of a target molecule to the TMD a greater extent than it is by the binding of any non-target molecule or in the absence of the target molecule. The recognition event between the nucleic acid sensor molecule and the target molecule need not be permanent during the time in which the resulting allosteric modulation occurs. Thus, the recognition event can be transient with  
30 respect to the ensuing allosteric modulation (*e.g.*, conformational change) of the nucleic acid precursor molecule or nucleic acid sensor molecule.

As defined herein, an **“array”** or **“microarray”** refers to a biosensor comprising a plurality of nucleic acid sensor molecules immobilized on a substrate.

As defined herein, a **“substrate”** refers to any physical supporting surface, whether rigid, flexible, solid, porous, gel-based, or of any other material or composition.

As defined herein, an **“amplicon”** is the sequence of a nucleic acid sensor molecule with ligase activity covalently ligated to an oligonucleotide substrate.

5 As defined herein, a **“cleavage substrate”** is an oligonucleotide or portion of an oligonucleotide cleaved upon target molecule recognized by a target modulation domain of an endonucleolytic nucleic acid sensor molecule.

As defined herein, an **“oligonucleotide substrate”** is an oligonucleotide that is acted upon by the catalytic domain of a nucleic acid sensor molecule.

10 As defined herein, an **“effector oligonucleotide”** is an oligonucleotide that base pairs with the effector oligonucleotide binding domain of a nucleic acid sensor molecule with ligase activity.

As defined herein, an **“effector oligonucleotide binding domain”** is the portion of the nucleic acid sensor molecule with ligase activity which is complementary to the effector  
15 oligonucleotide.

As defined herein, a **“capture oligonucleotide”** is an oligonucleotide that is used to attach a nucleic acid sensor molecule to a substrate by complementarity and/or hybridization.

As defined herein, an **“oligonucleotide substrate binding domain”** is the portion  
20 on the nucleic acid sensor molecule with ligase activity that is complementary to and can base pair with an oligonucleotide substrate.

As defined herein, a **“oligonucleotide supersubstrate”** is an oligonucleotide substrate that is complementary to and can base pair with the oligonucleotide substrate binding domain and to the effector oligonucleotide binding domain of a nucleic acid sensor  
25 molecule with ligase activity. The oligonucleotide supersubstrate may or may not carry an affinity tag.

As defined herein, a **“oligonucleotide supersubstrate binding domain”** is the region of a nucleic acid sensor molecule with ligase activity that is complementary to and can base pair with the oligonucleotide supersubstrate.

30 As defined herein, **“stem selection”** refers to a process performed on a pool of nucleic molecules comprising a target modulation domain, a catalytic domain and an oligonucleotide linker region wherein the linker region is fully or partially randomized.

As defined herein, **“rational design/engineering”** refers to a technique used to construct nucleic acid sensor molecules in which a non-conserved region of a ribozyme is

replaced with a target modulation domain and joined to the catalytic domain of the ribozyme by an oligonucleotide linker region.

As defined herein, “**amplicon dependent nucleic acid amplification**” refers to a technique by which one can amplify the signal of a nucleic acid sensor molecule by use of standard RT/PCR or Real-Time RT-PCR methods.”

As defined herein, “**switch factor**” is the enhancement observed in the catalytic activity and/or catalytic initial rate of a nucleic acid sensor molecule upon recognition of a target molecule by the target modulation domain.

As defined herein, a “**cis-ligase ribozyme**” is a ligase ribozyme that ligates its 3' end to its 5' end. The *cis*-ligase ribozyme is also referred herein as “**1-piece ligase**” and is a 1-component system where oligonucleotide substrate, oligonucleotide substrate binding domain, catalytic domain, effector oligonucleotide and effector oligonucleotide binding domains are fused in the format shown in Figure 39.

As defined herein, a “**trans-ligase ribozyme**” is a ligase ribozyme that ligates its 3' end to the 3' end of an oligonucleotide substrate.

As defined herein, a “**2-piece ligase**” is a 2-component *trans*-ligase ribozyme. The first component consists of the catalytic domain, the linker region, the target modulation domain, the substrate binding domain and the effector oligonucleotide binding domain. The second component is the oligonucleotide substrate that is complementary to the substrate binding domain and the effector oligonucleotide binding domain. This system follows the format shown in Figure 41.

As defined herein, a “**3-piece ligase**” is a 3-component *trans*-ligase ribozyme. The first component consists of the catalytic domain, the linker, the target modulation domain, the substrate binding domain and the effector oligonucleotide binding domain. The second component is the effector oligonucleotide that is complementary to the effector oligonucleotide binding domain. The third component is the oligonucleotide substrate that is complementary to the substrate binding domain. This system follows the format of the 3-piece ligase platform shown in Figure 39.

#### 1. Generating a Target Specific Nucleic Acid Sensor Molecule

Catalytic nucleic acid sensor molecules (NASMs) are selected which have a target molecule-sensitive catalytic activity (e.g., ligation or self-cleavage) from a pool of randomized oligonucleotides. The catalytic NASMs have a target modulation domain to which the target molecule specifically binds and a catalytic domain for mediating a catalytic

reaction. Binding of a target molecule to the target modulation domain triggers a conformation change and/or change in catalytic activity in the nucleic acid sensor molecule. In one embodiment, by modifying (*e.g.*, removing) at least a portion of the catalytic domain and coupling it to an optical signal generating unit, an optical nucleic acid sensor molecule  
5 is generated whose optical properties change upon binding of a target molecule to the target modulation domain. In one embodiment, the pool of randomized oligonucleotides comprises the catalytic site of a ribozyme.

**A. Selection and Generation of Catalytic Nucleic Acid Sensor Molecules**

10 In one embodiment, a heterogeneous population of oligonucleotide molecules comprising randomized sequences is screened to identify a nucleic acid sensor molecule having a catalytic activity which is modified (*e.g.*, activated) upon interaction with a target molecule. Each oligonucleotide in the population comprises a random sequence and at least one fixed sequence at its 5' and/or 3' end. In one embodiment, the fixed sequence comprises  
15 at least a portion of a catalytic site. In the embodiments shown in Figures 1 and 4, the random sequence is flanked at both ends with fixed sequences.

In one embodiment, the random sequence portion of the oligonucleotide is about 15-70 (*e.g.*, 30-40) nucleotides in length and can comprise ribonucleotides and/or deoxyribonucleotides. Random oligonucleotides can be synthesized from phosphodiester-  
20 linked nucleotides using solid phase oligonucleotide synthesis techniques well known in the art (*see, e.g.*, Froehler, et al., 1986a; 1986b). Oligonucleotides can also be synthesized using solution phase methods such as triester synthesis methods (*see, e.g.*, Sood, et al., 1977, and Hirose, et al., 1978). Typical syntheses carried out on automated DNA synthesis equipment yield  $10^{15}$ - $10^{17}$  molecules. Sufficiently large regions of random sequence in the sequence  
25 design increases the likelihood that each synthesized molecule is likely to represent a unique sequence.

To synthesize randomized sequences, mixtures of all four nucleotides are added at each nucleotide addition step during the synthesis process, allowing for random incorporation of nucleotides. In one embodiment, random oligonucleotides comprise  
30 entirely random sequences; however, in other embodiments, random oligonucleotide can comprise stretches of nonrandom or partially random sequences. Partially random sequences can be created by adding the four nucleotides in different molar ratios at each addition step.

To generate oligonucleotide populations which are resistant to nucleases and hydrolysis, modified oligonucleotides can be used and can include one or more substitute internucleotide linkages, altered sugars, altered bases, or combinations thereof. In one embodiment, oligonucleotides are provided in which the P(O)O group is replaced by P(O)S ("thioate"), P(S)S ("dithioate"), P(O)NR<sub>2</sub> ("amidate"), P(O)R, P(O)OR', CO or CH<sub>2</sub> ("formacetal") or 3'-amine (-NH-CH<sub>2</sub>-CH<sub>2</sub>-), wherein each R or R' is independently H or substituted or unsubstituted alkyl. Linkage groups can be attached to adjacent nucleotide through an -O-N-, or -S- linkage. Not all linkages in the oligonucleotide are required to be identical.

10 In further embodiments, the oligonucleotides comprise modified sugar groups, for example, one or more of the hydroxyl groups is replaced with halogen, aliphatic groups, or functionalized as ethers or amines. In one embodiment, the 2'-position of the furanose residue is substituted by any of an O-methyl, O-alkyl, O-allyl, S-alkyl, S-allyl, or halo group. Methods of synthesis of 2'-modified sugars are described in Sproat, et al., 1991; 15 Cotten, et al., 1991; and Hobbs, et al., 1973. The use of 2-fluoro-ribonucleotide oligomer molecules can increase the sensitivity of a nucleic acid sensor molecule for a target molecule by ten-to one hundred-fold over those generated using unsubstituted ribo- or deoxyribooligonucleotides (Pagratis, et al., 1997), providing additional binding interactions with a target molecule and increasing the stability of the nucleic acid sensor molecule's 20 secondary structure(s) (Kraus, et al., 1998; Pieken, et al., 1991; Lin, et al., 1994; Jellinek, et al. 1995; Pagratis, et al., 1997).

In the embodiments shown in Figures 1 and 4, the random sequence portion of the oligonucleotide is flanked by at least one fixed sequence which comprises a sequence shared by all the molecules of the oligonucleotide population. Fixed sequences include 25 sequences such as hybridization sites for PCR primers, promoter sequences for RNA polymerases (e.g., T3, T4, T7, SP6, and the like), restriction sites, or homopolymeric sequences, such as poly A or poly T tracts, catalytic cores (described further below), sites for selective binding to affinity columns, and other sequences to facilitate cloning and/or sequencing of an oligonucleotide of interest. In one embodiment, the fixed sequence is 30 approximately 50 nucleotides in length.

In a preferred embodiment, the fixed sequence comprises at least a portion of a catalytic site of an oligonucleotide molecule (e.g., a ribozyme) capable of catalyzing a chemical reaction. Catalytic sites are well known in the art and include, e.g. a ligase site (see Figure 2), the catalytic sites of Group I or Group II introns (see, e.g., U.S. Patent



Number 5,780,272), the catalytic core of a hammerhead ribozyme (*see, e.g.*, U.S. Patent Number 5,767,263 and U.S. Patent Number 5,700,923, and Figure 5, herein) or a hairpin ribozyme (*see, e.g.*, U.S. Patent Number 5,631,359. Other catalytic sites are disclosed in U.S. Patent Number 6,063,566, Koizumi et al., FEBS Lett. 239: 285-288 (1988), Haseloff and Gerlach, Nature 334: 585-59 (1988), Hampel and Tritz, Biochemistry 28: 4929-4933 (1989), Uhlenbeck, Nature, 328: 596-600 (1987), and Fedor and Uhlenbeck, Proc. Natl. Acad. Sci. USA 87: 1668-1672 (1990)).

Nucleic acid sensor molecules are generally selected in a 5 to 20 cycle procedure. In one embodiment, heterogeneity is introduced only in the initial selection stages and does not occur throughout the replicating process. Figure 1A shows a schematic diagram in which the oligonucleotide population is screened for a nucleic acid sensor molecule which comprises a target molecule activatable ligase activity. Figure 1B shows the hammerhead nucleic acid sensor molecule selection methodology. Each of these methods are readily modified for the selection of NASMs with other catalytic activities.

In the embodiment shown in Figure 1A, the ligation reaction involves covalent attachment of an oligonucleotide substrate to the 5'-end of the NASM through formation of a phosphodiester linkage. Other ligation chemistries can form the basis for selection of NASMs (*e.g.*, oligonucleotide ligation to the 3'-end, alkylation's (Wilson & Szostak), peptide bond formation (Zhang & Czech), Diels-Alder reactions to couple alkenes and dienes (Seelig & Jaschke), etc.). For some chemistries, the chemical functional groups that constitute the reactants in the ligation reaction may not naturally appear within nucleic acids. Thus, it may be necessary to synthesize an RNA pool in which one of the ligation reactants is covalently attached to each member of the pool (*e.g.*, attaching a primary amine to the 5'-end of an RNA to enable selection for peptide bond formation).

In this embodiment, the oligonucleotide population from which the NASMs will be selected is initially screened in a negative selection procedure to eliminate any molecules which have ligase activity even in the absence of target molecule binding. A solution of oligonucleotides (*e.g.*, 100 pM) comprising a 5' and 3' fixed sequence ("5'-fixed: random: 3'-fixed") is denatured with a 3' primer sequence ("3' prime") (*e.g.*, 200 pM) which binds to at least a portion of the 3' fixed sequence. In one embodiment, the 5'-fixed:random:3'-fixed sequence is 5'.

GGACUUCGGUCCAGUGCUCGUGCACUAGGCCGUUCGACC-N<sub>30-50</sub>

CUUAGACAGGAGGUUAGGUGCCUCGUGAUGUCCAGUCGC-3' (SEQ ID NO:1), where N represents a random sequence having 30 to 50 nucleotides and the 3' primer sequence used is 5'-GCGACTGGACATCACGAG-3' (SEQ ID NO:2).

Ligation buffer (*e.g.*, 30 mM Tris HCl, pH 7.4, 600 mM NaCl, 1 mM EDTA, 1% NP-40, 60 mM MgCl<sub>2</sub>) and a tagged oligonucleotide substrate sequence ("tag-substrate") (*e.g.*, Tag-UGCCACU) are added and the mixture is incubated for about 16 to about 24 hours at 25 °C in the absence of target molecule (STEP 1). Tags encompassed within the scope include, *e.g.*, radioactive labels, fluorescent labels, a chemically reactive species such as thiophosphate, the first member of a binding pair comprising a first and second binding member, each member bindable to the other (*e.g.*, biotin, an antigen recognized by an antibody, or a tag nucleic acid sequence). The reaction is stopped by the addition of EDTA. Alternatively, the reaction can be terminated by removal of the substrate or addition of denaturants (*e.g.* urea, formamide).

Ligated molecules are removed from pool of selectable molecules (STEP 2), generating a population of oligonucleotides substantially free of ligated molecules (as measured by absence of the tag sequence in the solution). In the embodiment shown in Figure 1A, the tag is the first member of a binding pair (*e.g.*, biotin) and the ligated molecules ("biotin-oligonucleotide substrate:5'-fixed:random:3'-fixed") are physically removed from the solution by contacting the sample to a solid support to which the second member of the binding pair is bound ("S") (*e.g.*, streptavidin). The eluant collected comprises a population of oligonucleotides enriched for non-ligated molecules (5'-fixed:random:3'-fixed). This step can be repeated multiple times until the oligonucleotide population is substantially free of molecules having target-insensitive ligase activity.

This step allows for suppression of the ability of constitutively active molecules to be carried through to the next cycle of selection. Physical separation of ligated and unligated molecules is one mechanism by which this can be achieved. Alternatively, the negative selection step can be configured such that catalysis converts active molecules to a form that blocks their ability to be either retained during the subsequent positive selection step or to be amplified for the next cycle of selection. For example, the oligonucleotide substrate used for ligation in the negative selection step can be synthesized without a capture tag. Target-independent ligases covalently self-attach the untagged oligonucleotide substrate during the negative selection step and are then unable to accept a tagged form of the oligonucleotide substrate provided during the positive selection step that follows. In another embodiment, the oligonucleotide substrate provided during the negative selection

step has a different sequence from that provided during the positive selection step. When PCR is carried out using a primer complementary to the positive selection oligonucleotide substrate, only target-activated ligases will be capable of amplification.

5 A positive selection phase follows. In this phase, more 3' primer and tagged oligonucleotide substrate are added to the pool resulting from the negative selection step. Target molecules are then added to form a reacted solution and the reacted solution is incubated at 25 °C for about 2 hours (STEP 3). Target molecules encompassed within the scope include, *e.g.*, proteins or portions thereof (*e.g.*, receptors, antigen, antibodies, enzymes, growth factors), peptides, enzyme inhibitors, hormones, carbohydrates, 10 polysaccharides, glycoproteins, lipids, phospholipids, metabolites, metal ions, cofactors, inhibitors, drugs, dyes, vitamins, nucleic acids, membrane structures, receptors, organelles, and viruses. Target molecules can be free in solution or can be part of a larger cellular structure (*e.g.*, such as a receptor embedded in a cell membrane). In one embodiment, a target molecule is one which does not naturally bind to nucleic acids.

15 In one embodiment, nucleic acid sensor molecules are selected which are activated by target molecules comprising molecules having an identified biological activity (*e.g.*, a known enzymatic activity, receptor activity, or a known structural role); however, in another embodiment, the biological activity of at least one of the target molecules is unknown (*e.g.*, the target molecule is a polypeptide expressed from the open reading frame of an EST 20 sequence, or is an uncharacterized polypeptide synthesized based on a predicted open reading frame, or is a purified or semi-purified protein whose function is unknown).

Although in one embodiment the target molecule does not naturally bind to nucleic acids, in another embodiment, the target molecule does bind in a sequence specific or non-specific manner to a nucleic acid sensor molecule. In a further embodiment, a plurality of 25 target molecules binds to the nucleic acid sensor molecule. Selection for NASMs specifically responsive to a plurality of target molecules (*i.e.* not activated by single targets within the plurality) may be achieved by including at least two negative selection steps in which subsets of the target molecules are provided.

In still a further embodiment, nucleic acid sensor molecules are selected which bind 30 specifically to a modified target molecule but which do not bind to non-modified target molecules. Targeted modifications include, *e.g.*, post-translational modifications of a protein, such as phosphorylation, ribosylation, methylation (Arg, Asp, N, S, or O-directed), prenylation (*e.g.*, farnesyl, geranylgeranyl, and the like), acetylation, acylation, allelic variations within a protein (*e.g.*, single amino acid changes in a protein) and cleavage sites

in a protein. In another embodiment, intermediates in a chemical synthesis pathway can be targeted, as well as starting and final products. In still a further embodiment, stereochemically distinct species of a molecules can be targeted.

The reacted solution is enriched for ligated molecules (biotin-oligonucleotide  
5 substrate: 5'-fixed:random:3'-fixed) by removing non-tagged molecules (5'-  
fixed:random:3'-fixed) from the solution. For example, in one embodiment, the tagged  
oligonucleotide substrate comprises a biotin tag and ligated molecules are isolated by  
passing the reacted solution over a solid support to which streptavidin (S) is bound (STEP  
4). Eluant containing non-bound, non-ligated molecules (5'-fixed:random:3'-fixed) is  
10 discarded and bound, ligated molecules (biotin-oligonucleotide substrate: 5'-  
fixed:random:3'-fixed) are identified as nucleic acid sensor molecules and released from the  
support by disrupting the binding pair interaction which enabled capture of the catalytically  
active molecules. For example, heating to 95° C in the presence of 10 mM biotin allows  
release of biotin-tagged catalysts from an immobilized streptavidin support. In another  
15 embodiment, the captured catalysts remain attached to a solid support and are directly  
amplified (described below) while immobilized. Multiple positive selection phases can be  
performed (STEPS 3 and 4). In one embodiment, the stringency of each positive selection  
phase is increased by decreasing the incubation time by one half.

Physically removing inactive species from the pool adds stringency to the selection  
20 process. However, to the extent that the ligation reaction increases the amplification  
potential of the NASMs, this step may be omitted. In the illustrated embodiment, for  
example, ligation of an oligonucleotide to the active species provides a primer binding site  
that enables subsequent PCR amplification using an oligonucleotide substrate  
complementary to the original oligonucleotide substrate. Unligated species do not  
25 necessarily need to be physically separated from other species because they are less likely to  
amplify in the absence of a covalently tethered primer binding site. Selected nucleic acid  
sensor molecules are amplified (or in the case of RNA molecules, first reverse transcribed,  
then amplified) using an oligonucleotide substrate primer ("S primer") (e.g., 5'-  
AAAAAATGCACTGGACT-3' (SEQ ID NO:3)) which specifically binds to the litgated  
30 oligonucleotide substrate sequence (STEP 5). In one embodiment, amplified molecules are  
further amplified with a nested PCR primer that regenerates a T7 promoter ("T7 Primer")  
from the 5' fixed and the litigated oligonucleotide substrate sequence (STEP 6). Following  
transcription with T7 RNA polymerase (STEP 7), the oligonucleotide pool may be further  
selected and amplified to eliminate any remaining unligated sequences (5'-fixed:random:3'-

fixed) by repeating STEPS 3-7. It should be obvious to those of skill in the art that in addition to PCR, and RT-PCR, any number of amplification methods can be used (either enzymatic, chemical, or replication-based, *e.g.*, such as by cloning), either singly, or in combination. Exemplary amplification methods are disclosed in Saiki, et al., 1985; Saiki, et al., 1988; Kwok, et al., 1989; Joyce, 1989; and Guatelli, et al., 1990.

Because the 3' primer (3' prime) (see STEP 3 in Figure 1A) is included in the ligation mixture, selected nucleic acid sensor molecules may require this sequence for activation. In cases where this is undesirable, the 3' primer may be omitted from the mix. Alternatively, the final nucleic acid sensor molecule can be modified by attaching the 3' primer via a short sequence loop or a chemical linker to the 3' end of the nucleic acid sensor molecule, thereby eliminating the requirement for added primer, allowing 3' primer sequence to self-prime the molecule.

As shown in Figure 1B, selection of a nucleic acid sensor molecule begins with the synthesis of a ribozyme sequence on a DNA synthesizer. Random nucleotides are incorporated generating pools of roughly  $10^{16}$  molecules. Most molecules in this pool are non-functional, but a handful will respond to a given target and be useful as nucleic acid sensor molecules. Sorting among the billions of species to find the desired molecules starts from the complex sequence pool. Nucleic acid sensor molecules are isolated by an iterative process: in addition to the target-activated ribozymes that one desires, the starting pool is usually dominated by either constitutively active or completely inactive ribozymes. The selection process removes both types of contaminants. In the following amplification stage, thousands of copies of the surviving sequences are generated to enable the next round of selection. During amplification, random mutations can be introduced into the copied molecules — this 'genetic noise' allows functional NASMs to continuously evolve and become even better adapted as target-activated molecules. The entire experiment reduces the pool complexity from  $10^{16}$  down to  $< 100$ .

**Pool preparation.** The starting library of DNA sequences is generated by automated chemical synthesis on a DNA synthesizer. This library of sequences is transcribed *in vitro* into RNA using T7 RNA polymerase and subsequently purified.

**Negative selection incubation.** In the absence of the desired target molecule of interest, the RNA library is incubated together with the binding buffer alone. During this incubation, non-allosteric (or non-target activated) ribozymes are expected to undergo cleavage.

**Size-based Purification.** Undesired members of the hammerhead pool, those that are constitutively active in the absence of the target molecule, are removed from the unreacted members by PAGE-chromatography; 7M Urea, 8% acrylamide, 1X TBE. Higher molecular weight species are eluted as a single broad band from the gel matrix into TBE buffer, then purified for subsequent steps in the selection cycle.

**Positive selection incubation.** The remaining RNA pool is then incubated under identical conditions but now in the presence of the target molecule of interest in binding buffer.

**Size-based Purification.** In this step, desired members of the hammerhead pool, those that are only active in the presence of the target molecule, are removed from the remaining unreacted members by PAGE-chromatography; 7M Urea, 8% acrylamide, 1X TBE. In this step, lower molecular weight species are eluted as a single broad band from the gel matrix into TBE buffer, then purified for subsequent steps in the selection cycle.

**Amplification.** RT-PCR amplified DNA is purified and transcribed to yield an enriched pool for a subsequent round of reselection.

**Iteratively repeat.** Rounds of selection and amplification (steps 2-5) are repeated until functional members sufficiently dominate the resultant library.

In another embodiment, as shown in Figure 4, an oligonucleotide population is screened for a nucleic acid sensor molecule which comprises a target molecule having activatable self-cleaving activity. In this embodiment, the starting population of oligonucleotide molecules comprises 5' and 3' fixed regions ("5'-fixed and 3' fixed A-3'fixed B") and at least one of the fixed regions, in this example, the 3' fixed region, comprises a ribozyme catalytic core including a self cleavage site (the junction between 3' fixed A-3' fixed B). In one embodiment, the 5'-fixed: random:3' fixed A-3'-fixed B molecule is GGGCGACCCUGAUGAGCCUGG-N<sub>20-50</sub>-UUAGACGAAACGGUGAAAGCCGUAGGUUGCCC (SEQ ID NO:4), where N<sub>20-50</sub> is a random sequence of 20-50 nucleotides.

The population of oligonucleotide molecules comprising random oligonucleotides flanked by fixed 5' and 3' sequences (5'-fixed:random:3'-fixed A: 3' fixed B) are negatively selected to remove oligonucleotides which self-cleave (*i.e.*, 5'-fixed:random:3'-fixed-A molecules) even in the absence of target molecules. The oligonucleotide pool is incubated in reaction buffer (*e.g.*, 50 mM Tris HCl, pH 7.5, 20 mM MgCl<sub>2</sub>) for 5 hours at 25 °C, punctuated at one hour intervals by incubation at 60 °C for one minutes (STEP 1). In one embodiment, the uncleaved fraction of the oligonucleotide population (containing 5'-

fixed and 3' fixed A-3'-fixed B molecules) is purified by denaturing 10% polyacrylamide gel electrophoresis (PAGE) (STEP 2). Target molecule dependent cleavage activity is then selected in the presence of target molecules in the presence of reaction buffer by incubation at 23 °C for about 30 seconds to about five minutes (STEP 3). Cleaved molecules (5'-  
 5 fixed:random:3'fixed-A molecules) are identified as nucleic acid sensor molecules and are purified by PAGE (STEP 4).

Amplification of the cleaved molecule is performed using primers which specifically bind the 5'-fixed and the 3'-fixed A sequences, regenerating the T7 promoter and the 3'-fixed B site (STEP 5), and the molecule is further amplified further by RNA transcription  
 10 using T7 polymerase (STEP 6). In one embodiment, the process (STEPS 1-6) is repeated until the nucleic acid sensor precursor population is reduced to about one to five unique sequences.

Alternative methods for separating cleaved from uncleaved RNAs can be used. Tags can be attached to the 3'-fixed B sequence and separation can be based upon  
 15 separating tagged sequences from non-tagged sequences at STEP 4. Chromatographic procedures that separate molecules on the basis of size (e.g., gel filtration) can be used in place of electrophoresis. One end of each molecule in the RNA pool can be attached to a solid support and catalytically active molecules isolated upon release from the support as a result of cleavage. Alternate catalytic cores may be used. These alternate catalytic cores  
 20 and methods using these cores are also encompassed within the scope of the invention.

Nucleic acid sensor molecules which combine both cleavage and ligase activities in a single molecule can be isolated by using one or a combination of both of the selection strategies outlined independently above for ligases and endonucleases. For example, the hairpin ribozyme is known to catalyze cleavage followed by ligation of a second  
 25 oligonucleotide substrate (Berzal-Herranz et al.). Target activated sensor precursors based on the hairpin activity can be isolated from a pool of randomized sequence RNAs prepared as described previously with a sequence of the form 5'-

GGAGTTACCTAACAAGAAACAGNgaagtcaaccagNgaaacNCACGGAGACGTGNNaNat  
 taNctggt(N<sub>20</sub>-

30 N<sub>50</sub>)GGACCTACTGAGCTGACAGTCCTGTTTGATGCATACCGAGTAAGTG-3' (SEQ ID NO:36) where N indicates any nucleotide, lower case letters represent doped nucleotides, and uppercase letters represent fixed nucleotides. Hairpin-based NASMs can be isolated on the basis of target molecule dependent release of the fragment 5'-  
 GUCCUGUUUGAUGCAUACCGAGUAAGUG-3' (SEQ ID NO:74) in the same way that

hammerhead-based NASMs are isolated (*e.g.*, target molecule dependent increase in electrophoretic mobility or target molecule dependent release from a solid support).

Alternatively, nucleic acid sensor molecules can be selected on the basis of their ability to substitute the 3'-sequence released upon cleavage for another sequence as described in an  
5 target molecule independent manner by Berzal-Heranz et al. In this scheme, the original 3'-end of the NASM is released in an initial cleavage event and an exogenously provided oligonucleotide substrate with a free 5'-hydroxyl is ligated back on. The newly attached 3'-end provides a primer binding site that can form the basis for preferential amplification of catalytically active molecules. Constitutively active molecules that are not activated by a  
10 provided target molecule can be removed from the pool by (1) separating away molecules that exhibit increased electrophoretic mobility in the absence of an exogenous oligonucleotide substrate or in the absence of target molecule, or (2) capturing molecules that acquire an exogenous oligonucleotide substrate (*e.g.*, using a 3'-biotinylated substrate and captured re-ligated species on an avidin column).

15 Like the hairpin ribozyme, the group I intron self-splicing ribozymes combine cleavage and ligation activities to promote ligation of the exons that flank it. In the first step of group I intron-catalyzed splicing, an exogenous guanosine cofactor attacks the 5'-splice site. As a result of an intron-mediated phosphodiester exchange reaction, the 5'-exon is released coincident with attachment of the guanosine cofactor to the ribozyme. In a  
20 second chemical step, the 3'-hydroxyl at the end of the 5'-exon attacks the phosphodiester linkage between the intron and the 3'-exon, leading to ligation of the two exons and release of the intron. Group I intron-derived NASMs can be isolated from degenerate sequence pools by selecting molecules on the basis of either one or both chemical steps, operating in either a forward or reverse direction. NASMs can be isolated by specifically enriching  
25 those molecules that fail to promote catalysis in the absence of target molecule but which are catalytically active in its presence. Specific examples of selection schemes follow. In each case, a pool of RNAs related in sequence to a representative group I intron (*e.g.*, the *Tetrahymena thermophila* pre-rRNA intron or the phage T4 td intron) serves as the starting point for selection. Random sequence regions can be embedded within the intron at sites  
30 known to be important for proper folding and activity (*e.g.*, substituting the P5abc domain of the *Tetrahymena* intron, Williams et al.). Intron nucleic acid sensor molecules, in this case, sensitive to thio-GMP can be generated as follows.

*First step, forward direction*



The intron is synthesized with a short 5'-exon. In the negative selection step, a guanosine cofactor is provided and constitutively active molecules undergo splicing. In the positive selection step, the target molecule is provided together with thio-GMP. Molecules responsive to the target undergo activated splicing and as a result acquire a unique  
5 thiophosphate at their 5'-termini. Thio-tagged NASMs can be separated from untagged ribozymes by their specific retention on mercury gels or activated thiol agarose columns.

*First step, reverse direction*

The method is performed as described in Green & Szostak. An intron is synthesized with a 5'-guanosine and no 5'-exon. An oligonucleotide substrate complementary to the 5'-  
10 internal guide sequence is provided during the negative selection step and constitutively active molecules ligate the substrate to their 5'-ends, releasing the original terminal guanosine. A second oligonucleotide substrate with a different 5'-sequence is provided together with target in the positive selection step. NASMs specifically activated by the target molecule ligate the second oligonucleotide substrate to their 5'-ends. PCR  
15 amplification using a primer corresponding to the second substrate can be carried out to preferentially amplify target molecule sensitive nucleic acid sensor molecules.

*Second step, reverse direction*

The method is performed as described in Robertson & Joyce. The intron is synthesized with no flanking exons. During the negative selection step, pool RNAs are  
20 incubated together with a short oligonucleotide substrate under conditions which allow catalysis to proceed. During the positive selection step, a second oligonucleotide substrate with a different 3'-sequence is provided together with the sensor target. NASMs are activated and catalyze ligation of the 3'-end of the second substrate. Reverse transcription carried out using a primer complementary to the 3'-end of the second substrate specifically  
25 selects NASMs for subsequent amplification.

Once nucleic acid sensor molecules are identified, they can be isolated, cloned, sequenced, and/or resynthesized using natural or modified nucleotides. Accordingly, synthesis intermediates of nucleic acid sensor molecules are also encompassed within the scope, as are replicatable sequences (e.g., plasmids) comprising nucleic acid sensor  
30 precursor molecules and nucleic acid sensor molecules.

**B. Converting a catalytic nucleic acid sensor molecule to an**

optical nucleic acid sensor molecule

The nucleic acid sensor molecules identified above through *in vitro* selection comprise a catalytic domain (i.e., a signal generating moiety), coupled to a target modulation domain, (i.e., a domain which recognizes a target molecule and which transduces that molecular recognition event into the generation of a detectable signal). In general, the target modulation domain is defined by the minimum number of nucleotides sufficient to create a three-dimensional structure which recognizes a target molecule. In addition, the nucleic acid sensor molecules of the present invention use the energy of molecular recognition to modulate the catalytic or conformational properties of the nucleic acid sensor molecule. The selection process as described in detail in the present invention identifies novel nucleic acid sensor molecules through target modulation of the catalytic core of a ribozyme. Hence, the *in vitro* selection procedures described herein are distinct from those previously described for affinity-based aptamer selections (e.g., SELEX) in that we show that selective pressure on the starting population of NASMs (starting pool size is as high as  $10^{14}$  to  $10^{17}$  molecules) results in nucleic acid sensor molecules with enhanced catalytic properties, but not in enhanced binding properties, Figure 59. Specifically, the NASM selection procedures place selective pressure on catalytic effectiveness of potential NASMS by modulating both target concentration and reaction time-dependence. Either parameter, when optimized throughout the selection, can lead to nucleic acid molecular sensor molecules which have custom-designed catalytic properties, e.g., NASMs that have high switch factors (Figure 47 and Figure 59), and or NASMs that have high specificity (Figure 57). The kinetic properties of the NASMs of the present invention are consistent with that obtained from a nucleoprotein-selection reported previously by Robertson and Ellington (2001) in which the resulting ribozyme (switch factor equal to 1,700 fold) has the same affinity for RNA (1  $\mu$ M) as did the starting pool.

In the examples described above, the catalytic site is a known sequence (a ligase site or a hammerhead catalytic core) and is at least a portion of either the 5' and/or 3' fixed region (the other portion being supplied by the random sequence), or is a complete catalytic site. However, in other embodiments, the catalytic site may be selected along with the target molecule binding activity of oligonucleotides within the oligonucleotide pool.

In one embodiment, in order to convert a catalytic nucleic acid sensor molecule into an optical nucleic acid sensor molecule, at least a portion of the catalytic domain is

modified (*e.g.*, deleted). In one embodiment, the deletion enhances the conformational stability of the optical nucleic acid sensor molecule in either the bound or unbound forms. In one embodiment, shown in Figures 6A and B, deletion of the entire catalytic domain of the catalytic NASM shown in Figure 5 is shown to stabilize the unbound form of the nucleic acid sensor molecule. In another embodiment, the deletion may be chosen so as to take advantage of the inherent fluorescence-quenching properties of unpaired guanosine (G) residues (Walter, N.G. and Burke, J.M., "Real-time monitoring of hairpin ribozyme kinetics through base-specific quenching of fluorescein-labeled substrates", RNA 3:392 (1997).)

In another embodiment, the target modulation domain from a previously identified nucleic acid sensor molecule is incorporated into an oligonucleotide sequence that changes conformation (*e.g.*, from a duplexed hairpin to a G-quadruplex) upon target binding. Optical Nucleic acid sensor molecules of this type can be derived from allosteric ribozymes, such as those derived from the hammerhead, hairpin, L1 ligase, or group 1 intron ribozymes and the like, all of which transduce molecular recognition into a detectable signal. For example, 3',5'-cyclic nucleotide monophosphate (cNMP)-dependent hammerhead ribozymes were reengineered into (RNA) optical nucleic acid sensor molecules which specifically bound to cNMP (Garretta et al., 2001). The catalytic cores for hammerhead ribozymes were removed and replaced with 5-base duplex forming sequences. The binding of these reengineered RNA optical NASMs to c-NMP was then confirmed experimentally. By adjusting the duplex length, they can be redesigned to undergo significant conformational changes. The conformational changes can be coupled to detection *via* FRET or simply changes in fluorescence intensity (as in the case of a molecular beacon). For example, by adding an appropriate probe on the each end of the duplex, the stabilization of duplex by target binding can be monitored with the change in fluorescence.

While the above experimental example is performed in solution and utilizes a cuvette-based fluorescence spectrometer, in alternative embodiments the methods are performed in microwell multiplate readers (*e.g.*, the Packard Fusion, or the Tecan Ultra) for high-throughput solution phase measurements.

In another embodiment, a nucleic acid sensor molecule is bound to a surface by a linker attached to one end of the molecule. For example, a nucleic acid sensor molecule is modified to include a 12 carbon linker terminated with an amine group. This free amine group allows the NASM to be attached to an aldehyde-derivatized glass surface via standard protocols for Schiff base formation and reduction. The nucleic acid sensor molecules can be bound in discrete regions or spots to form an array or uniformly distributed to cover an

extended area. In the absence of target molecule, the optical nucleic acid sensor molecule forms a stem-loop conformation with duplex formation along the stem due to the complementarity of the nucleotides at the 3' and 5' ends of the molecule. In the presence of the target molecule, the optical nucleic acid sensor molecule undergoes a conformational rearrangement. In some embodiments, this conformational rearrangement results in a change in the distance between the fluorophore attached to the 5' end and the quencher attached to the 3' end. With the quencher separated from the fluorophore, the detected fluorescence emission intensity from the fluorophore increases sharply. The detected increase in fluorescence intensity with target molecule concentration can be used to detect and quantify the amount of target present in a sample solution introduced onto the surface. A sample solution can be laterally confined about the sensor surface by a coverslip, microwell, incubation chamber seal, or flowcell.

In one embodiment, after deletion of at least a portion of the catalytic site from a catalytic nucleic acid sensor molecule, an optical signaling unit is either added to, or inserted within, the nucleic acid sensor molecule, generating a sensor molecule whose optical properties change in response to binding of the target molecule to the target modulation domain. In one embodiment, the optical signaling unit is added by exposing at least a 5' or 3' nucleotide that was not previously exposed. The 5' nucleotide or a 5' subterminal nucleotide (e.g., an internal nucleotide) of the molecule is couplable to a first signaling moiety while the 3' nucleotide or 3' subterminal nucleotide is couplable to a second signaling moiety. Target molecule binding to the optical nucleic acid sensor molecule alters the proximity of the 5' and 3' nucleotide (or subterminal nucleotides) with respect to each other, and when the first and second signaling moieties are coupled to their respective nucleotides, this change in proximity results in a target sensitive change in the optical properties of the nucleic acid sensor molecule. Detection of changes in the optical properties of the nucleic acid sensor molecule can therefore be correlated with the presence and/or quantity of a target molecule in a sample.

In another embodiment optical NASMs are generated by adding first and second signaling moieties, that are coupled to the 5' terminal or subterminal sequences, and 3' terminal and subterminal sequences respectively, of the catalytic NASM. Signaling molecules can be coupled to nucleotides which are already part of the nucleic acid sensor molecule or may be coupled to nucleotides which are inserted into the nucleic acid sensor molecule, or can be added to a nucleic acid sensor molecule as it is synthesized. Coupling chemistries to attach signaling molecules are well known in the art (see, for example, The

Molecular Probes Handbook, R. Haugland). Suitable chemistries include, *e.g.*, derivatization of the 5-position of pyrimidine bases (*e.g.*, using 5'-amino allyl precursors), derivatization of the 5'-end (*e.g.*, phosphoroamidites that add a primary amine to the 5'-end of chemically-synthesized oligonucleotide) or the 3'-end (*e.g.*, periodate treatment of RNA to convert the 3'-ribose into a dialdehyde which can subsequently react with hydrazide-bearing signaling molecules).

In another embodiment, a single signaling moiety is either added to, or inserted within, the catalytic nucleic sensor molecule. In this embodiment, binding of the target molecule results in changes in both the conformation and physical aspect (*e.g.*, molecular volume, and thus rotational diffusion rate, etc.) of the optical nucleic acid sensor molecule. Conformational changes in the optical nucleic acid sensor molecule upon target binding will modify the chemical environment of the signaling moiety, while changes in the physical aspect of the nucleic acid sensor molecule will alter the kinetic properties of the signaling moiety. In both cases, the result will be a detectable change in the optical properties of the nucleic acid sensor molecule.

In one embodiment, the optical nucleic acid sensor molecule is prepared without a quencher group. Instead of a quencher group a moiety with a free amine group can be added. This free amine group allows the sensor molecule to be attached to an aldehyde-derivatized glass surface via standard protocols for Schiff base formation and reduction. The nucleic acid sensor molecules can be bound in discrete regions or spots to form an array, or uniformly distributed to cover an extended area. In the absence of target, the optical nucleic acid sensor molecule will diffusionally rotate about its point of attachment to the surface at a rate characteristic of its molecular volume and mass. After target binding, the optical NASM-target complex will have a correspondingly larger volume and mass. This change in molecular volume (mass) will slow the rate of rotational diffusion, and result in a measurable change in the polarization state of the fluorescence emission from the fluorophore.

In one embodiment of the invention, a single signaling moiety is attached to a portion of a catalytic NASM that is released as a result of catalysis (*e.g.*, either end of a self-cleaving ribozyme or the pyrophosphate at the 5'-end of a ligase). Target molecule-activated catalysis leads to release of the signaling moiety from the optical NASM to generate a signal correlated with the presence of the target. Release can be detected by either (1) changes in the intrinsic optical properties of the signaling moiety (*e.g.*, decreased fluorescence polarization as the released moiety is able to tumble more freely in solution),

or (2) changes in the partitioning of the signaling moiety (*e.g.*, release of a fluorophore from a chip containing immobilized ribozymes such that the total fluorescence of the chip is reduced following washing).

In another embodiment of the invention, the catalytic nucleic acid sensor molecule is  
5 unmodified and the optical signaling unit is provided as a substrate for the NASM. One example of this embodiment includes a fluorescently tagged oligonucleotide substrate which can be joined to a NASM with ligase activity. In a heterogeneous assay using the ligase as a sensor molecule, analyte-containing samples are incubated with the fluorescent oligonucleotide substrate and the ligase under conditions that allow the ligase to function.  
10 Following an incubation period, the ligase is separated from free oligonucleotide substrate (*e.g.*, by capturing ligases onto a solid support on the basis of hybridization to ligase-specific sequences or by pre-immobilizing the ligases on a solid support and washing extensively).

Quantitation of the captured fluorescence signal provides a means for inferring the  
15 concentration of analyte in the sample. In a second example of this embodiment, catalytic activity alters the fluorescence properties of an oligonucleotide substrate without leading to its own modification. Fluorophore pairs or fluorophore/quencher pairs can be attached to nucleotides flanking either side of the cleavage site of an oligonucleotide substrate for a trans-acting endonuclease ribozyme (Jenne *et al.*). Target activated cleavage of the  
20 substrate leads to separation of the pair and a change in its optical properties.

In another embodiment of the invention, the ligase catalytic NASM and its oligonucleotide substrates are unmodified and detection relies on catalytically-coupled changes in the ability of the NASM to be enzymatically amplified. In one example, a target-activated ligase is incubated together with oligonucleotide substrate and an analyte-  
25 containing sample under conditions which allow the ligase to function. Following an incubation period, the reaction is quenched and the mixture subjected to RT/PCR amplification using a primer pair that includes the oligo sequence corresponding to the ligation substrate. Amplification products can be detected by a variety of generally practiced methods (*e.g.* Taqman®). Only those ribozymes that have self-ligated an  
30 oligonucleotide substrate are capable of amplification under these conditions and will generate a signal that can be coupled to the concentration of the sensor target.

i. Proximity Dependent Signaling Moieties

Many proximity dependent signaling moieties are known in the art and are encompassed within the scope of the present invention (*see, e.g.*, Morrison, 1992, in Nonisotopic DNA Probe Techniques, Kricka, ed., Academic Press, Inc., San Diego, Calif., chapter 13; and Heller and Morrison, 1985, in Rapid Detection and Identification of  
5 Infectious Agents, Academic Press, Inc., San Diego, Calif., pages 245-256). Systems using these signaling moieties rely on the change in fluorescence that occurs when the moieties are brought into close proximity. Such systems are described in the literature as fluorescence energy transfer (FET), fluorescence resonance energy transfer (FRET),  
10 nonradiative energy transfer, long-range energy transfer, dipole-coupled energy transfer, or Forster energy transfer (*see, e.g.*, U.S. Patent Number 5,491,063, Wu, and Brand, 1994).

Other proximity-dependent signaling systems that do not rely on direct energy transfer between signaling moieties are also known in the art and can be used in the methods described herein. These include, *e.g.*, systems in which a signaling moiety is  
15 stimulated to fluoresce or luminesce upon activation by the target molecule. This activation may be direct (*e.g.*, as in the case of scintillation proximity assays (SPA), via a photon or radionuclide decay product emitted by the bound target), or indirect (*e.g.*, as in the case of AlphaScreen™ assays, via reaction with singlet oxygen released from a photosensitized donor bead upon illumination). In both scenarios, the activation of detected signaling  
20 moiety is dependent on close proximity of the signaling moiety and the activating species. In general, for both fluorescence, fluorescence polarization, and scintillation-proximity-type assays, the nucleic acid sensor molecule may be utilized in either solution-phase or solid-phase formats. That is, in functional form, the nucleic acid sensor molecule may be tethered (directly, or via a linker) to a solid support or free in solution.

25 In one embodiment, a scintillation proximity assay (SPA) is used. In this embodiment, the nucleic acid sensor molecules, ligate on oligonucleotide substrate in the presence of a target molecule (*see* Figures 2A and B), are bound to a scintillant-impregnated microwell plate (*e.g.*, FlashPlates, NEN Life Sciences Products, Boston, MA) coated with, for example, streptavidin via a (biotin) linker attached to the 5' end of the effector  
30 oligonucleotide sequence (for example, GCGACTGGACATCACGAG (SEQ ID NO:51) in Figure 2A). The various plate-sensor coupling chemistries are determined by the type and manufacturer of the plates, and are well-known in the art. Upon the addition of a solution containing target molecule and excess radiolabeled (*e.g.*, with <sup>32</sup>P or <sup>35</sup>S) oligonucleotide

substrate in ligation buffer, the NASMs hybridize and ligate the substrate oligonucleotide. Some fraction of the radiolabeled oligonucleotide substrate will be ligated to surface-immobilized NASMs on the plate, while unligated oligonucleotide substrate will be free in solution. Only those oligonucleotide substrates ligated to surface-immobilized NASMs on the plate will be in close enough proximity to the scintillant molecules embedded in the plate to excite them, thereby stimulating luminescence which can be easily detected using a luminometer (e.g., the TopCount luminescence plate reader, Packard Biosciences, Meriden, CT). This type of homogeneous assay format provides straightforward, real-time detection, quantification, and kinetic properties of target molecule binding.

10 In another embodiment, a similar SPA assay format is performed using scintillant-impregnated beads (e.g., Amersham Pharmacia Biotech, Inc., Piscataway, NJ). In this embodiment, the NASMs which ligate on oligonucleotide substrate in the presence of a target molecule (see Figures 2A and B) are coupled to scintillant-impregnated beads which are suspended in solution in, for example, a microwell plate. The various bead-sensor coupling chemistries are determined by the type and manufacturer of the beads, and are well-known in the art. Upon the addition of a solution containing target molecule and excess radiolabeled (e.g., with  $^{32}\text{P}$  or  $^{35}\text{S}$ ) oligonucleotide substrate in ligation buffer, the NASMs hybridize and ligate the oligonucleotide substrate. Some fraction of the radiolabeled substrate will be ligated to surface-immobilized NASMs on the beads, while unligated substrate will be free in solution. Only those substrates ligated to surface-immobilized NASMs on the beads will be in close enough proximity to the scintillant molecules embedded in the beads to excite them, thereby stimulating luminescence which can be easily detected using a luminometer (e.g., the TopCount luminescence plate reader, Packard Biosciences, Meriden, CT). In addition to enabling real-time target detection and quantification, this type of homogeneous assay format can be used to investigate cellular processes *in situ* in real time. This could be done by culturing cells directly onto a microwell plate and allowing uptake of scintillant beads and radioisotope by cells. Biosynthesis, proliferation, drug uptake, cell motility, etc. can then be monitored via the luminescence signal generated by beads in presence of selected target molecules (see Cook et al., 1992, or Heath et al., 1992).

Figures 11 A and B show an exemplary embodiment of a non-isotopic proximity assay based on nucleic acid sensor molecules used in conjunction with AlphaScreen™ beads (Packard Biosciences, Meriden, CT). In this embodiment, the nucleic acid sensor



molecules, which ligate on oligonucleotide substrate in the presence of a target molecule (see Figures 2A and B), are bound to a chemiluminescent compound-impregnated acceptor bead coated with, for example, streptavidin, via a (biotin) linker attached to the 5' end of the effector oligonucleotide sequence (GCGACTGGACATCACGAG (SEQ ID NO:51) in Figure 2A). The various bead-sensor coupling chemistries are determined by the type and manufacturer of the beads, and are well-known in the art. The oligonucleotide substrate is coupled to a photosensitizer-impregnated donor bead coated with, for example, streptavidin, via a (biotin) linker attached to the 3' end of the substrate. The donor (substrate) and acceptor (ribozyme) beads and target molecules are then combined in solution in a microwell plate, some of the NASMs hybridize and ligate the oligonucleotide substrate, bringing the donor and acceptor beads into close proximity (< 200 nm). Upon illumination at 680 nm, the photosensitizer in the donor bead converts ambient oxygen into the singlet state at a rate of approximately 60,000/second per bead. The singlet oxygen will diffuse a maximum distance of approximately 200 nm in solution; if an acceptor bead containing a chemiluminescent compound is within this range, *i.e.*, if ligation has occurred in the presence of the target molecule, chemiluminescence at 370 nm is generated. This radiation is immediately converted within the acceptor bead to visible luminescence at 520-620 nm with a decay half-life of 0.3 sec. The visible luminescence at 520-620 nm is detected using a time-resolved fluorescence/ luminescence plate reader (*e.g.*, the Fusion multifunction plate reader, Packard Biosciences, Meriden, CT). This type of nonisotopic homogeneous proximity assay format provides highly sensitive detection and quantification of target molecule concentrations in volumes < 25 microliters for high throughput screening (*see* Beaudet et al. 2001).

Suitable fluorescent labels are known in the art and commercially available from, for example, Molecular Probes (Eugene, Oreg.). These include, *e.g.*, donor/acceptor (*i.e.*, first and second signaling moieties) molecules such as: fluorescein isothiocyanate (FITC) /tetramethylrhodamine isothiocyanate (TRITC), FITC/Texas Red TM Molecular Probes), FITC/N-hydroxysuccinimidyl 1-pyrenebutyrate (PYB), FITC/eosin isothiocyanate (EITC), N-hydroxysuccinimidyl 1-pyrenesulfonate (PYS)/FITC, FITC/Rhodamine X (ROX), FITC/tetramethylrhodamine (TAMRA), and others. In addition to the organic fluorophores already mentioned, various types of nonorganic fluorescent labels are known in the art and are commercially available from, for example, Quantum Dot Corporation, Inc. Hayward CA). These include, *e.g.*, donor/ acceptor (*i.e.*, first and second signaling moieties) semiconductor nanocrystals (*i.e.*, 'quantum dots') whose absorption and emission spectra

can be precisely controlled through the selection of nanoparticle material, size, and composition (*see*, for example, Bruchez et al., 1998, Chan and Nie, 1998, Han et al., 2001).

The selection of a particular donor/acceptor pair is not critical to practicing the invention provided that energy can be transferred between the donor and the acceptor. P-  
5 (dimethyl aminophenylazo) benzoic acid (DABCYL) is one example of a non-fluorescent acceptor dye which effectively quenches fluorescence from an adjacent fluorophore, *e.g.*, fluorescein or 5-(2'-aminoethyl) aminonaphthalene (EDANS).

Figures 3A and B and 6A and B show exemplary optical nucleic acid sensor molecules derived from catalytic nucleic acid molecules (Figures 2A and B and 5,  
10 respectively), according to two embodiments. Figure 3 shows a catalytic nucleic acid sensor molecule obtained from an-oligonucleotide pool in which the catalytic site was a ligase site. Figure 6 shows a catalytic nucleic acid sensor molecule obtained from an oligonucleotide pool in which the catalytic site was a site mediating self-cleavage.

In the embodiment shown in Figures 3A and B, a catalytic nucleic acid sensor  
15 molecule from which a portion of a ligase site (*e.g.*, the AGUCG sequence at the 3' end of the nucleic acid sensor precursor molecule, as shown in Figure 2) has been removed is coupled to a first signaling moiety (F) at a first nucleotide (1) and to a second signaling moiety (D) at a second nucleotide (2). In a further embodiment, the first and second  
20 signaling moieties molecules are attached to non-terminal sequences. The position of the non-terminal sequences coupled to signaling moieties is limited to a maximal distance from the 5' or 3' nucleotide which still permits proximity dependent changes in the optical properties of the molecule. Coupling chemistries are routinely practiced in the art, and oligonucleotide synthesis services provided commercially (*e.g.*, Integrated DNA Technologies, Coralville, IA) can also be used to generate labeled molecules. In a further  
25 embodiment, the nucleic acid sensor molecule is used, either tethered to a solid support or free in solution, to detect the presence and concentration of target molecules in a complex biological fluid.

In the embodiment shown in Figures 3A and B, the first signaling moiety (F) is a fluorescein molecule coupled to the 5' end and the second signaling molecule (D) is a  
30 DABCYL molecule (a quenching group) coupled to the 3' end. Because of the nearly complete base pairing of the non-target molecule activated form (*see* Figure 3B), this is the favored form of the nucleic acid sensor molecule in the absence of the target molecule. When the nucleic acid sensor molecule is not activated by target molecule, the fluorescent group and the quenching group are in close proximity and little fluorescence is detectable

from the fluorescent group. Addition of target molecule causes a change in the conformation of the optical nucleic acid sensor molecule shown in Figure 3B to that shown in Figure 3A. When the molecule assumes the conformation shown in Figure 3A, the first and second signaling moieties (F and D, respectively) are no longer in sufficient proximity for the quenching group to quench the fluorescence of the fluorescent group, resulting in a detectable fluorescent signal being produced upon recognition of the target molecule.

In one embodiment, the target modulation domain sequence from a previously identified nucleic acid sensor molecule is incorporated into a separate oligonucleotide sequence which changes conformation upon target recognition as shown in Figures 6A and B. During or after synthesis, an optical signal generating unit is either added or inserted into the oligonucleotide sequence comprising the derived nucleic acid sensor molecule. As noted previously, nucleic acid sensor molecules of this type can be derived from allosteric ribozymes, such as those derived from the hammerhead, hairpin, L1 ligase, or group 1 intron ribozymes and the like (see Soukup et al., 2001, or Hamaguchi et al., 2001), all of which transduce molecular recognition into a detectable optical signal.

In the embodiment shown in Figures 9 A, B, and C, a self-cleaving ribozyme such as the hammerhead (in this case attached to a solid support via a linker molecule is shown) is labeled with a fluorophore. In Figure 9 A, the labeled NASM in the unactivated state comprises two oligonucleotide including a transacting cleavage substrate which bears a second fluorescent label. In the unactivated state, *i.e.*, in the absence of target molecule, the donor fluorophore (on one oligonucleotide NASM) and the acceptor fluorophore (on the cleavage substrate) are in sufficiently close proximity for FRET to occur; thus, minimal fluorescent emission is detected from the donor fluorophore at wavelength 3,  $\lambda_3$ , upon epillumination excitation at wavelength 1,  $\lambda_1$ . Upon target molecule recognition, the cleavage fragment of the cleavage substrate bearing the acceptor fluorophore dissociates from the ribozyme-target complex. Once separated from the acceptor fluorophore, the donor fluorophore can no longer undergo de-excitation via FRET, resulting in a detectable increase in its fluorescent emission at wavelength 2,  $\lambda_2$  (see, for example, Singh. et al., 1999; Wu, and Brand, 1994; Walter and Burke, 1997; Walter et al., 1998). In a further embodiment, the change in the polarization state of the fluorescent emission from the donor fluorophore (due to the increased diffusional rotation rate of the smaller cleavage fragment) can be detected/monitored in addition to changes in fluorescent emission intensity (see, for Singh, 2000). In a further embodiment, the NASMs are free in solution.

In another embodiment, shown in Figure 9 B, the acceptor fluorophore attached to the cleavage substrate is replaced by a quencher group. This replacement will also result in minimal fluorescent donor emission at wavelength 2,  $\lambda_2$ , when the NASM is in the unbound state under epi-illumination excitation at wavelength 1,  $\lambda_1$ . Upon target molecule  
5 recognition, the cleavage fragments of the cleavage substrate bearing the donor and quencher groups dissociate from the NASM-target molecule complex. Once separated from the quencher, the donor fluorophore will exhibit a detectable increase in its fluorescent emission at wavelength 2,  $\lambda_2$ . In a further embodiment, the change in the polarization state of the fluorescent emission from the donor fluorophore (due to the increased diffusional  
10 rotation rate of the smaller cleavage fragment) can be detected/monitored in addition to changes in fluorescent emission intensity. In a further embodiment, NASMs are free in solution.

In a different embodiment, the optical configuration is designed to provide excitation via total internal reflection (TIR)-illumination, as shown in Figure 9 C. Also, the  
15 donor fluorophore is attached to the NASM body while the quencher is attached to the cleavage substrate. In this configuration, with the surface-immobilized NASM in the unbound state, the fluorescent donor emission at wavelength 2,  $\lambda_2$ , will be minimal. Upon target molecule recognition, the cleavage fragment of the cleavage substrate bearing the quencher group dissociates from the NASM-target molecule complex. Once separated from  
20 the quencher, the donor fluorophore will exhibit a detectable increase in its fluorescent emission at wavelength 2,  $\lambda_2$ . In an alternative embodiment to that shown in shown in Figure 9 C, the quencher group can be replaced with an acceptor fluorophore. In yet another alternative embodiment to those shown in Figures 9 A, B, and C, the donor fluorophore is coupled to the cleavage fragment of the cleavage substrate and the acceptor fluorophore or  
25 quencher group is deleted. Upon target molecule recognition and dissociation of the cleavage fragment, the polarization state of the fluorescent emission from the donor fluorophore will undergo a detectable change due to the difference in the diffusional rotation rates of the surface-bound ribozyme-target complex and the free cleavage fragment.

In one embodiment, a universal FRET trans-substrate is synthesized for all NASMs  
30 derived from self-cleaving allosteric ribozymes. This substrate would have complementary optical signaling units (*i.e.*, donor and acceptor groups) coupled to opposite ends of the synthetic oligonucleotide sequence. Such a universal substrate would obviate the need for coupling optical signaling units to the sensor (*i.e.*, ribozyme) molecule itself.

The relative stabilities of the activated and unactivated forms of the nucleic acid sensor molecules is optimized to achieve the highest sensitivity of detection of target molecule. In one embodiment, the nucleic acid sensor molecule is further engineered to enhance the stability of one form over another. In one embodiment, the boxed UA in  
5 Figures 3A and B is changed to a CC, favoring the formation of the target molecule activated form. Because these bases do not form base pairs when the nucleic acid sensor molecule is unactivated, the unactivated form is not stabilized.

In addition to the herein described methods, any additional proximity dependent signaling system known in the art can be used to practice the method according to the  
10 invention, and are encompassed within the scope.

A number of methods can be used to evaluate the relative stability of different conformations of the nucleic acid sensor molecule. In one embodiment, the free energy of the structures formed by the nucleic acid sensor molecule is determined using software programs such as mfold®, which can be found on the Rensselaer Polytechnic Institute (RPI)  
15 web site ([www.rpi.edu/dept](http://www.rpi.edu/dept)).

In another embodiment, a gel assay is performed which permits detection of different conformations of the nucleic acid sensor molecule. In this embodiment, the nucleic acid sensor molecule is allowed to come to equilibrium at room temperature or the temperature at which the nucleic acid sensor molecule will be used. The molecule is then  
20 cooled to 4 °C and electrophoresed on a native (non-denaturing) gel at 4 °C. Each of the conformations formed by the nucleic acid sensor molecule will run at a different position on the gel, allowing visualization of the relative concentration of each conformation. Similarly, the conformation of nucleic acid sensor molecules which form in the presence of target molecule is then determined by a method such as circular dichroism (CD). By comparing  
25 the conformation of the nucleic acid sensor molecule formed in the presence of target molecule with the conformations formed in the absence of target molecule, the conformation which corresponds to the activated conformation can be identified in a sample in which there is no target molecule. The nucleic acid sensor molecule can then be engineered to minimize the formation of the activated conformation in the absence of target  
30 molecule. The sensitivity and specificity of nucleic acid sensor molecule can be further tested using target molecule modulation assays with known amounts of target molecules.

In another embodiment, shown in Figures 6 A and B, a catalytical nucleic acid sensor molecule from which a portion of a self-cleaving site has been removed, is coupled to a first signaling moiety (F) at a first nucleotide and to a second signaling moiety (D) at a

second nucleotide. In this embodiment, the entire catalytic site of the catalytic nucleic acid molecule (see Figure 5) has been removed. In one embodiment (Figures 6 A and B), additional bases (*e.g.*, UGGUAU) are added to one end of the portion of the nucleic acid sensor molecule comprising the modulation domain, to stabilize the unbound form of the nucleic acid sensor molecule (Figure 6B). These bases are selected to be complementary to bases at the opposite end of the nucleic acid sensor molecule (ACCAUA). Additional bases may be added to either the 5' or the 3' end of the nucleic acid sensor molecule.

Modifications to stabilize one conformation of the nucleic sensor molecule over another may be identified using the mfold program or native gel assays discussed above. A labeled nucleic acid sensor molecule is generated by coupling a first signaling moiety (F) to a first nucleotide and a second signaling moiety (D) to a second nucleotide as discussed above. As above, the sensitivity and specificity of the nucleic acid sensor molecule can be further assayed by using target molecule modulation assays with known amounts of target molecules.

15           ii.     Optical Signal Generating Units With Single Signaling Moieties

In one embodiment, the optical nucleic acid sensor molecule comprises an optical signaling unit with a single signaling moiety introduced at either an internal or terminal position within the nucleic acid sensor molecule. In this embodiment, binding of the target molecule results in changes in both the conformation and physical aspect (*e.g.*, molecular volume or mass, rotational diffusion rate, etc.) of the nucleic acid sensor molecule. Conformational changes in the nucleic acid sensor molecule upon target recognition will modify the chemical environment of the signaling moiety. Such a change in chemical environment will in general change the optical properties of the signaling moiety. Suitable signaling moieties are described in Jhaveri, et al, 2000, and include, *e.g.*, fluorescein, acridine, and other organic and nonorganic fluorophores.

In one embodiment, a signaling moiety is introduced at a position in the catalytic nucleic acid molecule near the target activation site (identifiable by footprinting studies, for example). Binding of the target molecule will (via a change in conformation of the nucleic acid molecule) alter the chemical environment and thus affect the optical properties of the signaling moiety in a detectable manner.

Recognition of the target molecule by the NASM will result in changes in the conformation and physical aspect of the nucleic acid sensor molecule, and will thus alter the kinetic properties of the signaling moiety. In particular, the changes in conformation and

mass of the sensor-target complex will reduce the rotational diffusion rate for the sensor-target complex, resulting in a detectable change in the observed steady state fluorescence polarization (FP) from the signaling moiety. The expected change in FP signal with target concentration can be derived using a modified form of the well-known Michaelis-Menten model for ligand binding kinetics (Lakowicz, 1999). FP is therefore a highly sensitive means of detecting and quantitatively determining the concentration of target molecules in a sample solution (Jameson and Sawyer, 1995; Jameson and Seifried, 1999; Jolley, 1999; Singh, 2000; Owicki et al., 1997). FP methods are capable of functioning in both solution- and solid-phase implementations.

Numerous additional methods can be used that, *e.g.*, make use of a single fluorescent label and an unpaired guanosine residue (instead of a quencher group), to enable the use of FRET in target detection and quantitation as described in the embodiments above (*see* Walter and Burke, 1997).

In a further embodiment, shown in Figure 10 A, B, and C an unlabeled ligating ribozyme such as the lysozyme-dependent L1 ligase is shown (*see*, for example, Robertson, M.P. and Ellington, A.D, 2000). In the unactivated state, *i.e.*, in the absence of target, no fluorescent emission is detected from the surface-bound ribozymes under total internal reflection (TIR)-illumination (*see* Figure 10 A), or epi-illumination (*see* Figure 10 B). Upon recognition of target molecules in the presence of an oligonucleotide substrate with a tag (where the tag is capable of binding to a subsequently added fluorescent label via interactions including, but not limited to, biotin/streptavidin, amine/aldehyde, hydrazide, thiol, or other reactive groups) those oligonucleotide substrates hybridized to NASMs will undergo ligation and become covalently bonded to the thereto. In order to maximize the probability of hybridization for a given NASM, oligonucleotide substrate can be added in excess relative to NASM, the temperature of the ambient solution in which the reaction takes place can be kept below room temperature (*e.g.*, 4 °C), and agitation of the reaction vessel can be employed to overcome the kinetic limitation of diffusion-limited transport of species in solution. Given the above conditions, as well as sufficient time for maximal hybridization and subsequent ligation to occur, fluorescent label with the appropriate reactive group to bind the substrate tag is added to the reaction mixture. Again, the degree of substrate-label binding can be maximized through control of label concentration, solution temperature, and agitation. Once the fluorescent label has bound to all available ligated substrate-ribozyme-target complex, the solution temperature can be raised to drive off all of the hybridized but unligated substrate. With TIR-illumination, the spatial extent of the

excitation region above the solid substrate surface to which the ribozymes are bound is only on the order of 100 nm. Therefore, the bulk solution above the substrate surface is not illuminated and the detected fluorescent emission will be primarily due to fluorophores which are bound to ligated oligonucleotide substrate-NASM- target molecule complexes tethered to the substrate surface. The fluorescence emission from surface-bound NASM- target molecule complexes in this homogeneous solid phase assay format represents an easily detectable optical signal. In another embodiment, the fluorescence polarization (FP) of the labeled substrate can be monitored, as shown in Figure 10 C. Upon ligation, the steady state fluorescence polarization signal from the substrate-NASM complex will increase detectably relative to the FP signal from the free labeled oligonucleotide substrate in solution, due to the difference in the diffusional rotation rates between the free and ligated forms.

In another embodiment, an unlabeled ligating ribozyme such as the lysozyme-dependent L1 ligase (see, for example, Robertson, M.P. and Ellington, A.D, 2000) is bound to a solid surface. In this embodiment, the oligonucleotide substrate is coupled to an enzyme-linked luminescent moiety, such as horse radish peroxidase (HRP) by a tag (where the tag is capable of binding to a subsequently added label via interactions including, but not limited to, biotin/streptavidin, amine/aldehyde, hydrazide, thiol, or other reactive groups). In the absence of target molecule, no luminescent emission is detected from the surface-bound ribozymes. Upon recognition of target molecules in the presence of labeled oligonucleotide substrate, those oligonucleotide substrates hybridized to NASMs will undergo ligation and become covalently bonded to the NASMs. After removal of excess, unbound oligonucleotide substrate, the activation empty for the enzyme-linked luminescent label is added to the reaction volume. The resulting luminescent signal (e.g., from HRP, luciferase, etc.) is easily detectable using standard luminometers (e.g., the Fusion multifunction plate reader, Packard Bioscience). In a further embodiment, the activated solution can be precipitated, followed by colorimetric detection. In a particular embodiment, the enzyme linked signal amplification, TSA, (sometimes referred to as CARD-catalyzed reporter deposition) is an ultrasensitive detection method. The technology uses turnover of multiple tyramide substrates per horseradish peroxidase (HRP) enzyme to generate high-density labeling of a target protein or nucleic acid probe *in situ*. Tyramide signal amplification is a combination of three elementary processes: (1) Ligation (or not) of a biotinylated ligase oligonucleotide substrate oligo, followed by binding (or not) of a streptavidin-HRP to the probe; (2) HRP-mediated conversion of multiple copies of a



fluorescent tyramide derivative to a highly reactive radical; (3) Covalent binding of the reactive, short lived tyramide radicals to nearby nucleophilic residues, greatly reducing diffusion-related signal loss.

## 5    2.    Generating Biosensors

Optical nucleic acid sensor molecules for the detection of a target molecule of interest are generated by first selecting catalytic nucleic acid molecules with catalytic activity modifiable (*e.g.*, activatable) by a selected target molecule. In one embodiment, at  
10    least a portion of the catalytic site of the catalytic NASM is then removed and an optical signal generating unit is either added or inserted. Recognition of the target molecule by the nucleic acid sensor molecule activates a change in the properties of the optical signaling unit.

In one embodiment, a biosensor is provided which comprises a plurality of optical  
15    nucleic acid sensor molecules labeled with first and second signaling moieties specific for a target molecule. In another embodiment, the optical NASMs are labeled with a single signaling moiety. In one embodiment, the labeled nucleic acid sensor molecules are provided in a solution (*e.g.*, a buffer). In another embodiment, the labeled nucleic acid sensor molecules are attached directly or indirectly (*e.g.*, through a linker molecule) to a  
20    substrate. In further embodiments, nucleic acid sensor molecules can be synthesized directly onto the substrate. Suitable substrates which are encompassed within the scope include, *e.g.*, glass or quartz, silicon, encapsulated or unencapsulated semiconductor nanocrystal materials (*e.g.*, CdSe), nitrocellulose, nylon, plastic, and other polymers. Substrates may assume a variety of configurations (*e.g.*, planar, slide shaped, wafers, chips, tubular, disc-  
25    like, beads, containers, or plates, such as microtiter plates, and other shapes).

Numerous attachment chemistries, both direct and indirect, can be used to immobilize the sensor molecules on a solid support. These include, *e.g.*, amine/aldehyde, biotin/streptavidin (avidin, neutravidin), ADH/oxidized 3' RNA. In a particular  
embodiment, the nucleic acid sensor molecules ligate a substrate in the presence of a target  
30    molecule (see Figures 2A and B). In this embodiment the ribozymes are bound to a solid substrate via the effector oligonucleotide sequence (for example, GCGACTGGACATCACGAG (SEQ ID NO:51) in Figure 2A).

In one embodiment, a manual or computer-controlled robotic microarrayer is used to generate arrays of nucleic acid sensor molecules immobilized on a solid substrate. In one

embodiment, the arrayer utilizes contact-printing technology (*i.e.*, it utilizes printing pins of metal, glass, etc., with or without quill-slots or other modifications). In a different embodiment, the arrayer utilizes non-contact printing technology (*i.e.*, it utilizes ink jet or capillary-based technologies, or other means of dispensing a solution containing the material to be arrayed). Numerous methods for preparing, processing, and analyzing microarrays are known in the art (*see* Schena et al., 2000; Mace et al., 2000; Heller et al., 1999; Basararsky et al., 2000; Schermer, 1999). Robotic and manual arrayers are commercially available for example, the SpotArray from Packard Biosciences, Meriden, CT, and the RA-1 from GenomicSolutions, Ann Arbor, MI).

10 In one embodiment, larger substrates can be generated by combining a plurality of smaller biosensors forming an array of biosensors. In a further embodiment, nucleic acid sensor molecules placed on the substrate are addressed (*e.g.*, by specific linker or effector oligonucleotide sequences on the nucleic acid sensor molecule) and information relating to the location of each nucleic acid sensor molecule and its target molecule specificity is stored within a processor. This technique is known as spatial addressing or spatial multiplexing. Techniques for addressing nucleic acids on substrates are known in the art and are described in, for example, U.S. Patent Number 6,060,252, U.S. Patent Number 6,051,380, U.S. Patent Number 5,763,263, U.S. Patent Number 5,763,175, and U.S. Patent Number 5,741,462.

20 In one embodiment different nucleic acid sensor molecules are immobilized on a streptavidin-derivatized glass substrate via biotin linkers. The individual sensor spots can be manually arrayed. Solution measurements of target molecule concentration can be made by bathing the surface of the biosensor array in a solution containing the targets (analytes) of interest. In practice this is accomplished either by incorporating the array within a microflowcell (with a flow rate of ~25 microliters/min), or by placing a small volume (~6-10 microliters) of the target solution on the array surface and covering it with a cover slip. Detection and quantification of target concentration is accomplished by monitoring changes in the fluorescence polarization (FP) signal emitted from the fluorescein label under illumination by 488 nm laser radiation. The rotational diffusion rate is inversely proportional to the molecular volume; thus the rotational correlation time for the roughly 20-nucleotide unbound sensor (*i.e.*, in the absence of target molecule) will be significantly less than that for the target-NASM complex. The fluorescence emission from the target-NASM complex will therefore experience greater residual polarization due to the smaller angle through which the emission dipole axis of the sensor fluorophore can rotate within its radiative lifetime. In another embodiment, different surface attachment chemistries are used

to immobilize the NASMs on a solid substrate. As previously noted, these include, *e.g.*, interactions involving biotin/streptavidin, amine/aldehyde, hydrazide, thiol, or other reactive groups.

The specificity of the biosensors and NASMs according to the invention is determined  
5 by the specificity of the target modulation domain of the nucleic acid sensor molecule. In one embodiment, a biosensor is provided in which all of the nucleic acid sensor molecules recognize the same molecule. In another embodiment, a biosensor is provided which can recognize at least two different target molecules allowing for multi-analyte detection. Multiple analytes can be distinguished by using different combinations of first and second  
10 signaling molecules. In addition to the wavelength/color and spatial multiplexing techniques previously described, biosensors may be used to detect multiple analytes using intensity multiplexing. This is accomplished by varying the number of fluorescent label molecules on each biosensor in a controlled fashion. Since a single fluorescent label is the smallest integral labeling unit possible, the number of fluorophores (*i.e.*, the intensity from) a given  
15 biosensor molecule provides a multiplexing index. Using the combination of 6-wavelength (color) and 10-level intensity multiplexing, implemented in the context of semiconductor nanocrystals derivatized as bioconjugates, would theoretically allow the encoding of million different analyte-specific biosensors (Han et al., 2001).

In one embodiment, multiple single target biosensors can be combined to form a  
20 multianalyte detection system which is either solution-based or substrate-based according to the needs of the user. In this embodiment, individual biosensors can be later removed from the system, if the user desires to return to a single analyte detection system (*e.g.*, using target molecules bound to supports, or, for example, manually removing a selected biosensor(s) in the case of substrate-based biosensors). In a further embodiment, nucleic  
25 acid sensor molecules binding to multiple analytes are distinguished from each other by referring to the address of the nucleic acid sensor molecule on a substrate and correlating its location with the appropriate target molecule to which it binds (previously described as spatial addressing or multiplexing).

In one embodiment, subsections of a biosensor array can be individually subjected to  
30 separate analyte solutions by use of substrate partitions or enclosures that prevent fluid flow between subarrays, and microfluidic pathways and injectors to introduce the different analyte solutions to the appropriate sensor subarray.

### 3. Nucleic acid sensor molecule and biosensor systems

In one embodiment, a nucleic acid sensor molecule or biosensor system is provided comprising a nucleic acid sensor molecule in communication with a detector system. In a further embodiment, a processor is provided to process optical signals detected by the detector system. In still a further embodiment, the processor is connectable to a server  
5 which is also connectable to other processors. In this embodiment, optical data obtained at a site where the NASM or biosensor system resides can be transmitted through the server and data is obtained, and a report displayed on the display of the off-site processor within seconds of the transmission of the optical data. In one embodiment, data from patients is stored in a database which can be accessed by a user of the system.

10 Data obtainable from the biosensors according to the invention include diagnostic data, data relating to lead compound development, and nucleic acid sensor molecule modeling data (*e.g.*, information correlating the sequence of individual sensor molecules with specificity for a particular target molecule). In one embodiment, these data are stored in a computer database. In a further embodiment, the database includes, along with  
15 diagnostic data obtained from a sample by the biosensor, information relating to a particular patient, such as medical history and billing information. Although, in one embodiment, the database is part of the nucleic acid sensor molecule system, the database can be used separately with other detection assay methods and drug development methods.

Detectors used with the nucleic acid sensor molecule systems according to the  
20 invention, can vary, and include any suitable detectors for detecting optical changes in nucleic acid molecules. These include, *e.g.*, photomultiplier tubes (PMTs), charge coupled devices (CCDs), intensified CCDs, and avalanche photodiodes (APDs). In one embodiment, a nucleic acid sensor molecule comprising labeled nucleic acid sensor molecules is excited by a light source in communication with the biosensor. In a further embodiment, when the  
25 optical signaling unit comprises first and second signal moieties that are donor/acceptor pairs (*i.e.*, signal generation relies on the fluorescence of a donor molecule when it is removed from the proximity of a quencher acceptor molecule), recognition of a target molecule will cause a large increase in fluorescence emission intensity over a low background signal level. The high signal-to-noise ratio permits small signals to be measured  
30 using high-gain detectors, such as PMTs or APDs. Using intensified CCDs, and PMTs, single molecule fluorescence measurements have been made by monitoring the fluorescence emission, and changes in fluorescence lifetime, from donor/acceptor FRET pairs (see Sako, et al., 2000; Lakowicz et al, 1991)).

Light sources include, *e.g.*, filtered, wide-spectrum light sources, (*e.g.*, tungsten, or xenon arc), laser light sources, such as gas lasers, solid state crystal lasers, semiconductor diode lasers (including multiple quantum well, distributed feedback, and vertical cavity surface emitting lasers (VCSELs)), dye lasers, metallic vapor lasers, free electron lasers, and lasers using any other substance as a gain medium. Common gas lasers include Argon-ion, Krypton-ion, and mixed gas (*e.g.*, Ar-Kr) ion lasers, emitting at 455, 458, 466, 476, 488, 496, 502, 514, and 528 nm (Ar ion); and 406, 413, 415, 468, 476, 482, 520, 531, 568, 647, and 676 nm (Kr ion). Also included in gas lasers are Helium Neon lasers emitting at 543, 594, 612, and 633 nm. Typical output lines from solid state crystal lasers include 532 nm (doubled Nd:YAG) and 408/816 nm (doubled/primary from Ti:Sapphire). Typical output lines from semiconductor diode lasers are 635, 650, 670, and 780 nm.

Excitation wavelengths and emission detection wavelengths will vary depending on the signaling moieties used. In one embodiment, where the first and second signaling moieties are fluorescein and DABCYL, the excitation wavelength is 488 nm and the emission wavelength is 514 nm. In the case of semiconductor nanocrystal-based fluorescent labels, a single excitation wavelength or broadband UV source may be used to excite several probes with widely spectrally separated emission wavelengths (*see* Bruchez et al., 1998; Chan et al., 1998).

In one embodiment, detection of changes in the optical properties of the nucleic acid sensor molecules is performed using any of a cooled CCD camera, a cooled intensified CCD camera, a single-photon-counting detector (*e.g.*, PMT or APD), or other light sensitive sensor. In one embodiment, the detector is optically coupled to the nucleic acid sensor molecule through a lens system, such as in an optical microscope (*e.g.*, a confocal microscope). In another embodiment, a fiber optic coupler is used, where the input to the optical fiber is placed in close proximity to the substrate surface of a biosensor, either above or below the substrate. In yet another embodiment, the optical fiber provides the substrate for the attachment of nucleic acid sensor molecules and the biosensor is an integral part of the optical fiber.

In one embodiment, the interior surface of a glass or plastic capillary tube provides the substrate for the attachment of nucleic acid sensor molecules. The capillary can be either circular or rectangular in cross-section, and of any dimension. The capillary section containing the biosensors can be integrated into a microfluidic liquid-handling system which can inject different wash, buffer, and analyte-containing solutions through the sensor tube. Spatial encoding of the sensors can be accomplished by patterning them longitudinally

along the axis of the tube, as well as radially, around the circumference of the tube interior. Excitation can be accomplished by coupling a laser source (*e.g.*, using a shaped output beam, such as from a VCSEL) into the glass or plastic layer forming the capillary tube. The coupled excitation light will undergo TIR at the interior surface/solution interface of the tube, thus selectively exciting fluorescently labeled biosensors attached to the tube walls, but not the bulk solution. In one embodiment, detection can be accomplished using a lens-coupled or proximity-coupled large area segmented (pixelated) detector, such as a CCD. In a particular embodiment, a scanning (*i.e.*, longitudinal/axial and azimuthal) microscope objective lens/emission filter combination is used to image the biosensor substrate onto a CCD detector. In a different embodiment, a high resolution CCD detector with an emission filter in front of it is placed in extremely close proximity to the capillary to allow direct imaging of the biosensors. In a different embodiment, highly efficient detection is accomplished using a mirrored tubular cavity that is elliptical in cross-section. The sensor tube is placed along one focal axis of the cavity, while a side-window PMT is placed along the other focal axis with an emission filter in front of it. Any light emitted from the biosensor tube in any direction will be collected by the cavity and focused onto the window of the PMT.

In still another embodiment, the optical properties of a nucleic acid sensor molecule are analyzed using a spectrometer (*e.g.*, such as a luminescence spectrometer) which is in communication with the biosensor. The spectrometer can perform wavelength discrimination for excitation and detection using either monochromators (*i.e.*, diffraction gratings), or wavelength bandpass filters. In this embodiment, biosensor molecules are excited at absorption maxima appropriate to the signal labeling moieties being used (*e.g.*, acridine at 450 nm, fluorescein at 495 nm) and fluorescence intensity is measured at emission wavelengths appropriate for the labeling moiety used (*e.g.*, acridine at 495 nm; fluorescein at 515 nm). Achieving sufficient spectral separation (*i.e.*, a large enough Stokes shift) between the excitation wavelength and the emission wavelength is critical to the ultimate limit of detection sensitivity. Given that the intensity of the excitation light is much greater than that of the emitted fluorescence, even a small fraction of the excitation light being detected or amplified by the detection system will obscure a weak biosensor fluorescence emission signal. In one embodiment, the biosensor molecules are in solution and are pipetted (either manually or robotically) into a cuvette or a well in a microtiter plate within the spectrometer. In a further embodiment, the spectrometer is a multifunction plate reader capable of detecting optical changes in fluorescence or luminescence intensity (at

one or more wavelengths), time-resolved fluorescence, fluorescence polarization (FP), absorbance (epi and transmitted), etc., such as the Fusion multifunction plate reader system (Packard Biosciences, Meriden, CT). Such a system can be used to detect optical changes in biosensors either in solution, bound to the surface of microwells in plates, or immobilized  
5 on the surface of solid substrate (e.g., a biosensor microarray on a glass substrate). This type of multiplate/multisubstrate detection system, coupled with robotic liquid handling and sample manipulation, is particularly amenable to high-throughput, low-volume assay formats.

In embodiments where nucleic acid sensor molecules are attached to substrates,  
10 such as a glass slide or in microarray format, it is desirable to reject any stray or background light in order to permit the detection of very low intensity fluorescence signals. In one embodiment, a small sample volume (~10 nL) is probed to obtain spatial discrimination by using an appropriate optical configuration, such as evanescent excitation or confocal imaging. Furthermore, background light can be minimized by the use of narrow-bandpass  
15 wavelength filters between the sample and the detector and by using opaque shielding to remove any ambient light from the measurement system.

In one embodiment, spatial discrimination of nucleic acid sensor molecules attached to a substrate in a direction normal to the interface of the substrate (i.e., excitation of only a small thickness of the solution layer directly above and surrounding the plane of attachment  
20 of the biosensor molecules to the substrate surface) is obtained by evanescent wave excitation. This is illustrated in Figure 12. Evanescent wave excitation utilizes electromagnetic energy that propagates into the lower-index of refraction medium when an electromagnetic wave is totally internally reflected at the interface between higher and lower-refractive index materials. In this embodiment a collimated laser beam is incident on  
25 the substrate/solution interface (at which the biosensors are immobilized) at an angle greater than the critical angle for total internal reflection (TIR). This can be accomplished by directing light into a suitably shaped prism or an optical fiber. In the case of a prism, as shown in Figure 12, the substrate is optically coupled (via index-matching fluid) to the upper surface of the prism, such that TIR occurs at the substrate/solution interface on which  
30 the biosensors are immobilized. Using this method, excitation can be localized to within a few hundred nanometers of the substrate/solution interface, thus eliminating autofluorescence background from the bulk analyte solution, optics, or substrate. Target recognition is detected by a change in the fluorescent emission of the nucleic acid sensor,

whether a change in intensity or polarization. Spatial discrimination in the plane of the interface (*i.e.*, laterally) is achieved by the optical system.

In one embodiment, a large area of the biosensor substrate is uniformly illuminated, either via evanescent wave excitation or epi-illumination from above, and the detected  
5 signal is spatially encoded through the use of a pixelated detector, such as CCD camera. An example of this type of uniform illumination/CCD detection system (using epi-illumination)) for the case of microarrayed biosensors on solid substrates is the GeneTAC 2000 scanner (GenomicSolutions, Ann Arbor, MI). In a different embodiment, a small area (*e.g.*, 10 x 10 microns to 100 x 100 microns) of the biosensor substrate is illuminated by a  
10 micro-collimated beam or focused spot. In one embodiment, the excitation spot is rastered in a 2-dimensional scan across the static biosensor substrate surface and the signal detected (with an integrating detector, such as a PMT) at each point correlated with the spatial location of that point on the biosensor substrate (*e.g.*, by the mechanical positioning system responsible for scanning the excitation spot). Two examples of this type of moving spot  
15 detection system for the case of microarrayed biosensors on solid substrates are: the DNAScope scanner (confocal, epi-illumination, GeneFocus, Waterloo, ON, Canada), and the LS IV scanner (non-confocal, epi-illumination, GenomicSolutions, Ann Arbor, MI). In yet another embodiment, a small area (*e.g.*, 10 x 10 microns to 100 x 100 microns) of the biosensor substrate is illuminated by a stationary micro-collimated beam or focused spot,  
20 and the biosensor substrate is rastered in a 2-dimensional scan beneath the static excitation spot, with the signal detected (with an integrating detector, such as a PMT) at each point correlated with the spatial location of that point on the biosensor substrate (*e.g.*, by the mechanical positioning system responsible for scanning the substrate). An example of this type of moving substrate detection (using confocal epi-illumination) system for the case of  
25 microarrayed biosensors on solid substrates is the ScanArray 5000 scanner (Packard Biochip, Billerica, MA).

In the embodiment shown in Figure 12, a TIR evanescent wave excitation optical configuration is implemented, with a static substrate and dual-capability detection system. The detection system is built on the frame of a Zeiss universal fluorescence microscope.  
30 The system is equipped with 2 PMTs on one optical port, and an intensified CCD camera (Cooke, St. Louis, MO) mounted on the other optical port. The optical path utilizes a moveable mirror which can direct the collimated, polarized laser beam through focusing optics to form a spot, or a beam expander to form a large (> 1cm) beam whose central portion is roughly uniform over the field of view of the objective lens. Another movable



mirror can direct the light either to the intensified CCD camera when using large area uniform illumination, or to the PMTs in the scanned spot mode. In spot scanning mode, a polarizing beamsplitter separates the parallel and perpendicular components of the emitted fluorescence and directs each to its designated PMT. An emission filter in the optical column rejects scattered excitation light from either type of detector. In CCD imaging mode, manually adjusted polarizers in the optical column of the microscope must be adjusted to obtain parallel and perpendicular images from which the fluorescence polarization or anisotropy can be calculated. A software program interfaces with data acquisition boards in a computer which acquires the digital output data from both PMTs and CCD. This program also controls the PMT power, electromechanical shutters, and galvanometer mirror scanner, calculates and plots fluorescence polarization in real time, and displays FP and intensity images.

In another embodiment, the detection system is a single photon counter system (*see, e.g.,* U.S. Patent Number 6,016,195 and U.S. Patent Number 5,866,348) requiring rastering of the sensor substrate to image larger areas and survey the different binding regions on the biosensor.

In another embodiment of the invention, the biosensor is used to detect a target molecule through changes in the electrochemical properties of the nucleic acid sensor molecule or molecules in close proximity to it which occur upon recognition of the NASM to the target molecule.

In a preferred embodiment, the biosensor system would consist of three major components: One, optical nucleic acid sensor molecules immobilized on an array of independently addressable gold electrodes. The nucleic acid sensor molecules immobilized on each electrode may be modulated by the same or different target molecules, including proteins, metabolites and other small molecules, etc.; two, an oligonucleotide substrate which acts as a signaling probe, hybridizing to the oligonucleotide substrate binding domain of the ligase sensor and forming a covalent phosphodiester bond with the nucleic acid sensor molecule nucleotide adjacent to its 3' terminus in the presence of the appropriate target. This oligonucleotide substrate is typically a nucleic acid sequence containing one or more modified nucleotides conjugated to redox active metallic complexes, *e.g.,* ferrocene moieties, which can act as electron donors; and three, an immobilized mixed self-assembled surface monolayer (SAM), comprised of conductive species separated by insulating species, covering the surface of the electrodes. Examples of conductive species include thiol-terminated linear molecules, such as oligophenylethyl molecules, while examples of

nonconductive thiol-terminated linear molecules, include alkane-thiol molecules terminated with polyethylene glycol (PEG). All immobilized species can be covalently attached to the electrode surface by terminal thiol groups. Figure 15 schematically shows the structure of the mixed self-assembled surface monolayer (SAM) coating the gold electrode, as well as the immobilized nucleic acid sensor molecule (NASM) with a ligated oligonucleotide substrate conjugated to several redox active moieties. Upon recognition of the target molecule by the target modulation domain and subsequent ligation of the oligonucleotide substrate at the site indicated in the figure, the redox active signaling moieties coupled to the substrate oligo will be brought into close proximity to the conductive surface layer, resulting in a detectable increase in electronic surface signal.

In another preferred embodiment, the biosensor system would consist of two major components: (1) Optical nucleic acid sensor molecules immobilized on an array of independent addressable gold electrodes. The nucleic acid sensor molecules immobilized on each electrode may be modulated by the same or different target molecules, including proteins, metabolites and other small molecules, etc. The NASM will contain one or more nucleotides conjugated to redox active metallic complexes, *e.g.*, ferrocene moieties, which can act as electron donors; and (2) an immobilized mixed self-assembled surface monolayer (SAM), comprised of conductive species separated by insulating species, covering the surface of the electrodes. Examples of conductive species include thiol-terminated linear molecules, such as oligophenylethynyl molecules, while examples of nonconductive thiol-terminated linear molecules include alkane-thiol molecules terminated with polyethylene glycol (PEG). Figure 16 shows the SAM-coated molecule immobilized via a capture oligonucleotide. In this case, the redox active signaling moieties are coupled to the body of the NASM, as shown in the figure. Upon recognition of the target molecule by the target modulation domain and subsequent cleavage at the site indicated in the figure, the bulk of the NASM, including the nucleotides coupled to the redox active signaling moieties will dissociate from the surface, resulting in a detectable loss of electronic current signal.

In another embodiment, the array would be subjected, *e.g.*, by an integrated microfluidic flowcell, to an analyte solution containing the target(s) of interest at some unknown concentration. The range of possible sample analyte solutions may include standard buffers, biological fluids, and cell or tissue extracts. The sample solution will also contain the signaling probe at a saturating concentration relative to the immobilized nucleic acid sensor molecule. This ensures that at any given time during analysis, there is a high probability that each nucleic acid sensor molecule will have a signaling probe hybridized to

it. In the presence of the target molecules in the sample solution, the nucleic acid sensor molecule will form a covalent phosphodiester bond, *i.e.*, ligate, with the signaling probe, thus immobilizing it with its redox active electron donor species in electrical contact with the conductive molecules within the mixed self-assembled surface monolayer. After some  
5 integration time, during which signal probe ligation occurs, it may be necessary to denature the hybridized but unligated signaling probes. This denaturation step, which effectively removes 'background' signaling probes and their associated redox moieties from the vicinity of the electrode, can be accomplished by a small temperature increase (*e.g.*, from 21 °C to 25 °C), or by a brief negative voltage spike applied to the sensor electrodes followed  
10 by the application of a large positive DC voltage to a separate electrode that would collect unligated signaling. For the case of a sufficiently short hybridization region, *e.g.*, 5 base-pairs, on the signaling probe, a separate denaturation step may not be necessary. In either case, following nucleic acid sensor molecule activation by target molecules, a linear electrical potential ramp is applied to the electrodes. The redox species conjugated to the  
15 immobilized signaling probe-nucleic acid sensor molecule will be electrochemically oxidized, liberating one or more electrons per moiety. The conductive molecules within the surface monolayer will provide an electrical path for the liberated electrons to the electrode surface, as shown in Figure 17.

The net electron transfer to or from the electrode will be measured as a peak in the  
20 faradaic current, centered at the redox potential of the electron donor species (specified for a given reference electrode) and superposed on top of the capacitive current baseline which is observed in the absence of surface-immobilized signaling probes. This is shown schematically in Figure 18.

Quantitative analysis of the sensor signal, and therefore accurate determination of  
25 target molecule concentration, is based on the fact that the measured faradaic peak height is directly proportional to number of redox moieties immobilized at the electrode, that is, the number of nucleic acid sensor molecules ligated to signaling probes times the multiplicity of redox moieties per signaling probe molecule. Signal generation by the nucleic acid sensor molecules is thus amplified by virtue of multiple redox species per signaling probe. In  
30 addition, if an alternating current (AC) bias voltage is applied (superposed) on top of the DC linear voltage ramp applied to the sensor electrodes, *i.e.*, in the case of AC voltammetry, signal amplification would result from the cyclic repetition of the signal-generating redox reaction.

The system described above for the case of a surface-immobilized nucleic acid sensor molecule which ligates a signaling probe containing one or more modified nucleotides conjugated to redox active species suggests a general method and instrumentation for the detection and quantitation of an arbitrary target molecule in solution in real time. Detection of a particular target would require development of a nucleic acid sensor molecule against that recognizes the target molecule. Additionally, nucleic acid sensor molecules have been developed which are activated only in the presence of two different target molecules. Such dual-effector sensors could be used to detect the simultaneous presence of two or more targets, or could be used in conjunction with single-target molecule sensors to form biological logic (*i.e.*, AND, OR, etc.) circuits.

Multiplexed detection of multiple target molecules simultaneously in a complex sample solution could be accomplished by immobilizing nucleic acid sensor molecules against the target molecules of interest on separate electrodes within a two-dimensional array of electrodes. A complex sample solution containing multiple target molecules and a common signaling probe could then be introduced to the array. All nucleic acid sensor molecules would be exposed simultaneously to all targets, with the target-activated nucleic acid sensor molecule response(s) being observed and recorded only at the spatial location(s) known to contain a nucleic acid sensor molecule specific for the target molecules present in the (unknown) sample. The utility of such a nucleic acid sensor molecule array would be greatly enhanced by the integration of a microfluidic sample and reagent delivery system. Such an integrated microfluidic system would allow the application of reagents and samples to the sensor array to be automated, and would allow the reduction of sample volume required for analysis to < 1 uL.

The sensor array electrodes may be of any configuration, number, and size. In a preferred embodiment, the sensor and reference electrodes would be circular gold pads on the order of 100-500 uM in diameter, separated by a center-to center distance equal to twice their diameter. Each electrode would be addressed by separate electrical interconnects. The application of electrical signals to the sensor electrodes can be accomplished using standard commercially available AC and DC voltage sources. Detection of faradaic electrical signals from the sensor electrodes can be accomplished easily using standard commercially available data acquisition boards mounted within and controlled by a microcomputer. Specifically, the raw sensor current signals would need to be amplified, and then converted to a voltage and analyzed via a high resolution (*i.e.*, 16 bit) analog to digital converter (ADC). It is possible to reduce the signal background and to increase the signal to noise

ratio (SNR) by using the common technique of phase-sensitive detection. In this detection method, an alternating current (AC) bias voltage (at a frequency between, for example, 100 to 1000 Hz) is superposed on top of the DC linear voltage ramp applied to the sensor electrodes. The frequency of the applied bias voltage is called the fundamental frequency. It can be shown that the sensor response signal contains multiple frequency components, including the fundamental frequency and its harmonics (integral multiples of the fundamental frequency). It can further be shown that the *n*th harmonic signal is proportional to the *n*th derivative of the signal. Detecting these derivative signals (by means of a lock-in amplifier) minimizes the effects of constant or sloping backgrounds, and can enhance sensitivity by increasing the signal to noise ratio and allowing the separation of closely spaced signal peaks. It should be noted that digital, computer-controlled AC and DC voltage sources (*i.e.*, digital to analog converters, DACs), current preamplifiers, analog to digital converters (ADCs), and lock-in amplifiers are all available as integrated signal generation/acquisition boards that can be mounted within and controlled by a single microcomputer.

In a preferred embodiment, an integrated nucleic acid sensor molecule system with electrochemical detection would include the following elements one, a independently addressable multielement electrode array with immobilized surface layer composed of conductive species separated by insulating species and sensors; two, optical nucleic acid sensor molecules immobilized on the electrode array; three, an oligonucleotide substrate/signaling probe which ligates with the nucleic acid sensor molecule in the presence of the appropriate target; four, an automated or semi-automated microfluidic reagent and sample delivery system; and five, a reader instrument/data acquisition system consisting of a microcomputer controlling the appropriate voltage sources, current and lock-in amplifiers, data acquisition boards, and software interface for instrument control and data collection.

In another embodiment, the change in activity of the nucleic acid sensor molecule can be detected by watching the change in fluorescence of a nucleic acid sensor molecule when it is immobilized on a chip. A ligase can be attached to a chip and its ligase activity monitored. Ligase ERK nucleic acid sensor molecules, labeled with one fluorophore, *e.g.*, Cy3 is attached via an amino modification to an aldehyde chip. The initial Cy3 fluorescence indicates the efficiency of immobilization of the nucleic acid sensor molecules. Next, the chip is exposed to a substrate labeled with a second fluorophore, *e.g.*, Cy5, with or without the ERK protein target. In the presence of ERK the nucleic acid sensor molecule

ligates the substrate to itself, and becomes Cy5-labeled. Without ERK, the ligation does not occur.

The use of a labeled effector oligonucleotide does not change the rate of ligation of the nucleic acid sensor molecule whether ERK is present or not. When using nucleic acid sensor molecules in the context of a chip based system, in one embodiment, an effector oligonucleotide is used to attach the nucleic acid sensor molecule to the chip. Whether one uses effector oligonucleotide or not, the TaqMan (real-time PCR) traces obtained in the presence or absence of target (ERK) for ERK dependent ligases are identical. The presence or absence of this effector oligonucleotide does not affect the activity of the nucleic acid sensor molecule.

In another embodiment, a hammerhead nucleic acid sensor molecule could be used to measure the concentration of an analyte through the use of fluorescence. Figure 82 shows how many nucleic acid sensor molecules with different effector molecules and/or analytes could be integrated onto one chip to study the concentration of many molecules at once.

Any optical method known in the art, in addition to those described above can be used in the detection and/or quantification of all targets of interest in all sensor formats, in both biological and nonbiological media. These targets include, *e.g.*, those listed in Table 1, below.

Any other detection method can also be used in the detection and/or quantification of targets. For example, radioactive labels could be used, including  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{14}\text{C}$ ,  $^3\text{H}$ , or  $^{125}\text{I}$ . Also enzymatic labels can be used including horse radish peroxidase or alkaline phosphatase. The detection method could also involve the use of a capture tag for the bound nucleic acid sensor molecule.

#### 4. Methods of Using Nucleic Acid Sensor Molecules

##### A. Diagnostic Assays

The nucleic acid sensor molecules according to the invention can be used to detect virtually any target molecule. In one embodiment, the target molecule is a target molecule associated with a pathological condition and detection of changes in the optical properties of the nucleic acid sensor molecules of the biosensor provides a means of diagnosing the condition. Target molecules which are contemplated within the scope include, *e.g.* proteins, modified forms of proteins, metabolites, organic molecules, and metal ions, as discussed

above. Because signal generation in this system is reversible, washing of the biosensor(s) in a suitable buffer will allow the biosensor(s) to be used multiple times, enhancing the reproducibility of the any diagnostic assay since the same reagents can be used over and over. Suitable wash buffers include, e.g., binding buffer without target or, for faster  
5 washing, a high salt buffer or other denaturing conditions, followed by re-equilibration with binding buffer.

Re-use of the biosensor is enhanced by selecting optimal fluorophores. For example, Alexa Fluor 488, produced by Molecular Probes, has similar optical characteristics compared to fluorescein, but has a much longer lifetime. However, in one embodiment, a  
10 site recognized by a nuclease is engineered proximal to the signal generating site, and sequences comprising signaling moieties are removed from the biosensor and replaced by new sequences, as needed.

#### Profiling Biosensors for Use in Diagnostic Assays

In one embodiment, the expression pattern of a plurality of target molecules is  
15 determined to obtain a profile of target molecules associated with a trait in an individual to determine an expression pattern which is diagnostic of that trait. In this embodiment, combinations of biosensors targeted to individual target molecules are selected until a signature optical profile is determined which is characteristic of a trait. Traits include, e.g., a disease, a genetic alteration, a combination of genetic alterations (e.g., a polygenic  
20 disorder), a physiological reaction to an environmental condition, or a wild type state (e.g., of an organism or of an organ system). The target molecules which generate the signature optical profile are identified (based on the type of biosensors used) as signature target molecules. The expression of the signature target molecules can thereafter be determined to identify the presence of the trait in a patient.

25 The expression of the target molecules can be identified using any molecular detection system known in the art; however, in a preferred embodiment, the detection system comprises optical nucleic acid sensor molecules and the trait is identified by detecting the signature optical profile. In one embodiment, data relating to the signature optical profile is stored in the memory of a computer. Signature optical profiles can be  
30 generated for individual patients or can be generated for populations of individuals. In the latter embodiment, data relating to a composite signature profile (e.g., comprising normalized data) is stored in the memory of a computer or in a computer program product.

Because the biological function of the target molecules does not need to be known, biosensors according to the invention can be generated which are diagnostic of diseases/traits whose biological basis is not yet known or are the result of complex polygenic interactions and/or of environmental influences. In one embodiment, nucleic acid sensor molecules are identified which are activatable by synthetic polypeptides obtained from putative open reading frames identified in the human genome project and/or in other sequencing efforts. Combinations of these activatable nucleic acid sensor molecules (along with activatable nucleic acid sensor molecules specific for target molecules with known functions) are identified which generate a diagnostic optical signal, and signature target molecules are in turn identified which are linked to a particular trait, allowing a biological activity to be associated with a previously uncharacterized molecule.

Data relating to signature target molecules or to the optical signals generated upon activation of nucleic acid sensor molecules upon binding to signature target molecules is stored in a database, which can include further information such as sequence information or chemical structure information relating to the signature target molecule. A signature profile relating to a particular trait is generated based on normalized data from a plurality of tests. In one embodiment, a signature profile is obtained by determining any or all of the level, chemical structure, or activity, of signature target molecules associated with a disease in samples from a population of healthy individuals to determine a signature profile corresponding to a healthy state. In a further embodiment, signature profiles are obtained using data from subsets of populations which are divided into groups based on sex, age, exposure to environmental factors, ethnic background, and family history of a disease.

#### **B. Drug Discovery**

Generally, methods of drug discovery comprise steps of 1) identifying target(s) molecules associated with a disease; 2) validating target molecules (e.g., mimicking the disease in an animal or cellular model); 3) developing assays to identify lead compounds which affect that target (e.g., such as using libraries to assay the ability of a compound to bind to the target); 4) prioritizing and modifying lead compounds identified through biochemical and cellular testing; 5) testing in animal models; and 6) testing in humans (clinical trials). Through the power of genomics and combinatorial chemistry, large numbers of lead compounds can be identified in high throughput assays (step 3); however, a bottleneck occurs at step 4 because of the lack of efficient ways to prioritize and optimize lead compounds and to identify those which actually offer potential for clinical trials. The



target activatable nucleic acid sensor molecules according to the present invention offer a way to solve this problem by providing reagents which can be used at each step of the drug development process. Most importantly, the target activatable nucleic acid sensor molecules according to the present invention offer a way to correlate biochemical data, from *in vitro* 5 biochemistry and cellular assays, with the effect of a drug on physiological response from a biological assay.

In one embodiment of invention, a method for identifying a drug compound is provided, comprising identifying a profile of target molecules associated with a disease trait in a patient, administering a candidate compound to the patient, and monitoring changes in 10 the profile. In another embodiment, the monitored profile is compared with a profile of a healthy patient or population of healthy patients, and a compound which generates a profile which is substantially similar to the profile of target molecules in the healthy patient(s) (based on routine statistical testing) is identified as a drug. In a further embodiment, both the profiling and the drug identification step is performed using at least one nucleic acid 15 sensor molecule whose properties change upon binding to a target molecule.

In a further embodiment, a method for identifying a drug compound comprises identifying a plurality of pathway target molecules, each belonging to a pathway, monitoring the level, chemical structure, and/or activity of pathway target molecules in a patient having a disease trait, administering a candidate compound to the patient, and 20 monitoring changes in the level, chemical structure, and/or activity of the pathway target molecules. In another embodiment, the monitored level, chemical structure, and/or activity of the pathway target molecules is compared to the level, chemical structure, and/or activity of pathway target molecules in a wild type patient or patients. In a further embodiment, both profiling and the identification of drug compounds is performed using at least one nucleic 25 acid sensor molecule whose properties change upon binding to a pathway target molecule.

Properties according to this aspect include, *e.g.*, optical properties, change in sequence, chemical structure, catalytic activity, and/or molecular weight. In a preferred embodiment, sensor molecules are target activated optical nucleic acid sensor molecules.

i. Nucleic acid sensor molecules for Use in Identifying Lead Compounds

30 In one embodiment, biosensors activatable by signature target molecules, identified as described above, are provided and are validated by testing against multiple patient samples *in vitro* to verify that the optical signal generated by these molecules is diagnostic of a particular disease. Validation can also be performed *ex vivo*, *e.g.*, in cell culture, (using

microscope-based detection systems and other optical systems as described in U.S. Patent Number 5,843,658, U.S. Patent Number 5,776,782, U.S. Patent Number 5,648,269, and U.S. Patent Number 5,585,245) and/or *in vivo*, for example, by providing a profile biosensor in communication with an optical fiber.

5       The incorporation of biosensors into fiber optic waveguides is known in the art (see, e.g., U.S. Patent Number 4,577,109, U.S. Patent Number 5,037,615, U.S. Patent Number 4,929,561, U.S. Patent Number 4,822,746, and U.S. Patent Number 4,762,799). The selection of fluorescent energy transfer molecules for *in vivo* use is described in EP-A 649848, for example. In this embodiment, nucleic-acid based biosensors are introduced into  
10 the body by any suitable medical access device, such as an endoscope or a catheter. The optical fiber is provided within a working lumen of the access device and is in communication with an optical imaging system.

In one embodiment, the same methods which are used to validate the diagnostic value of particular sets of target molecule/nucleic acid sensor molecule combinations are  
15 used to identify lead compounds which can function as drugs. Thus, in one embodiment, the effects of a compound on target dependent optical signaling is monitored to identify changes in a signature profile arising as a result of treatment with a candidate compound.

In one embodiment, samples from a treated patient are tested *in vitro*; however, samples can also be tested *ex vivo* or *in vivo*. When the diagnostic profile identified by the  
20 biosensor changes from a profile which is a signature of a disease to one which is substantially similar to the signature of a wild type state (e.g., as determined using routine statistical tests), the lead compound is identified as a drug. Target molecules which activate the biosensor can comprise molecules with characterized activity and/or molecules with uncharacterized activity. Because large number of target molecules can be monitored  
25 simultaneously, the method provides a way to assess the affects of compounds on multiple drug targets simultaneously, allowing identification of the most sensitive drug targets associated with a particular trait (e.g., a disease or a genetic alteration).

Examples of suitable target molecules include, e.g., nuclear hormone receptor (NHR) polypeptides; G-coupled protein receptor (GPCR) polypeptides, phosphodiesterase  
30 (PDE), and protein kinases.

*NHR polypeptides*

Included in the invention are methods of identifying nucleic acid sensor molecules for detection of conformational isoforms of nuclear hormone receptors, as well as the nucleic acid sensor molecules identified by the methods described herein.

Nuclear hormone receptors (NHRs) act as ligand-inducible transcription factors by  
5 directly interacting as monomers, homodimers, or heterodimers in complex with DNA response elements of target genes. The activation of these transcription regulators is induced by the change in conformation of the NHR upon complex formation with ligand.

Provided are methods for generating unique NASMs for each NHR ligand binding domain (LBD). The NASMs described herein can include, *e.g.*, those derived from the  
10 hammerhead, hairpin, L1 ligase or group I intron ribozymes and the like, any of which transduce molecular recognition into a detectable signal.

Also provided is a direct mechanistic assay for the action of small molecule ligand-agonism, -antagonism and partial antagonism of members of the NHR family. The mechanistic assays function in both *in vitro* biochemical as well as with *in vitro* cell-based  
15 settings. In the *in vitro* assay setting, the nucleic acid sensor molecules are designed to recognize one conformational isomer of the NHR. In one embodiment, the nucleic acid sensor molecule recognizes the unique conformation that exists for the agonist bound form of a hormone receptor; such as that observed for the estrogen receptor ligand binding domain ER-LBD when bound to estrogen (Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. Cell. 1998;95(7):927-37) and then produces a  
20 detectable signal, such as release of fluorescently labeled oligonucleotide, radiolabeled oligonucleotide, or reveals a change in nucleic acid sensor molecule conformation driven by ligand binding through a change in fluorescence or the like. Hence, in this embodiment, the nucleic acid sensor molecule transduces molecular recognition of the ER-LBD-estrogen agonist complex into a detectable signal. The level of the signal is then used to quantify the amount of ER-LBD-estrogen agonist complex present in solution. In another embodiment, the ER-LBD-estrogen specific nucleic acid sensor molecule is used as a screening tool in assays designed to detect inhibitors of ER-LBD-estrogen complex formation. These screening tools can be used to determine the inhibition potency of compounds in *in vitro*  
25 biochemical assays or in *in vitro* cell-based assays. Inhibitors of estrogen binding to ER-LBD are useful as anti-proliferative agents for treatment of breast cancers (*e.g.*, tamoxifen) and other estrogen dependent diseases. In another embodiment, nucleic acid sensor molecules are introduced into cell lines by known methods of electroporation, transfection or coupling to peptide translocating agents such as tat or antennapedia peptides. In another  
30

embodiment, the ER-LBD-estrogen complex specific nucleic acid sensor molecule is an allosteric intron imbedded in a reporter gene such as GFP or luciferase or the like. When the intron derived nucleic acid sensor molecule is inserted into the reporter gene it renders reporter gene expression effector dependent. Thus, in one embodiment functional GFP  
5 protein is expressed only when the ER-LBD-estrogen complex is present in the cell, and inhibitors of ER-LBD-estrogen complex formation thus block functional GFP protein expression in appropriate mammalian, such as MCF7 or T47D, yeast or bacterial cell lines. In a further embodiment, the MCF7 or T47D tumor cell lines transfected with GFP-ER-LBD-estrogen nucleic acid sensor molecule sensitive construct are used to form tumor  
10 xenografts in nude mice. Thus, the transfected tumor xenograft cell lines can be used to form tumors in mice which are not only estrogen dependent but also regulate reporter gene expression in ER-LBD-estrogen dependent manner. These cell lines and tumor models are used to discover inhibitors of ER-LBD-estrogen complex formation *in vivo*.

NHR ligand binding domains bind antagonists, forming additional conformational  
15 isomers. When antagonists are bound to the receptor a new conformer results such as that observed upon tamoxifen binding to the estrogen receptor to form a stable ER-LBD-tamoxifen complex (Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. Cell. 1998 Dec 23;95(7):927-37). Accordingly, the invention includes use of an ER-LBD-tamoxifen specific nucleic acid sensor molecule that is used to detect the levels  
20 of antagonist specific complex in both *in vitro* biochemical, cell-based and, in *in vivo* assays as described above.

Nucleic acid sensor molecules can be developed that are specific for the ligand binding domains of all NHRs. In addition, it should be clear that nucleic acid sensor molecules for agonist, antagonist, dimeric or multimeric forms of NHR LBDs can be used  
25 to screen for inhibitors of LBD function and therefore for inhibitors of NHR dependent transcriptional activation or repression. It should be clear to one skilled in the art that nucleic acid sensor molecules specific for individual LBD complexes can be used to screen for agents that modify NHR function in *in vitro* and in *in vivo* assays.

NHRs are multidomain proteins containing a variable NH<sub>2</sub>-terminal region (A/B), a  
30 conserved DNA binding domain (DBD) or region (C), a linker region (D), and a conserved region (E) that contains the ligand binding domain (LBD). NHRs also contain regions required for transcriptional activation, of particular interest is the region AF-2 which is located in the COOH-terminus and whose function is strictly ligand dependent. Provided herein is a method for generating unique nucleic acid sensor molecules to each of the 63

known human NHR LBDs. In addition, methods are described that enable the generation of nucleic acid sensor molecules capable of recognizing the activated state of the NHR by selection for nucleic acid sensor molecules geometries which signal the presence of either the activated or inactivated conformation (NHR with bound ligand), but whose signaling  
5 action is quiescent in the presence of other forms of the NHR.

The nucleic acid sensor molecules allow the direct, simultaneous, and rapid detection of the activation states of all NHRs. This tool can be used in *in vitro* assays for receptor activation with agonists and antagonists, and can be used to generate cell lines and animal models that report on the activation state of such receptors in a biological setting and  
10 as a function of drug or drug lead.

*GPCR Nucleic acid sensor molecules*

Also provided by the invention are nucleic acid sensor molecules for detection of conformational isoforms of G-protein coupled receptors.

G-protein coupled receptors (GPCRs) play fundamental roles in regulating the  
15 activity of virtually every cell. Upon binding of extracellular ligands, GPCRs interact with a specific subset of heterotrimeric G-proteins that can then, in their activated forms, inhibit or activate various effector enzymes and/or ion channels. Molecular cloning studies have identified multiple human GPCRs, and have identified the ligands for many of these.

GPCRs include three domains: an extracellular N-terminus, a central domain of  
20 seven trans-membrane helices connected by unstructured loops, and a cytoplasmic C-terminus. Activation of GPCRs is induced by ligand binding, which causes a conformational change in the receptor transmitting a signal across the plasma membrane to intracellular members of a signaling pathway. The methods described herein allow for generating unique nucleic acid sensor molecules to any GPCR. In addition, methods are  
25 described that enable the generation of nucleic acid sensor molecules capable of recognizing the activated state of the GPCR by selecting for nucleic acid sensor molecule geometries which signal the presence or absence of the activated conformation of the receptor through recognition of one or all of the mobile domains, but whose signaling action is quiescent in the presence of other forms of the GPCR. The biosensors described herein include nucleic  
30 acid sensor molecules such as allosteric ribozymes (AR), including those derived from the hammerhead, hairpin, L1 ligase, or group I intron ribozymes and the like. Nucleic acid sensor molecules may also be derived from aptamer beacons or signaling aptamers which transduce molecular recognition into a detectable signal.

Upon the activation by an extracellular ligand or stimuli, G-protein coupled receptor(GPCR) polypeptides activate intracellular G $\alpha$ -protein. A single GPCR can activate a number of G $\alpha$ -proteins. For example, adrenergic receptors activate Gi, which inhibit adenylyl cyclases, Gs, which stimulate adenylyl cyclases, and Gq, which regulate cellular Calcium ion level (Wenzel-Seifert and Seifert 2000). Thus, it is highly desirable to distinguish which class of G $\alpha$ -proteins are activated through the GPCR of interest in cells.

The initial drug screening of the GPCRs is normally performed by competition assay with radiolabeled ligands. For a cell based GPCR assay, incorporation of radiolabeled GTP can be measured to detect the coupling of G $\alpha$ -protein and GPCR, however this assay does not distinguish the type of G $\alpha$ -proteins involved. The assays for the effect on individual effectors, such as the Ca<sup>++</sup> flow or cellular cAMP level, are also commonly used, but only one downstream signal can be measured at a time using these assays.

Upon activation, a G $\alpha$ -protein undergoes significant conformational change which results in release of GDP and association with GTP (Coleman and Sprang 1998). The G $\alpha$ -protein also dissociates from its  $\beta\gamma$ -subunits. This activated form of G $\alpha$ -protein then becomes capable of interacting with its effector (Li, Sternweis et al. 1998). The well-characterized conformation change takes place in three switches; switch I (residues 177-187 in G $\alpha$ 1), switch II (residues 199-219 in G $\alpha$ 1), and switch III (residues 231-242 in G $\alpha$ 1). The sequences and the conformational changes in these switches are well conserved among G $\alpha$ -proteins.

Ras is a member of the small GTPase protein family, which shares significant similarity with other family members. GTP-bound ras and GDP-bound ras can be distinguished by the RBD (ras binding domain) of Raf-1 (Taylor, Resnick et al. 2001). The activated state of Raf-1 can be identified by RalRDS (Franke, Akkerman et al. 1997). This indicates significant change in the surface of the protein, and the effector binding surface are only available for interaction in GTP complex form.

The invention provides methods for selecting nucleic acid sensor molecules which recognize the conformational change upon GTP binding and/or specifically interact with newly exposed G-protein effector binding sites upon the activation. Class-specific activated G $\alpha$ -protein nucleic acid sensor molecules recognize the activated G $\alpha$ -proteins or its effector binding site, which allow us to interpret the multiple type of downstream signal affect by the GPCR. It can used in both *in vitro* HTS (high throughput screening) and cell-based HTS:

Also described is a method for developing a direct mechanistic assay of the action of small molecule ligand-agonism, -antagonism, and partial antagonism of members of the GPCR family. The mechanistic assays function in both *in vitro* biochemical and *in vitro* cell-based settings. In the *in vitro* assay setting, the nucleic acid sensor molecules are  
5 designed to recognize one conformational isomer of the GPCR.

In one embodiment, the nucleic acid sensor molecule recognizes the unique conformation that exists for the activated state when in complex with ligand, *e.g.*, such as that observed for the beta-2 adrenergic receptor when in complex with the artificial ligand isoproterenol (Ghanouni et al., PNAS USA, 98:5997-6002(2001)) and then produces a  
10 detectable signal, *e.g.*, by release of a fluorescently labeled oligonucleotide, release of a radiolabeled oligonucleotide, or a change in conformation of the NASM driven by ligand binding through a change in fluorescence or the like. Hence, in this embodiment, the nucleic acid sensor molecule transduces molecular recognition of the beta-2 adrenergic receptor – in complex with epinephrine, norepinephrine or an artificial ligand such as isoproterenol into a  
15 detectable signal. The level of the signal is then used to quantify the amount of beta-2 adrenergic receptor-agonist complex present in solution.

In another embodiment, the beta-2 adrenergic-agonist nucleic acid sensor molecule is used as a screening tool in assays designed to detect agonists of the beta-2 adrenergic receptor. These screening tools can be used to determine the activation potency of  
20 compounds in *in vitro* biochemical assays or in *in vitro* cell-based assays. Agonists of the beta-2 receptor are useful in the treatment of asthma (Robinson, et al. Lancet 357:2007-2011(2001)). In another embodiment, nucleic acid sensor molecules are introduced into cell lines by known methods of electroporation, transfection, or coupling to peptide translocating agents such as tat or antennapedia peptides.

In another embodiment, the beta-2 adrenergic receptor-agonist complex specific  
25 nucleic acid sensor molecule is an allosteric intron imbedded in a reporter gene such as GFP or luciferase or the like. When the intron-derived reporter is inserted into the reporter gene it renders reporter gene expression effector dependent. Thus, in one embodiment functional GFP protein is expressed only when the beta-2 adrenergic receptor-agonist complex is  
30 present in the cell, and inhibitors of beta-2 adrenergic receptor-agonist complex formation block functional GFP protein expression in appropriate cells such as mammalian human peripheral blood leukocytes, yeast, insect, or bacterial cell lines.

In a further embodiment, Chinese hamster fibroblasts, which do not express beta-adrenergic receptors (Sheppard, et al., PNAS USA 80:233-236(1983)), are transfected with

both the nucleic acid sensor molecule and the gene coding for the beta-adrenergic receptor under a constitutive promoter, and are used to create a model cell line suitable for HTS screening of candidate beta-2 agonists. Furthermore, cells can be caused to express known allelic variants, such as gln27-to-glu associated with obesity (Large, et al., J Clin. Invest 5 100:3005-3013), to create cells lines which model specific disease states. Furthermore, chimeric mice can be created by "knock-in" (Monroe et al., Immunity 11:201-212(1999)) which will express the nucleic acid sensor molecule in every cell as the result of blastocyst fusion (Chen et al., PNAS USA 90:4528-4532(1993)), and used for pharmacokinetic or bioavailability studies in which the GPCR activation states of various tissues in the 10 organism are of concern.

GPCRs bind antagonists, which cause the GPCRs to become resistant to conformational changes, or result in conformations not susceptible to activation, or blockade the ligand binding domain from interaction with the appropriate ligand and thus prevent activation of the GPCR, such as the beta-2 adrenergic receptor antagonist 15 butoxamine (Horinouchi et al., Pharmacology 62:98-102(2001)). Hence, the invention also provides a method for using a nucleic acid sensor molecule to detect conformers which result from binding of GPCRs to antagonists. Furthermore, when the cell line described above is transfected with a mutant variants of GPCRs which spontaneously adopt the active conformation, such as lys272-to-ala (Pei, et al., PNAS USA 91:2699-2702(1994) and 20 references therein) the nucleic acid sensor molecule can be employed in a screen for compounds which are beta-2 antagonists (Ramsay et al, Br J Pharmacol 133:315-323(2001)). Antagonists of the beta-2 receptor are useful in the treatment of cardiovascular diseases (Nagatomo, et al., Cardiovasc Drug Rev 19:9-24(2001)). The invention accordingly provides a method for using a Beta-2 adrenergic receptor – butoxamine 25 complex-specific nucleic acid sensor molecule that is used to detect the levels of an antagonist specific complex in both *in vitro* biochemical, cell-based, and *in vivo* assays as described above.

Nucleic acid sensor molecules can also be developed that are specific for the occupancy state of the ligand-binding domains of all GPCRs. In addition, nucleic acid 30 sensor molecules for the agonist, antagonist, dimeric, or multimeric forms of all GPCRs can be used to screen for inhibitors or activators of GPCR function and therefore for inhibitors or activators of GPCR-dependent cell signaling pathways. Nucleic acid sensor molecules specific for individual GPCR complexes can be additionally used to screen for agents that modify GPCRs in *in vitro* and *in vivo* assays.



*Phosphodiesterase-specific nucleic acid sensor molecules*

Multiple classes of phosphodiesterases (PDEs) have been identified in humans. These enzymes catalyze a reaction that converts second messenger cAMP and cGMP into 5'-AMP and 5'-GMP. Different class of PDEs have different substrate specificity as well as different physiological function. For example, PDE4s are specific for cAMP and PDE5 are specific for cGMP.

The invention provides multiple classes of PDE nucleic acid sensor molecules. The first class of nucleic acid sensor molecules can distinguish cAMP vs. 5'AMP (cGMP vs 5'GMP) (Koizumi, Kerr et al. 1999) (Koizumi, Soukup et al. 1999). The second class of nucleic acid sensor molecule binds to the active site of PDE in a class specific manner and inhibits PDE catalytic activity. This class of nucleic acid sensor molecule can be raised using PDEs in the presence and absence of high affinity known inhibitors (e.g., Ropalin for PDE4). The third class of nucleic acid sensor molecule recognizes PDE in a class-specific (e.g., PDE1-11) or subclass-specific (PDE4A-D) manner.

*Protein kinase-specific nucleic acid sensor molecules*

The invention also provides nucleic acid sensor molecules raised against protein kinases. In one embodiment, the invention provides nucleic acid sensor molecules that are modulated by the phosphorylation state in a given peptide sequence. Alternatively, native proteins can be used with different phosphorylation states in order to raise nucleic acid sensor molecules that can distinguish the different phosphorylation states. For example, ERK1/2 and phosphorylated ERK1/2 can be distinguished by specific nucleic acid sensor molecules (Seiwert, Stines Nahreini et al. 2000). The nucleic acid sensor molecule also can be a competitive inhibitor for a kinase by binding at ATP or substrate binding sites.

Alternatively, an ADP-dependent nucleic acid sensor molecule can be obtained at lower pH. These nucleic acid sensor molecules can be used to detect the production of ADP.

**ii. Pathway Profiling Biosensors**

As shown in Figure 7, physiological function is modulated by complex pathways, each of which may have multiple overlapping and intersecting steps. Furthermore, the proteins involved in these pathways are highly homologous and can have overlapping substrates and drug specificities. Using current techniques, it is possible only to monitor the

response of single elements of a pathway. These techniques are inadequate to understand the mechanism of drug interactions. For example, a particular drug found to have a particular *in vitro* activity against a single target in biochemical assays might interact with other proteins in the same pathway, or in other unrelated pathways. Consequently,  
5 physiological function is often uncorrelated with the results of biochemical assays of a single target.

The nucleic acid sensor molecules according to the invention provide reagents to simultaneously quantify the level and chemical state of all components in a molecular pathway. As used herein, "pathway target molecules" are target molecules involved in the  
10 same pathway and whose accumulation and/or activity is dependent on other pathway target molecules, or whose accumulation and/or activity affects the accumulation and/or activity of other pathway target molecules. Pathway target molecules according to the invention include, *e.g.*, proteins, such as enzymes, modified forms of proteins, such as phosphorylated, sulfated, ribosylated proteins, methylated proteins (Arg, Asp; N, S or O  
15 directed), prenylated proteins (such as by farnesyl, geranylgeranyl, and other types of prenylation) acetylated or acylated proteins, cleaved or clipped proteins, bound or unbound forms of proteins, allelic variants of a protein (*e.g.*, proteins differing from each other by single amino acid changes in a protein), as well as substrates, intermediates, and products of enzymes (including both protein and non-protein molecules).

20 In another embodiment, diagnostic pathway target molecules are identified by pre-selecting a plurality of nucleic acid sensor molecules activatable by pathway-specific target molecules. In one embodiment, a profiling biosensor is provided comprising at least one nucleic acid sensor molecule specific for every molecular species within a pathway (*e.g.*, a signaling pathway), to generate a biosensor which can monitor the levels, chemical  
25 structure, and/or activity of every molecular species in the pathway.

Because of the finite number of target molecules (as determined from data obtained from the Human Genome Project) and the high throughput of the biosensors of the instant invention (greater than 10,000 target molecules can be monitored simultaneously), the profiling biosensors of the instant invention make it feasible to evaluate the response of all  
30 the components of a pathway to a drug compound simultaneously.

In one embodiment, a profiling biosensor reactive to the components of an entire pathway, is contacted with a sample from a patient having a disease, and an optical signal corresponding to a disease state is determined to identify diagnostic pathway target molecules which are diagnostic of that disease. Samples from a plurality of patients are

obtained and tested using the profiling biosensor to identify a pathway profile that is diagnostic of the disease, the pathway profile comprising normalized data relating to any or all of the level, structure, and activity, of the signature pathway molecules. A pathway profile corresponding to a wild type state is determined by testing the profiling biosensor molecules against samples from a population of healthy patients, or subsets of populations of healthy patients. In one embodiment, data relating to the optical signals generated by nucleic acid sensor molecules activated by the diagnostic pathway target molecules is stored within the memory of a computer or within a computer program product.

The pathway profiles can be used in diagnostic testing as discussed above. In one embodiment, a profiling biosensor is used in which the pathway is one which is known or suspected of being disrupted in patients having a particular trait (e.g., having a particular disease or genetic alteration(s)). For example, in one embodiment, one profiling biosensor used to evaluate samples from a patient with cardiovascular disease is a cholesterol metabolism pathway profiling biosensor. However, random combinations of profiling biosensors can be used to assess the physiological state of a patient, to identify diagnostic pathway profiles which are diagnostic of diseases whose molecular basis has not yet been identified or characterized.

In one embodiment, profiling biosensors according to the invention are used to assess the affect of a candidate drug on any or all of the level, chemical structure, or activity of diagnostic pathway target molecules to generate a drug treatment pathway profile. In this embodiment, a profiling biosensor is contacted with a sample from a cell or physiological system (e.g., a group of cells, a tissue system, an organ system, or a patient), and changes in optical signals are obtained which are correlated to any, or all of, the level, chemical structure, or activity of a particular pathway target molecule by relating the optical signal obtained to the address of the nucleic acid sensor molecule, as described above. In one embodiment, a drug treatment profile which is substantially similar to a diagnostic pathway profile obtained from a healthy population of patients (as determined by observing no significant differences in the profile by routine statistical testing, to within 95% confidence levels) is used to identify a candidate drug as one which is suitable for further testing. The profile produced by such a drug is used to produce an effective drug treatment profile, against which other candidate drugs can be compared.

In another embodiment, a candidate drug is tested against a plurality of profiling biosensors including the one which will generate a diagnostic signature profile, to identify drugs which produce an effective drug treatment profile without effecting significant

changes in other pathway profiles. In this embodiment, the systemic effects of a candidate drug can be predicted.

In further embodiments, it is desirable to use a biosensor representing less than an entire pathway. In one embodiment, a biosensor is provided comprising nucleic acid sensor molecules diagnostic pathway target molecules. In a further embodiment, a biosensor is provided which comprises nucleic acid sensor molecules necessary to evaluate particular components of a pathway suspected of being involved in a disease. For example, compounds being screened to identify candidate drugs that affect diseases relating to defective DNA repair can be tested against a pathway biosensor comprising only S phase cell cycle target molecule reactive nucleic acid sensor molecule.

Exemplary pathway target molecules include various proteins and modified form thereof. Modifications include post-translational modifications such as phosphorylation, prenylation, glycosylation, methionine removal, N-acetylation, acylation, acylation of cysteines, myristoylation, alkylation, ubiquitinylation, prolyl-4-hydroxylation, carboxylation of glutamyl residues, advanced glycosylation (e.g., of hemoglobin), deamination of glutamine and asparagine, addition of glycophosphatidylinositol, disulfide bond formation, hydroxylation, and lipidation. Examples of such proteins include ERK and phosphorylated ERK, CDK and phosphorylated CDK, modified cyclin A, cyclin D, cyclin E, k-Ras, h-Ras, Rho A, MEK-1, MEKK-1, Raf-1, Raf-A, JNK, PKA, ATK, PTEN, p53, P16, and INK4. Other proteins include those listed in Table 1 as well as post-translationally modified forms thereof.

Table 1

apoptotic pathways		Bcl, Bak, ICE proteases, Ich-1, CrmA, CPP32, APO-1/Fas, DR3, FADD containing proteins, perform, p53 tumor necrosis factor (TNF) receptor, NAIP, TAP, TRADD-TRAF2 and TRADD-FADD, TNF, D4-GDI, NF-kB, CPP32/apopain, CD40, IRF-i, p53, apoptin
blood clotting pathways		thrombin, fibrinogen, factor V, Factor VIII- FVa, FVIIIa, Factor XI, Factor Xia, Factors IX and X, thrombin receptor, thrombomodulin (TM), protein C (PC) to activated protein C (aPC). aPC, plasminogen activator inhibitor-i (PAT-i), tPA (tissue plasminogen activator)
calcium signaling pathways		calmodulin, calcineurin,
Cell cycle	G0	MPS, CYTOSTATIC FACTOR (CSF) (INCLUDING MOS)
Pathway	G1	mid G1 phase: cdk4/cycin D1-3 and cdk6/cyclin D1-3 late G 1 phase: cdk2/cycin E others: p53, p21, p16, Rb, p2'1, E2F, Cdc25A, Cdc25B

	S	cyclin A/CDK2, cyclin B/Cdc2, SPA-i, Cdc25A, Cdc25B
	G2	G2/M transition phase: cdkl/cyclin B 1-3, cdkl/cyclin A, Cdc25A, Cdc25B, Cdc25C. PIN1, Chki, Myt 1, Wee 1
	M	Cdc2/cyclin B, Plk, Cdc25C,
Cholesterol metabolism pathway		LDL, LDL-receptor, VLDL, HDL, cholesterol acyltransferase, apoprotein E, Cholesteryl esters, ApoA-I and A-II, HMGCoA reductase, cholesterol
Flt-3 pathway		flt-3 pathway flt-3, GRP-2, SHP-2, SHIP, She
JAK/STATS signaling pathway		Jak1, Jak2, IL-2, IL-4 and IL-7, Jak3, Ptk-2, Tyk2, EPO, GH, prolactin, IL-3, GM-CSF, G-CSF, IFN gamma, LW, OSM, IL-12 and IL-6, IFNR-alpha, IFNR- gamma , IL-2R beta , IL-6R, CNTFR, Stat 1 alpha, Stat 1 beta , and Stats2-6
MAP kinase signaling pathways		flt-3, ras, raf, Grb2, Erk-i, Erk-2, and Src, Erb2, gpl30, MEK-1, MEK-2, hsp 90, JNK, p38, Sin1, Sty1/Spcl, MKK's, MAPKAP kinase-2, JNK,SAPK
P53 pathway		bax, bid, caspases, cytochrome c
PI 3 kinase pathway		SHIP, Akt
ras activation pathways		p120-Ras GAP, neurofibromin, Gap1, Ra1-GDS, Rsbs 1, 2, and 4, Rin1, MEKK- 1, and phosphatidylinositol-3-OH kinase (P13K), ras
SIP signaling pathways		GRB2, SIP, ras, P1 3-kinase
SHC signaling pathways		trkA, trkb, NGF, BDNF, NT-4/5, trkC, fnt-3, Shc, PLC gamma 1, P1-3 kinase, SNT, ras, rafi, MEK and MAP kinase
TGF-13 signaling pathways		BMP, Smad 2, Smad4, activin, TGF
T-cell receptor complex		lck, fyn, CD4, CD8, T cell receptor proteins
MHC-I pathways		TAP proteins, LMP 2, LMP 7, gp 96, HSP 90, HSP 70

If desired, nucleic acid sensor molecules can be raised against particular amino acid sequences in the polypeptides. Some representative peptide regions are presented in Table 2.

5

Table 2

Sequence	Enzyme
L-R-A-S-L-G (SEQ ID NO:52)	PKA
A-A-K-I-Q-A-S-F-R-G-H-M-A-R-K-K (SEQ ID NO:53)	PKC
P-K-T-P-K-K-A-K-K-L(SEQ ID NO:54)	cdc2
E-P-P-L-S-Q-E-A-F-A-D-L-W-K-K(SEQ ID NO:55)	DNA-PK
D-D-D-B-B-S-I-T-R-R(SEQ ID NO:56)	CK-1
R-R-R-B-B-E-T-E-E-E (SEQ ID NO:57)	CK-2
K-K-A-L-R-R-Q-E-T-V-D-A-L (SEQ ID NO:58)	CaM KII
S-T-K-V-P-Q-T-P-L-H-T-S-R-V (SEQ ID NO:59)	P38
R-R-R-R-S-I-I-F-I (SEQ ID NO:60)	PKA
R-R-R-R-R-K-G-S-F-R-R-K-A (SEQ ID NO:61)	PKC $\alpha$
R-K-L-K-R-K-G-S-F-R-R-K-A (SEQ ID NO:62)	PKC $\beta$ I, II
R-R-R-R-R-K-G-S-F-K-K-F-A (SEQ ID NO:63)	PKC $\gamma$
A-A-R-K-R-K-G-S-F-F-Y-G-G (SEQ ID NO:64)	PKC $\delta$

Y-Y-X-K-R-K-M-S-F-F-E-F-D (SEQ ID NO: 65)	PKC $\epsilon$
A-R-L-R-R-R-R-S-F-R-R-X-R (SEQ ID NO: 66)	PKC $\eta$
R-R-F-K-R-Q-G-S-F-F-Y-F-F (SEQ ID NO: 67)	PKC $\zeta$
A-A-L-V-R-Q-M-S-V-A-F-F-F (SEQ ID NO: 68)	PKC $\mu$
K-R-Q-Q-S-F-D-L-F (SEQ ID NO: 69)	CaM KII
F-R-M-M-S-F-F-L-F (SEQ ID NO: 70)	Phosphorylase kinase
R-R-F-G-S-L-R-R-F (SEQ ID NO: 71)	SLK1
R-R-R-H-S-R-R-R-R (SEQ ID NO: 72)	SRPK2
R-K-R-X-R-T-Y-S-F-G (SEQ ID NO: 73)	AKT/PKB

In one embodiment, a profiling biosensor array is generated comprising target activatable nucleic acid sensor molecules which are activatable by components of a cell cycle pathway. In this embodiment, a cell cycle biosensor is generated comprising nucleic acid nucleic acid sensor molecules activatable by at least two members selected from the group consisting of: MPS, Cytostatic factor (CSF) (including Mos), cdk4, cyclins D1-3, cdk6, cdk2, cyclin E, p53, p21, p16, Rb, p27, E2F, cyclin A, cyclin B, cdk1, cyclin B1-3, Cdc2, SPA-1, and other biomolecules involved in cell cycle regulation.

In another embodiment, the cell cycle biosensor array generated is used to evaluate samples from patients suspected of having a disorder affecting cell proliferation (*e.g.*, cancer) and a signature target molecule profile is determined which is diagnostic of this disorder. Changes in the signature target molecule profile upon treatment with a candidate compound are subsequently monitored by any or all of *in vitro*, *ex vivo*, and *in vivo* methods, as described above, to identify and/or validate lead compounds for use in cancer therapies.

In further embodiments, a cell cycle biosensor is provided comprising a plurality of locations, each location comprising a set of nucleic acid sensor molecules activatable by target molecules which identify a different portion of the cell cycle. Thus, in one embodiment, a cell cycle biosensor comprises at a first location, nucleic acid sensor molecules activatable by G0 specific target molecules (*e.g.*, MPS, Cytostatic factor (CSF) (including Mos)), at a second location, nucleic acid sensor molecules activatable by G1 specific target molecules (cdk4, cyclin D1-3, cdk6, cdk2, cyclin E, p53, p21, p16, Rb, p27, E2F), at a third location, nucleic acid sensor molecules which are activatable by S specific target molecules (*e.g.*, cyclin A/CDK2, cyclin B/Cdc2, SPA-), at a fourth location, nucleic acid sensor molecules activatable by G2 specific target molecules (*e.g.*, cdk1, cyclin B 1-3, cyclin A), and at a fifth location, nucleic acid sensor molecules activatable by M specific

target molecules (*e.g.*, Cdc2, cyclin B). In this way the effects of diseases and/or drugs on specific phases of the cell cycle can be assessed.

Similarly, pathway specific biosensors can be generated for any of apoptotic pathways, blood clotting pathways, calcium regulation pathways, cholesterol metabolism  
5 pathways, the JAK/STATS signaling pathway, MAP kinase signaling pathways, p53 pathway, PI 3 kinase pathway, ras activation pathways, SIP signaling pathways, SHC signaling pathways, TGF- $\beta$  signaling pathways, T-cell receptor complex, and MHC-I pathways, using exemplary target molecules listed above, or other target molecule components of the respective pathways.

10 It should be apparent to those of ordinary skill in the art, that many other pathways exist whose components have been characterized and that target molecules within these pathways are also encompassed within the scope of the present invention (*e.g.*, including, but not limited to, phosphatase pathways, transcription factor pathways, hormone dependent pathways, as  
15 well as intermediary metabolism pathways, and developmental pathways). Further, additional pathways can be identified using the nucleic acid based biosensor profiling techniques discussed above (*e.g.*, identifying pathway molecules involved in the functioning of a wild type or diseased organ system, such as the cardiovascular system, central nervous system, digestive system, reproductive system, pulmonary system, skin system, and the  
20 like), and these also are encompassed within the scope of the invention.

Alternatively, or additionally, pathway specific molecules can be identified by other techniques known in the art (*see, e.g.*, U.S. Patent Number 6,087,477, U.S. Patent Number 6,054,558, U.S. Patent Number 6,048,709, and U.S. Patent Number 6,046,165) and used to engineer additional pathway target activatable nucleic acid sensor molecules. Because there  
25 is a finite number of pathway target molecules in each pathway (constrained by the absolute number of gene products which have been identified) (*see, e.g.*, Drews, Science 287: 1960-1964), it is feasible using the target activatable nucleic acid sensor molecules to generate biosensors representative of an entire pathway.

In further embodiments, sets of profiling biosensors are used to monitor the  
30 expression/activity of target molecules representing complex systems. Thus, for example, the effect of target molecules on the cardiovascular system and pulmonary system can be monitored simultaneously. In one embodiment, an array representative of a plurality of systems in the human body is used in methods to assess the effects of drug compounds on multiple systems in the body.

iii. Using Pathway Nucleic Acid Sensor Molecules in Drug Optimization

5 The profiling nucleic acid sensor molecules according to the invention can be used in every step of a drug optimization process, as shown in Figure 8, and are suitable reagents for use in conventional high throughput screening systems making them extremely adaptable for use alone, or in conjunction with, other drug development assays.

Step 1. Drug Target Discovery or Drug Target Validation

10 As discussed above, profiling nucleic acid sensor molecules can be used to identify signature target molecules which are correlatable to particular traits, such as disease. Signature target molecules are drug targets whose levels, structure, and/or activity can be used to evaluate the efficacy of compounds. A large number of signature drug targets, both characterized and uncharacterized, can be identified simultaneously using a single profiling biosensor according to one embodiment. In one embodiment, a profiling biosensor  
15 recognizes and is independently activated by about 1-5,000 molecules. In another embodiment, a profiling biosensor recognizes and is independently activated by about 500-10,000 molecules, and in another embodiment, by greater than 10,000 molecules.

Step 2. High Throughput Screening

20 In one embodiment, the drug targets identified in step 1 are evaluated in high throughput screening assays, using either solution-based biosensors or substrate-based biosensors, to characterize the biological activity of a drug target. For example, in one embodiment, nucleic acid sensor molecules are used to assess levels of substrate, product and intermediates produced by an enzyme in a wild type vs. a disease state, to identify other components of a pathway that would be affected by a drug acting on that target (*i.e.*,  
25 secondary drug targets). In another embodiment, the levels, structure, and/or activity of all of the modified forms of a drug target, or the active and inactive forms of a drug target (*e.g.*, a receptor) is determined in a wild type vs. a disease state, to further develop a diagnostic profile of a diagnostic pathway target molecule and to evaluate changes of that profile in the presence of a drug.

30 In a further embodiment, the same type of profiling biosensor used to identify a diagnostic profile is contacted with samples from patients exposed to a compound. A compound-treated sample which produces substantially similar levels, structure, and/or



activity of target and secondary drug targets in a sample from a healthy patient is used to identify a compound as a candidate drug. Because this testing is done in a high throughput format, a single dose of a candidate drug is evaluated in any given test.

5           **Step 3.           In vitro Biochemical Assays**

In one embodiment, the nucleic acid sensor molecules used in step 2, are tested in an *in vitro* biochemical assay to determine compound potency. In this embodiment, a preliminary dosing effect is determined to identify the IC<sub>50</sub> of candidate drug. In one embodiment, multiple biosensors of the type used in step 2 are contacted with samples from  
10 patients exposed to different doses of the candidate drugs identified in step 2, to identify candidate drugs with the highest potency (e.g., requiring the least amount of drug to generate a wild type profile or an effective drug profile.

**Step 4.           Cellular Assays**

15           In one embodiment, nucleic acid sensor molecules are used in cellular assays where the effect of adding a compound on cell physiology is known and the researcher wants to determine that the drug is in fact acting through the drug target selected in steps 1-3. Here a candidate drug is added to a physiological system (e.g., cell(s), tissue(s), organ(s), or a patient). Cells from the physiological system are lysed and the substrate or product of an  
20 enzyme reaction is monitored using the nucleic acid sensor molecule either in an ELISA format or other solid support-based format (e.g., a profiling array) or a solution phase format. In another embodiment, cell lysates are contacted with a profiling biosensor specific for a target or pathway of interest to determine the profile of target molecules in the lysed sample. The profile is then compared to the wild type profile and the disease profile to  
25 determine if the drug is operating *in vivo* to restore a cell to its wild type state. Thus, the physiological effect of a candidate drug on a physiological system is correlated with the *in vivo* mechanism of action of the candidate drug.

In a preferred embodiment, pathway profiling arrays comprised of nucleic acid sensor molecules affixed to a solid support are used in cellular assays to determine the  
30 selectivity of a compound for one target in a pathway relative to other candidate targets in a signal transduction pathway(s) or in another biochemical pathway(s). This data can be used to validate a drug lead or drug target.

In one embodiment, nucleic acid sensor molecules are expressed *in vivo* or intracellularly using plasmids, viruses or other extra-chromosomal DNA vectors and the

cellular nucleic acid sensor molecules are extracted and used to determine the activity of a drug or drug target. These cellular assays can also determine the selectivity of a compound for one target in a pathway relative to other candidate targets in a signal transduction pathway(s) or in another biochemical pathway(s). This data can be used to validate a drug lead or drug target.

*In vivo* detection:

With (Amersham) SPA scintillant beads coupled to nucleic acid sensor molecules, one can look at cellular processes in situ in real time, by culturing cells directly onto a microtiter plate and allowing uptake of scintillant beads and radioisotope by cells. One can then monitor biosynthesis, proliferation, drug uptake, cell motility, etc. via luminescence generated by beads in the presence of selected target.

**Step 5. Medicinal Chemistry**

In one embodiment, drug-lead potency, specificity, and/or *in vivo* activity is optimized by an iterative repetition of any or all of steps 1-4. In one embodiment, steps 1-4 are repeated until the desired potency, selectivity and *in vivo* mechanism of action of a candidate drug is obtained. Potency can range from picomolar affinity to nanomolar affinity as measured by *in vitro* IC50 values. The desired selectivity of a drug candidate for its target can vary from 2 to a million-fold, and can be obtained by measuring the potency (IC50) of a drug lead toward the drug target, versus the drug's potency (IC50) values against other pertinent targets (target pertinence is determined by the requirements of the biological system under investigation). A drug lead is deemed optimal when the parameters of potency, selectivity and cellular action are optimized with respect to each other.

In another embodiment, known drug leads from Steps 1-4 are found to be specific for targets that were not known to the researcher in step 2. This is also termed target discovery and validation, and occurs when steps 1-4 are repeated in an iterative fashion of any or all steps and the drug target is identified by the profiling array to, in fact, exist in an alternative signal transduction pathway, or to be a novel protein or enzyme in the pathway originally under investigation. Thus, MPP arrays can identify the site of action of a drug lead, and can determine the relative selectivity of a drug for one drug target of a drug target pathway.

**Step 6. Animal Model Assays**

In this embodiment, target cells (*e.g.*, tissue(s)) are removed from an animal model of the disease being targeted for treatment and lysed for testing. The lysate is contacted with nucleic acid sensor molecules either in a solid phase assay, a solution phase assay, or in a pathway profiling biosensor array format to assess the *in vivo* biological activity of a candidate drug identified by any of the previous steps or by some other method, on a target or pathway. Thus, in this embodiment, the physiological effect of a drug on a diseased or normal tissue is correlated with the *in vivo* mechanism of action of the drug.

**Step 7.      Optimization of the Drug Lead**

In one embodiment, drug-lead potency, specificity, and/or *in vivo* activity are optimized by an iterative repetition of any or all of steps 1-6. In one embodiment, steps 1-5 are repeated until the desired potency, selectivity and *in vivo* mechanism of action of a candidate drug are obtained.

**Step 8.      Pharmacokinetic Studies**

In one embodiment, the nucleic acid sensor molecules are used in pharmacokinetic studies, where the effect of a drug on the physiology of a cell, group of cells, tissue(s), organ(s), or animal model is assessed by obtaining blood, plasma, tissue, or a cell, and contacting this material with nucleic acid sensor molecules either in a solid phase (*e.g.*, ELISA), solution or array format to assess the *in vivo* pharmacological or toxicological activity of a compound. In this embodiment, the nucleic acid sensor molecules used are developed against the candidate drug itself, its metabolic products, and/or the metabolic products of proteins and small ligands involved in a xenobiotic or toxicological response to drug treatment.

In one embodiment, nucleic acid sensor molecules are employed to follow the fate of a drug or its metabolic by-products. In this embodiment, nucleic acid sensor molecules are generated to the drug and its metabolites. The drug is administered to the test animal either subcutaneously, intraperitoneally or by gavage. Subsequent to administration, the blood plasma or disease tissue is removed and its contents are screened for the remaining drug by Liquid chromatography (LC) or LC-mass spectrometry. Drug exposure is then determined as a function of time, dose and method of administration and is reported in values of half-life, bioavailability, AUC and C<sub>max</sub>. Metabolic products of a drug lead can be similarly followed.

Nucleic acid sensor molecules generated against enzymes or proteins known to those skilled in the art to be involved in drug metabolism (P450 enzymes, multi-drug transporter) can be used to follow the effect of a drug on xenobiotic or toxicological response to drug treatment.

5

**Step 9.      Optimization of the Drug Lead**

In one embodiment, drug-lead potency, specificity, and/or *in vivo* activity, and pharmacokinetic, or toxicological properties are optimized by an iterative repetition of any or all of steps 1-7. In one embodiment, steps 1-7 are repeated until the  
10    desired potency, selectivity and *in vivo* activity and pharmaco-kinetic, or toxicological properties of a candidate drug are obtained.

**Step 10.     Clinical Trials**

In one embodiment, nucleic acid sensor molecules are used in clinical trials to  
15    determine the fate of a drug in human or animal models, or used to follow the effect of drug treatment on a target or molecular pathway of choice, as described above. In one embodiment, the nucleic acid sensor molecules, in a solid phase assay (*e.g.*, ELISA format), a solution phase assay, or in a pathway profiling biosensor array format, are used to assess the *in vivo* biological activity of a drug being tested using lysed cell samples as described  
20    above.

In another embodiment, the appropriate profiling biosensor is used *in vivo*, to monitor the effects of the compound on the patient, for example, by providing the biosensor in communication with a fiber optic probe inserted into the patient, or *ex vivo*, monitoring optical signals in a cell using a microscope based detection system. In another embodiment,  
25    an *in vivo* assay is done by introducing a nucleic acid sensor molecule which retains its catalytic activity into a physiological system (*e.g.*, by injection at a target site in the body, through liposome carriers, and other means of administration routinely used in the art), obtaining cells from the physiological system and detecting the effect of the compound on the catalytic activity of the nucleic acid sensor molecule (*e.g.*, by evaluating the sequence of  
30    the nucleic acid sensor molecule) as a means of determining the level, structure, or activity of a drug target, and relating the level, structure, or activity of the target molecules to the efficacy of the drug.

Step 11. Optimization of the Drug Lead

In one embodiment, any or all of steps 1-10 are repeated to further optimize the properties of the candidate drug.

5 Step 12. Diagnostic Applications

In one embodiment, individuals who would be suitable for treatment with the candidate drugs identified steps 1-11, are identified using nucleic acid sensor molecules in the diagnostic assays discussed previously.

10 Step 13. Chemical Genomics

In one embodiment, nucleic acid sensor molecules are used in chemical genomic assays in which a drug or plurality of drug leads, with known or unknown physiological effects, and with unknown targets, are contacted with a physiological system and the site of action of the drug or plurality of drugs is determined using a plurality of the profiling biosensors described previously. Drug optimization then occurs as in steps 1-11.

15 5. Use of a Profiling Biosensor in Target Molecule Separation

In addition to, or instead of, their use in detection methods, and drug discovery methods, the nucleic acid sensor molecules according to the invention can also be used to retrieve the target molecules which they specifically recognize. Additional embodiments exploiting the recognition capacity of the biosensors disclosed are contemplated and encompassed within the scope.

20 6. Reagents for Generating and Using Nucleic Acid Sensor Molecules

In one embodiment, reagents are provided for generating and using nucleic acid sensor molecules. In one embodiment, a kit is provided comprising standardized reagents for making and/or using the nucleic acid sensor molecules according to the invention. In one embodiment, the kit comprises at least a first optical nucleic acid sensor molecule whose optical properties change upon recognition of a target molecule. In another embodiment, the kit additionally comprises any of: a control target molecule, a second nucleic acid sensor molecule which recognizes a different target molecule, suitable buffers, printed instructions, and combinations thereof. In a further embodiment, a nucleic acid sensor

molecule is provided with reagents for attaching a label and/or quencher or with reagents for attaching charge transfer molecules to the nucleic acid sensor molecule, which can sensitize the optical properties of the nucleic acid molecule to the presence of a target molecule.

5 In another embodiment, a composition is provided comprising a target molecule and a nucleic acid sensor molecule. The composition provides a reference against which to compare modified nucleic acid sensor molecules which recognizes to the same target, in order to select those with preferred catalytic activity or conformational change in the presence of the target. In a further embodiment, sets of complexes are provided. In still a further embodiment, a set of pathway target molecules and nucleic acid sensor molecules are provided. In another embodiment, a set of profiling target molecules and nucleic acid sensor molecules are provided. In still a further embodiment, solid supports are provided for isolation of target molecules from nucleic acid sensor molecules.

15 In yet another embodiment, a computer program product is provided comprising stored data relating to optical signals generated by profiling and or pathway target molecules. In another embodiment, a means to compare this data to other optical signals is provided. In a further embodiment, the memory comprises data relating to patient information or chemical structure information relating to either target molecules or nucleic acid sensor molecules.

20 The nucleic acid sensor molecules and biosensors according to the invention are amenable for use with high throughput screening systems and methods and the use of the nucleic acid sensor molecules and biosensors in these systems and methods is encompassed within the scope. In one embodiment, the system is a robotic workstation, comprising, at least one of an: arrayer, microplate or microarray feeders, stackers, washers, and dispensers, 25 an optical system, a carousel, a conveyer for conveying microplates or microarrays from one part of the system to another (in a vertical or horizontal direction), a shaker system or other mixing system, a temperature control system, a synthesizer, a solid phase extraction system, and sample concentrators. Components of the robotic workstation can be part of a single integrated system or can be provided separately for use at any stage of the drug 30 optimization process according to the invention. In a further embodiment, the system comprises a processor connectable to the network which comprises or can access applications comprising stored data relating to profiling information obtained using nucleic acid sensor molecules according to the invention, and/or statistical applications, applications for performing structure/activity analysis of target molecules and nucleic acid sensor

molecules, applications for performing nucleic acid sequence alignment and simultaneous structure superposition of proteins (*e.g.*, MOE-Align™), applications for predicting binding conformations of molecules to receptor structures, and applications for controlling the processing functions of the robotic workstations.

5           The invention is further illustrated in the following non-limiting examples.

## EXAMPLES

### Example 1: General Procedures

A. Generating nucleic acid sensor molecules from pools of ribozymes  
10   comprised of randomized linker and target modulation domains.

Direct selection of nucleic acid sensor molecules. Target modulated nucleic acid sensor molecules are isolated by *in vitro* selection. Pools of partially randomized ribozymes with  $10^{15}$ - $10^{17}$  unique sequences serve as the starting point for *in vitro* selection. As with the engineering approach, both the L1 ligase and hammerhead ribozyme are used as  
15   platforms for the selection of allosterically-controlled molecules. Selections are designed to yield ribozymes that specifically respond to any target. Nucleic acid sensor molecules with cross-specificity (*i.e.* modulated by alternate target molecules, or by alternate ligand-bound states, or by alternate post-translationally modified forms of protein or peptides) are selected against by including the undesired form of the target in an initial negative selection step.  
20   The specific sequence of operations comprising the selection experiment is outlined in Figure 1A and 1B for ligase and hammerhead-based selections, respectively

1.    **Pool preparation.** The starting library of DNA sequences is generated by automated chemical synthesis on a DNA synthesizer. This library of sequences is transcribed *in vitro* into RNA using T7 RNA polymerase, purified, and  
25    captured onto beads using an oligonucleotide tag complementary to the 3'-end.

2.    **Negative selection incubation.** In the absence of the desired modulator (target), the RNA library are incubated together with the undesired form of the modulator and an arbitrary sequence oligonucleotide substrate (substrate 1). (During this incubation, non-modulated ribozymes undergo ligation (Figure 1A) or  
30    cleavage (Figure 1B).

3.    **Positive selection incubation.** The undesired form of the modulator and oligonucleotide substrate 1 are removed by washing. The immobilized RNA

pool is then incubated under identical conditions but now in the presence of the intended modulator and a second biotinylated oligonucleotide substrate (substrate 2), in the case of ligase selections (Figure 1A).

4. **Partitioning on the basis of activity (coupled to amplification).**

5 Ligases active only in the presence of the desired modulator are isolated using streptavidin capture and selective PCR amplification (relying on sequence differences between substrates 1 and 2 to distinguish allosteric and non-allosteric activities). Hammerheads active only in presence of the desired modulator are isolated by gel electrophoresis.

10 5. **Purification.** PCR amplified DNA are purified and transcribed to yield an enriched pool for subsequent reselection.

6. **Iteratively repeat.** Rounds of selection and amplification (steps 2-5) are repeated until functional members sufficiently dominate the resultant library.

15 Once NASMs have been selected as described above, they are characterized as follows:

Nucleic acid sensor molecules which are derived from *in vitro* selection are tested as target modulated biosensors. The pool of NASMs is cloned into various plasmids that contain a T7 promoter transformed into *E. coli*. Individual NASM encoded DNA clones are isolated, linearized and the NASM is transcribed *in vitro* to generate NASM RNA. The  
20 NASM RNAs are then tested in target modulation assays which determine the rate or extent of ribozyme modulation. For hammerhead NASMs, the extent of target dependent and independent reaction is determined by quantifying the extent of endonucleolytic cleavage of an oligonucleotide substrate. The extent of reaction can be followed by electrophoresing the reaction products on a denaturing PAGE gel, and subsequently analyzed by standard  
25 radiometric methods. For Ligase NASMs, the extent of target dependent and independent reaction is determined by quantifying the extent of ligation of an oligonucleotide substrate, resulting in an increase in NASM molecular weight, as determined in denaturing PAGE gel electrophoresis.

Individual NASM clones which display high target dependent switch factor values,  
30 or high  $k_{act}$  rate values are subsequently chosen for further modification and evaluation.

Hammerhead derived NASM clones are then further modified to render (the NASM sequences) them suitable for the optical detection applications that are described in detail below. In brief, these NASMs are used as fluorescent biosensors affixed to solid supports,



as fluorescent biosensors in homogeneous FRET-based assays, and as biosensors in SPR applications.

Ligase derived NASM clones are further modified to render them suitable for a number of detection platforms and applications; including, but not limited to, the PCR and nucleotide amplification detection methods; fluorescent-based biosensors detectable in

solution and chip formats; and as *in vivo*, intracellular detection biosensors. The various detection applications of hammerhead ligase and intron-based NASMs are described in detail below.

10 **B. Preparation of an array of immobilized effector oligonucleotides or NASMs**

The following protocol describes a method for preparing an array of immobilized effector oligonucleotides with terminal amine groups attached to a solid substrate derivatized with aldehyde groups. The resulting array can then be used to spatially address (*i.e.*, the sequence of nucleotides for each effector oligonucleotide can be synthesized as a cognate to the effector oligonucleotide binding domain of a nucleic acid sensor molecule specific for a particular target molecule) and immobilize the nucleic acid sensor molecules prior to their use in a solid-phase assay (see, *e.g.*, Zammatteo et al., 2000):

Protocol for attachment of effector oligonucleotides to aldehyde derivatized substrate

20 ([www.arrayit.com](http://www.arrayit.com)):

1. Print discrete spots of solution containing effector oligonucleotides with amine-reactive terminal groups or linkers with terminal amine groups using microarraying pins, pipette, etc.
2. Allow spotted substrate to dry for 12 hrs. at room temperature and < 30% relative humidity.
3. Rinse substrate 2 times in dH<sub>2</sub>O with 0.2% SDS for 2 min. with vigorous agitation at room temperature.
4. Rinse substrate 1 time in dH<sub>2</sub>O for 2 min. with vigorous agitation at room temperature.
- 30 5. Transfer substrate to boiling (100° C) dH<sub>2</sub>O for 3 min. to denature DNA.
6. Dry substrate by centrifugation at 500x g for 1 min.
7. Treat substrate in 0.1 M NaBH<sub>4</sub> in phosphate buffered saline (PBS, pH 7) for 5 min. with mild agitation at room temperature.

8. Rinse substrate 2 times in dH<sub>2</sub>O with 0.2% SDS for 1 min. with vigorous agitation at room temperature.
9. Rinse substrate 1 time in dH<sub>2</sub>O for 2 min. with vigorous agitation at room temperature.
- 5 10. Transfer substrate to boiling (100 degrees C) dH<sub>2</sub>O for 10 sec. to denature DNA.
11. Dry substrate by centrifugation at 500x g for 1 min.
12. Store effector oligonucleotide-bound substrate at 4 °C prior to hybridization.

The nucleic acid sensor molecules can be, *e.g.*, those which possess either ligating or cleaving activity in the presence of a target molecule. (*See, e.g.*, Figures 2A and B for the  
10 ligater, Figure 5 for the cleaver).

In the case where it is desirable to immobilize an array of NASMs by direct attachment to a solid surface, the nucleic acid sensor molecules are bound to a solid substrate directly via their 3' termini. The attachment is accomplished by oxidation (using, *e.g.*, Na periodate) of the 3' vicinal diol of the nucleic acid sensor molecule to an aldehyde  
15 group. This aldehyde group will react with a hydrazide group to form a hydrazone bond. The hydrazone bond is quite stable to hydrolysis, etc., but can be further reduced (for example, by treatment with NaBH<sub>4</sub> or NaCNBH<sub>3</sub>). The use of adipic acid dihydrazide (ADH, a bifunctional linker) to derivatize an aldehyde surface results in a hydrazide-derivatized surface which provides a linker of approximately 10 atoms between the  
20 substrate surface and point of biomolecular attachment (see Ruhn et al., 1994; O'Shaughnessy, 1990; Roberston et al., 1972, Schluep et al., 1999; Chan et al., 1998). Preparation of a hydrazide-terminated surface via ADH treatment can be accomplished by treating an aldehyde-derivatized substrate according to the following protocol:

25 Protocol for ADH treatment of aldehyde substrate:

1. To 50 mL of 0.1 M phosphate buffer (pH 5) add 100-fold excess of adipic acid dihydrazide (ADH) relative to concentration of aldehyde groups on substrate surface.
2. Place substrate in a 50 mL tube containing the ADH in phosphate buffer and shake  
30 mixture for 2 hrs.
3. Remove the substrate and wash 4 times with 0.1 M phosphate buffer (pH 7).
4. Reduce free aldehyde groups on substrate surface by placing substrate in a 50 mL tube containing a 25-fold excess of NaBH<sub>4</sub> or NaCNBH<sub>3</sub> in 0.1 M phosphate buffer.
5. Shake the mixture for 90 min.

6. Wash 4 times with 0.1 M phosphate buffer (pH 7).
7. Store ADH-treated substrates in 0.1 M phosphate buffer (pH 7) at 4° C.

Preparation of the nucleic acid molecules for specific coupling to the ADH-terminated surface via their 3' termini can be accomplished according to the following protocol (see,

- 5 Proudnikov et al., 1996; Wu et al., 1996):

Protocol for Periodate oxidation of RNA:

1. Dissolve up to 20 micrograms of RNA in 5 microliters of H<sub>2</sub>O at 20° C.
2. Add 1 ML of 0.1 M NaIO<sub>4</sub> (~20-fold excess relative to RNA).
3. Incubate for 30 min. in a light-tight tube or enclosure.
- 10 4. Add 1 ML of 0.2 M Na sulphite (~2-fold excess relative to NaIO<sub>4</sub>) to stop reaction.
5. Incubate for 30 min. at room temperature.
6. Ethanol precipitate or use spin-separation column to recover oxidized RNA.

15 **Example 2. Selection for a nucleic acid sensor molecule selective for the estrogen receptor LBD**

A nucleic acid sensor molecule which is modulated by the estrogen receptor (ER) ligand binding domain (LBD) is obtained by *in vitro* selection methods to identify candidate nucleic acid sensor molecules that are modulated by an estrogen receptor LBD.

- 20 The full length gene for the estrogen receptor is known. One source of the full-length estrogen receptor clone is Acc. No. M12674 (see also Greene et al., Science 231:1150-54, 1986). The clone includes a 2092 nucleotide mRNA with the sequence presented in Table 3 below:

**Table 3**

25	1	gaattccaaa	attgtgatgt	ttcttgtatt	tttgatgaag	gagaaatact	gtaatgatca
	61	ctgtttacac	tatgtacact	ttaggccagc	cctttgtagc	gttatacaaa	ctgaaagcac
	121	accggaccgc	caggctccc	gggcagggcc	ggggccagag	ctcgcgtgtc	ggcgggacat
	181	gcgctgcgtc	gcctctaacc	tcgggctgtg	ctctttttcc	aggtggcccg	ccggtttctg
	241	agccttctgc	cctgcgggga	cacggtctgc	accctgcccc	cggccacgga	ccatgaccat
30	301	gaccctccac	accaaagcat	ctgggatggc	cctactgcat	cagatccaag	ggaacgagct
	361	ggagcccctg	aaccgtccgc	agctcaagat	ccccctggag	cggcccctgg	gcgaggtgta
	421	cctggacagc	agcaagccc	ccgtgtacaa	ctaccccgag	ggcgccgcct	acgagttcaa
	481	cgcgcgggcc	gcgcgcaacg	cgcaggctca	cggtcagacc	ggcctcccct	acggcccccg
	541	gtctgaggct	gcggcggttc	gctccaacgg	cctggggggg	ttccccccac	tcaacagcgt
35	601	gtctccgagc	ccgtgatgac	tactgcaccc	gccgccgcag	ctgtgcctt	tcctgcagcc
	661	ccacggccag	caggtgccct	actacctgga	gaacgagccc	agcggtaca	cggtgcgcga
	721	ggccggcccc	ccggcattct	acaggccaaa	ttcagataat	cgacgccagg	gtggcagaga
	781	aagattggcc	agtaccaatg	acaagggaag	tatggctatg	gaatctgcca	aggagactcg
	841	ctactgtgca	gtgtgcaatg	actatgcttc	aggctaccat	tatggagtct	ggtcctgtga
40	901	gggctgcaag	gccttcttca	agagaagtat	tcaaggacat	aacgactata	tgtgtccagc
	961	caccaaccag	tgcaccattg	ataaaaacag	gaggaagagc	tgccaggcct	gccggctccg
	1021	caaatgctac	gaagtgggaa	tgatgaaagg	tgggatacga	aaagaccgaa	gaggagggag
	1081	aatgttgaag	cacaagcgcc	agagagatga	tggggagggc	aggggtgaag	tggggtctgc
	1141	tggagacatg	agagctgcca	acctttggcc	aagccccgctc	atgatcaaac	gctctaagaa

1201 gaacagcctg gccttgctcc tgacggccga ccagatggtc agtgccttgt tggatgctga  
 1261 gccccccata ctctattccg agtatgatcc taccagaccc ttcagtgaag cttcgatgat  
 1321 gggcttactg accaacctgg cagacagggg gctgggtcac atgatcaact gggcgaagag  
 1381 ggtgccaggc tttgtggatt tgacctcca tgatcaggtc caccctctag aatgtgcctg  
 5 1441 gctagagatc ctgatgattg gtctcgtctg gcgctccatg gagcaccag tgaagctact  
 1501 gtttgcctct aacttgctct tggacaggaa ccagggaata tgtgtagagg gcatgggtga  
 1561 gatcttcgac atgctgctgg ctacatcatc tcggttccgc atgatgaatc tgcagggaga  
 1621 gatcttctg tgctcaaat ctattatctt gcttaattct ggagtgtaca catttctgtc  
 1681 cagcaccctg aagtctctgg aagagaagga ccatatccac cgagtccctg acaagatcac  
 10 1741 agacactttg atccacctga tggccaaggc aggcctgacc ctgcagcagc agcaccagcg  
 1801 gctggccccag ctctctctca tctctctcca catcaggcac atgagtaaca aaggcatgga  
 1861 gcatctgtac agcatgaagt gcaagaacgt ggtgccccct tatgacctgc tgcctggagat  
 1921 gctggacgcc caccgctac atcgccccc tagccgtgga ggggcatccg tggaggagac  
 1981 ggaccaaagc cacttggcca ctgcgggctc tacttcatcg cattccttgc aaaagtatta  
 15 2041 catcacgggg gaggcagagg gtttccctgc cacagtctga gagctccctg gc (SEQ ID  
 NO:9)

20 The polynucleotide encodes a polypeptide with the amino acid sequence presented  
 in Table 4 below:

Table 4

25 MTMTLHTKASGMALLHQIQGNELEPLNRPQLKIPLERPLGEVYLDSSKPAVYNYPEGAAYEFNAAAAANAQVYG  
 QTGLPYGPGSEAAAFSGNGLGGFPPLNSVSPSPMLLHPPQLSPFLQPHGQQVPYYLENESGYTVREAGPPA  
 FYRPNNDNRROGGRRERLASTNDKGSMAKESAKETRYCAVCNDYASGYHGVWVSCGCKAFFKRISIQGHNDYMCP  
 30 ATNQCTIDKNRRKSCQACRLRKCYEVGMMKGGIRKDRRGGRMLKHKRQRDDGEGRGEVGSAGDMRAANLWPSPL  
 MIKRSKNSLALSALTADQMVSALLDAEPPILYSEYDPTRPFSEASMMGLLTNLADRELHVHMINWAKRVPGFVDL  
 TLHDQVHLLECAWLEILMIGLVWRSMHPVKLLFAPNLLDRNQKCVBGMVEIFDMLLATSSRFRMMNLQGER  
 FVCLKSIILLNSGVYTFLSSTLKSLEEKDHIHRVLDKITDTLIHLMAKAGLTQQQHQRLLQLLILSHIRHMS  
 NKGMEHLYSMCKNVVPLYDLEMLDAHRLHAPTSRGGASVEETDQSHLATAGSTSSHSLQKYYITGEAEGFP  
 ATV (SEQ ID NO:10)

The gene encoding either full length ER or the ligand binding domain is cloned and  
 expressed in BL21 (DE3)-pLysS *E. coli* cells [Shiau AK, Barstad D, Loria PM, Cheng L,  
 Kushner PJ, Agard DA, Greene GL. Cell. 1998 Dec 23;95(7):927-37]. Human ER alpha  
 35 LBD (residues 297-554) are purified from estradiol-sepharose column by published  
 procedures [Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL.  
 Cell. 1998 Dec 23;95(7):927-37].

Selection of nucleic acid sensor molecules which are activated by ER-LBD not  
 bound to ligand: A library of up to  $10^{17}$  variants of *in vitro* synthesized (1  $\mu$ M) nucleic acid  
 40 sensor molecules is allowed to react with purified apo-ER-LBD at a final concentration of  
 1  $\mu$ M. Selection of catalytic nucleic acid sensor molecules and optionally, generation of an  
 optical NASM, is carried out by procedures outlined in prior examples and elsewhere  
 herein.

Selection of nucleic acid sensor molecules which are modulated by the ER-LBD-  
 45 Estradiol complex:

Stable complexes of ER-LBD and estradiol ligand are formed with from 1-10 equivalents of ligand. A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes is then allowed to react with purified ER-LBD-Estradiol at a final complex concentration of 1uM. Selection of allosterically activated ribozymes is carried out by procedures outlined the detailed description. Selection of nucleic acid sensor molecules which are modulated by the ER-LBD-Tamoxifen complex stable complexes of ER-LBD and tamoxifen ligand are formed with from 1-10 equivalents of ligand. A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes (1  $\mu$ M containing a plurality of potential target modulation domains and linker domains coupled to the catalytic domain of the ribozyme, is then allowed to react with purified ER-LBD-Tamoxifen at a final complex concentration of 1uM. Selection of catalytic nucleic acid sensor molecules target modulated ribozymes (nucleic acid sensor molecules) is carried out by procedures outlined in prior examples.

**Example 3. Selection for a library of nucleic acid sensor molecules which signal the presence of all known nuclear hormone receptor LBDs:**

A wide variety of nuclear hormone receptor ("NHR") ligand binding domains and their ligands, many of which are described in Table 5 below, are known for which a nucleic acid sensor molecule can be selected.

Table 5

Symbol	Description	Ligand
ATB3	nuclear receptor coactivator RAP250; peroxisome proliferator-activated receptor interacting protein; thyroid hormone receptor binding protein	thyroid hormone
AR	androgen receptor (dihydrotestosterone receptor; testicular feminization; spinal and bulbar muscular atrophy; Kennedy disease)	dihydroxytestosterone
CID	nuclear DNA-binding protein	
ESR1	estrogen receptor 1	estrogen
ESR2	estrogen receptor 2 (ER beta)	estrogen
ESRRA	estrogen-related receptor alpha	estrogen and TFIIB
ESRRB	estrogen-related receptor beta	estrogen and TFIIB
ESRRG	estrogen-related receptor gamma	estrogen and TFIIB
HNF4A	hepatocyte nuclear factor 4, alpha	
HNF4G	hepatocyte nuclear factor 4, gamma similar to retinoid X receptor, alpha (H. sapiens) similar to nuclear receptor subfamily 1, group D, member 1 (H. sapiens)	

NCOA1	nuclear receptor coactivator 1	Binds to steroid hormone receptors
NCOR1	nuclear receptor co-repressor 1	Thyroid hormone receptor without TH
NCOR2	nuclear receptor co-repressor 2	RXR without retinoic acid and THR without TH
NR0B1	nuclear receptor subfamily 0, group B, member 1	
NR0B2	nuclear receptor subfamily 0, group B, member 2	
NR1D1	nuclear receptor subfamily 1, group D, member 1	
NR1H2	nuclear receptor subfamily 1, group H, member 2	
NR1H3	nuclear receptor subfamily 1, group H, member 3	Interacts with RXR
NR1H4	nuclear receptor subfamily 1, group H, member 4	Bile acid, farnesol, or chenodeoxycholic acid
NR1I2	nuclear receptor subfamily 1, group I, member 2	pregnane
NR1I3	nuclear receptor subfamily 1, group I, member 3	androstane S
NR2C1	nuclear receptor subfamily 2, group C, member 1	
NR2C2	nuclear receptor subfamily 2, group C, member 2	
NR2E1	nuclear receptor subfamily 2, group E, member 1	
NR2E3	nuclear receptor subfamily 2, group E, member 3	
NR2F1	nuclear receptor subfamily 2, group F, member 1	
NR2F2	nuclear receptor subfamily 2, group F, member 2	
NR2F6	nuclear receptor subfamily 2, group F, member 6	Thyroid hormone
NR3C1	nuclear receptor subfamily 3, group C, member 1	glucocorticoid receptor, cortisol, corticosterone
NR3C2	nuclear receptor subfamily 3, group C, member 2	Aldosterone
NR4A1	nuclear receptor subfamily 4, group A, member 1	
NR4A2	nuclear receptor subfamily 4, group A, member 2	
NR4A3	nuclear receptor subfamily 4, group A, member 3	
NR5A1	nuclear receptor subfamily 5, group A, member 1	
NR5A2	nuclear receptor subfamily 5, group A, member 2	
NR6A1	nuclear receptor subfamily 6, group A, member 1	
PAX8	paired box gene 8	
PGR	progesterone receptor	progesterone
PPARA	peroxisome proliferative activated receptor, alpha	fenofibrate, clofibrate, WY14643
PPARBP	PPAR binding protein	binds to PPAR gamma
PPARD	peroxisome proliferative activated receptor, delta	WY1463
PPARG	peroxisome proliferative activated receptor, gamma	9-HODE, 13-HODE
PTHRI	parathyroid hormone receptor 1	parathyroid hormone
RARA	retinoic acid receptor, alpha	retinoic acid
RARB	retinoic acid receptor, beta	retinoic acid
RARG	retinoic acid receptor, gamma	retinoic acid
RORA	RAR-related orphan receptor A	

RORB	RAR-related orphan receptor B	
RORC	RAR-related orphan receptor C	
RXRA	retinoid X receptor, alpha	9-cis retenoic acid, complexes with activated VDR and THR
RXRB	retinoid X receptor, beta	9-cis retenoic acid, complexes with activated VDR and THR
RXRG	retinoid X receptor, gamma	9-cis retenoic acid, complexes with activated VDR and THR
SMAP	thyroid hormone receptor coactivating protein	activated THR
THRA	thyroid hormone receptor, alpha (avian erythroblastic leukemia viral (v-erb-a) oncogene homolog)	thyroid hormone
THRB	thyroid hormone receptor, beta (avian erythroblastic leukemia viral (v-erb-a) oncogene homolog 2)	thyroid hormone
TNRC11	trinucleotide repeat containing 11 (THR-associated protein, 230 kDa subunit)	activated THR
TRAP150	thyroid hormone receptor-associated protein, 150 kDa subunit	activated THR
TRAP240	thyroid hormone receptor-associated protein, 240 kDa subunit	activated THR
TRAP95	thyroid hormone receptor-associated protein, 95-kD subunit	activated THR
TRIP13	thyroid hormone receptor interactor 13	inactive THR
VDR	vitamin D (1,25- dihydroxyvitamin D3) receptor	vitamin D

N-terminally GST-tagged or N-/C-terminally His-tagged nuclear hormone receptor ligand binding domains, defined on the basis of structural homology are cloned and expressed in BL21 (DE3)-pLysS *E. coli* cells, or are cloned and expressed in standard  
5 baculovirus expression systems.

Human NHR LBDs (homologous to ER-alpha residues including the region aa 297-554) are purified from GSH-sepharose or nickel affinity columns by published procedures available from the manufacturers. LBDs are produced in a either a parallel or serial fashion and the purified proteins are stored in buffer containing 50 mM TrisHCl, 1 mM EDTA,  
10 1mM DTT and 50-250 NaCl/SCN salt, pH 7 to pH 8.5, 10% glycerol or other stabilizing agent. Protein sequence and MW is verified by electrospray LC-MS mass spectrometry.

1. Selection of nucleic acid sensor molecules which are modulated by NHR-LBD not bound to ligand: A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes (1  $\mu$ M) containing a plurality of potential target modulation domains and linker domains

coupled to the catalytic domain of the ribozyme, is allowed to react with purified apo-NHR-LBDs at a final concentration of 1  $\mu$ M LBD. Selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in the Detailed Description. Selections are carried out in parallel fashion. Selections can also be carried out in mixed pools of  
5 anywhere from 5-10 NHR-LBDs. In the final rounds of nucleic acid sensor molecule selection, the RNA pools may be separated into aliquots which may then be used to carry out *in vitro* selection against single NHR-LBD proteins to yield unique nucleic acid sensor molecules selective for multiple NHR-LBDs.

2. Selection of nucleic acid sensor molecules which are activated by ligand bound  
10 forms of NHR-LBDs.

Stable complexes of each NHR-LBD are formed with from 1-10 equivalents of ligand. A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes (1  $\mu$ M) containing a plurality of potential target modulation domains and linker domains coupled to the catalytic domain of the ribozyme is then allowed to react with purified NHR-LBD-Ligand  
15 complexes at a final complex concentration of 1  $\mu$ M. Selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in the Detailed Description and Example 1 below. Selections are carried out in parallel fashion. Selections can also be carried out in mixed pools of anywhere from 5-10 NHR-LBD-Ligand complexes. In the final rounds of nucleic acid sensor molecule selection, the RNA pools may be separated into  
20 aliquots which may then be used to carry out *in vitro* selection against single NHR-LBD-Ligand complexes to yield unique nucleic acid sensor molecules selective for all NHR-LBDs, bound to their ligands.

Thus, the invention provides an *in vitro* selection protocol against purified LBDs, bound to their ligands or not, for each known NHR. *In vitro* selections can be carried out  
25 with less than 1 mg of the purified forms of the LBDs. In addition the selection of nucleic acid sensor molecules can be done *in vitro* with mixed pools of LBD and subsequently deconvoluted after selection is complete. Alternatively, the final selection can be carried out with fractionally purified extracts containing a slight excess of recombinant LBD. In one embodiment the LBD is expressed in *E. coli*, or insect cell lines or mammalian cell  
30 lines. In another embodiment, the selection is carried out in cell free lysates in which the LBD is expressed in an *in vitro* transcription-translation procedure such as is described in the literature or can be purchased using common reagents from Roche or Promega. In another embodiment, the fractionated or purified LBDs are combined with known ligands, (as described above) agonist, antagonists or partial agonist/antagonists to form stable



complexes, and these complexes are then used for *in vitro* selection of nucleic acid sensor molecules. Upon interaction of the nucleic acid sensor molecule with the NHR-LBD, a signal will be generated detectable to an external monitoring device. In this manner, the activation state of any or all NHRs can be monitored *in vivo* or *in vitro* as will be described in detail in subsequent examples.

**Example 4. Selection for a nucleic acid sensor molecule selective for the Beta-2 adrenergic receptor.**

The full-length gene for the Beta-2 adrenergic receptor is described (Emorine et al., Proc. Natl. Acad. Sci. USA 84:6995-99, 1987) and is available at Acc. No. AAA88017.

The nucleic acid sequence is set forth in Table 6 below:

**Table 6**

```

GCACCGGAGCCCTAGCACCCGACAAGCTGAGTGTGCAGGACGAGTCCCCACCACACCC
ACACCACAGCCGCTGAATGAGGCTTCCAGGCGTCCGCTCGCGGCCCGCAGAGCCCCGCG
15 TGGGTCGCGCCGCTGAGGCGCCCCAGCCAGTGGCCTTACCTGCCAGACTGCGCGCCATG
GGGCAACCCGGGAACGGCAGCGCCTTCTTGCTGGCACCAATGGAAGCCATGCGCCGGAC
CAGCAGCTCAGCAGGAAAGGACGAGGTGTGGGTGGTGGGCATGGGCATCGTCATGTCT
CTCATCGTCTTGCCATCGTGTGTTGGCAATGTGCTGGTCATCACAGCCATTGCCAAGTTC
GAGCGTCTGCAGACGGTCACCAACTACTTCATCACTTCACTGGCCTGTGCTGATCTGGTC
20 ATGGGCCCTGGCAGTGGTGGCCCTTTGGGGCCGCCATATTCTTATGAAAATGTGGACTTTT
GGCAACTTCTGGTGCGAGTTTGGACTTCCATTGATGTGCTGTGCGTCACGGCCAGCATT
GAGACCCTGTGCGTGATCGCAGTGGATCGTACTTTGCCATTACTTCACCTTTCAAGTAC
CAGAGCCTGTGACCAAGAATAAGGCCCGGGTGATCATCTGATGGTGTGGATTGTGTCA
GGCCTTACCTCCTTCTTGCCCATTCAGATGCACTGGTACCGGGCCACCCACCAGGAAGCC
25 ATCAACTGCTATGCCAATGAGACCTGCTGTGACTTCTTACGAACCAAGCCTATGCCATT
GCCTCTTCCATCGTGTCTTCTACGTTCCCTTGGTGATCATGGTCTTCGTCTACTCCAGG
GTCTTTTCAAGGAGGCCAAAAGGCAGCTCCAGAAGATTGACAAATCTGAGGGCCGCTTCCAT
GTCCAGAACCTTAGCCAGGTGGAGCAGGATGGCGGGACGGGGCATGGACTCCGCAGATCT
TCCAAGTTCTGCTGAAGGAGCACAAAGCCCTCAAGACGTTAGGCATCATCATGGGCACT
30 TTCACCTCTGCTGGCTGCCCTTCTTCATCGTTAACTTGTGATGTGATCCAGGATAAC
CTCATCCGTAAGGAAGTTTACATCCTCCTAAATTGGATAGGCTATGTCAATTCTGGTTTC
AATCCCTTATCTACTGCCGAGCCAGATTTCAGGATTGCCTTCCAGGAGCTTCTGTGC
CTGCGCAGGTCTTCTTGAAGGCCTATGGGAATGGCTACTCCAGCAACGGCAACACAGGG
GAGCAGAGTGGATATCAGTGGAACAGGAGAAAGAAAATAAACTGCTGTGTGAAGACCTC
35 CCAGGCACGGAAGACTTTGTGGGCCATCAAGTACTGTGCCTAGCGATAACATTGATTCA
CAAGGGAGGAATTGTAGTACAAATGACTCACTGCTGTAAAGCAGTTTTTCTACTTTTAAA
GACCCCCCCCCCAACAGAACACTAAACAGACTATTAACTTGAGGGTAATAAACTTA
GAATAAAATTGTAAAAATTGTATAGAGATATGCAGAAGGAAGGGCATCCTTCTGCCTTTT
TTATTTTAAAGCTGTAAAAAGAGAGAAAACCTATTGAGTGATTATTTGTTATTTGTA
40 CAGTTCAGTTCCTCTTGTGATGGAATTGTAGTTTATGTCTAAAGAGCTTAGTCCTAG
AGGACCTGAGTCTGCTATATTTTCATGACTTTCCATGTATCTACCTCACTATTCAAGTA
TTAGGGTAATATATTGCTGTAATTGTATCTGAAGGAGATTTCCTTCTACACCCTT
GGACTTGAGGATTTTGTAGTATCTCGGACCTTTTCAGCTGTGAACATGGACTCTTCCCCAC
TCCTCTATTTGCTCACACGGGTATTTTAGGCAGGATTTGAGGAGCAGCTTCAGTTGT
45 TTTCCCGAGCAAAGGTCTAAAGTTTACAGTAAATAAAATGTTTGACCATG (SEQ ID NO:11)

```

The amino acid sequence of the polypeptide encoded by the nucleic acid sequence is set forth in Table 7 below:

**Table 7**

1 mgqpgngsaf llapngshap dhdtvqqrde vvvvgmgivm slivlaivfg nvlvitaiaak  
 61 ferlqgtvtny fitslacadl vmglavvpfg aahilmkmwt fgnfwcefwt sidvlecvtas  
 121 ietlcviavd ryfaitspfk yqslitknka rviilmwiv sgltstflpiq mhwyrathqe  
 181 aincyanetc cdfftnqaya iassivsfyv plvimfvys rvfgeakrql qkidksegrf  
 5 241 hvqnlsqveq dgrtghglrr sskfclkehk alktlgiimg tftlcwlpff invnihviqd  
 301 nlirkevyil lnwigyvnsf fnpliyrcsp dfriaqell clrrsslkay gngyssngnt  
 361 geqsgyhveq ekenkllced lpgtedfvgh ggtvpsdnid sggrncstnd sll (SEQ ID  
 NO:12)

10

The gene encoding either the full-length Beta-2 adrenergic receptor or cytoplasmic loop II or III or helix VII is cloned, expressed and purified from *E. coli* or baculovirus infected cells (Hampe, et al., J Biotechnol 77:219-234 (2000)) according to published procedures, and incorporated into detergent micelles to simulate the cellular milieu (Min, et al., J Biol Chem 268:9400-9404 (1993)). A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes (1  $\mu$ M) containing a plurality of potential target modulation domains and linker domains coupled to the catalytic domain of the ribozyme is allowed to react with purified Beta-2 adrenergic receptor at a final concentration of 1  $\mu$ M. Selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in Example 1.

20

Selection of nucleic acid sensor molecules which are activated by the butoxamine-Beta-2 adrenergic complex. 1:1 complexes of butoxamine and purified Beta-2 adrenergic receptor are formed and selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in Example 1.

Selection of nucleic acid sensor molecules which are activated by the isoproterenol-Beta-2 adrenergic complex is accomplished as follows. 1:1 complexes of isoproterenol and purified Beta-2 adrenergic receptor are formed and selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in Example 1 and the Detailed Description.

**Example 5: Selection for a library of nucleic acid sensor molecules which signal the presence of all known GPCRs**

30

The full-length gene sequences for over 400 G-protein coupled receptors ("GPCR") is known. GPCR polypeptides, peptides and peptide fragments are also known as set forth in Table 8 and Table 9 below. The entire gene, or peptides derived from these sequences, is N-terminally tagged or N-/C-terminally His-tagged and cloned, expressed, and purified, or synthesized by chemical means. All 400 plus GPCRs are produced in either a serial or parallel fashion and the purified proteins or peptides stored in buffer containing 50 mM Tris-HCl, 1 mM EDTA, 1 mM DTT, and 50-250 mM NaCl/SCN salt, pH 7 to pH 8.5, 10% glycerol or other stabilizing agent.

35

Table 8

	ref	NP_006134.1	G protein-coupled receptor 19 [Homo sapiens]
	ref	XP_049562.1	G protein-coupled receptor 19 [Homo sapiens]
5	ref	NP_004876.1	neuropeptide G protein-coupled receptor; n...
	ref	XP_011102.1	46228 [Homo sapiens] >gi 14723215 ref XP_0...
	ref	NP_057624.1	G protein-coupled receptor 72; reserved; G...
	ref	XP_011520.3	orexin receptor 2 [Homo sapiens]
	ref	NP_001517.1	orexin receptor 2 [Homo sapiens]
10	ref	NP_001471.1	galanin receptor 1; Galanin receptor [Homo...
	ref	NP_071429.1	neuropeptide FF 1; RFamide-related peptide...
	ref	XP_005747.4	tachykinin receptor 2 [Homo sapiens]
	ref	NP_001048.1	tachykinin receptor 2; Tachykinin receptor...
	ref	NP_000901.1	neuropeptide Y receptor Y2 [Homo sapiens] ...
15	ref	NP_003848.1	galanin receptor 2 [Homo sapiens] >gi 1365...
	ref	XP_004030.2	adrenergic, beta-2-, receptor, surface [Ho...
	ref	NP_000015.1	adrenergic, beta-2-, receptor, surface [Ho...
	ref	XP_001777.1	orexin receptor 1 [Homo sapiens]
	ref	XP_011871.3	neuropeptide FF 1; RFamide-related peptide...
20	ref	NP_001516.1	orexin receptor 1 [Homo sapiens]
	ref	NP_001041.1	somatostatin receptor 2 [Homo sapiens] >gi...
	ref	NP_001040.1	somatostatin receptor 1 [Homo sapiens] >gi...
	ref	NP_001044.1	somatostatin receptor 5 [Homo sapiens]
	ref	XP_012565.1	somatostatin receptor 5 [Homo sapiens]
25	ref	NP_115940.1	G protein-coupled receptor; G protein-coup...
	ref	XP_037563.1	G protein-coupled receptor [Homo sapiens]
	ref	NP_001050.1	tachykinin receptor 3; NK-3 receptor; neur...
	ref	XP_011942.1	prolactin-releasing hormone receptor [Homo...
	ref	XP_017624.1	G protein-coupled receptor 58 [Homo sapiens]
30	ref	NP_004239.1	prolactin-releasing hormone receptor [Homo...
	ref	NP_071640.1	histamine receptor H2; gastric receptor 1 ...
	ref	NP_055441.1	G protein-coupled receptor 58 [Homo sapiens]
	ref	XP_009594.2	somatostatin receptor 4 [Homo sapiens]
	ref	NP_003605.1	galanin receptor 3; galanin receptor, fami...
35	ref	NP_001049.1	tachykinin receptor 1, isoform long; Tachy...
	ref	XP_039747.1	opioid receptor, mu 1 [Homo sapiens] >gi 1...
	ref	NP_000905.1	opioid receptor, mu 1 [Homo sapiens]
	ref	XP_004341.2	53355 [Homo sapiens]
	ref	XP_052174.1	50635 [Homo sapiens]
40	ref	XP_052175.1	5-hydroxytryptamine (serotonin) receptor 4...
	ref	XP_052165.1	5-hydroxytryptamine (serotonin) receptor 4...
	ref	XP_052164.1	50636 [Homo sapiens] >gi 14732317 ref XP_0...
	ref	NP_000861.1	5-hydroxytryptamine (serotonin) receptor 4...
	ref	NP_001043.1	somatostatin receptor 4 [Homo sapiens]
45	ref	NP_000721.1	cholecystokinin A receptor [Homo sapiens] ...
	ref	NP_006670.1	putative opioid receptor, neuromedin K (ne...
	ref	NP_055442.1	G protein-coupled receptor 57 [Homo sapiens]
	ref	NP_000698.1	arginine vasopressin receptor 1B; arginine...
	ref	NP_001718.1	bombesin-like receptor 3 [Homo sapiens] >g...
50	ref	XP_040306.1	similar to SOMATOSTATIN RECEPTOR TYPE 2 (S...
	ref	NP_056542.1	tachykinin receptor 1, isoform short; Tach...
	ref	NP_001042.1	somatostatin receptor 3 [Homo sapiens] >gi...
	ref	NP_000722.1	cholecystokinin B receptor [Homo sapiens]
	ref	NP_000789.1	dopamine receptor D5; Dopamine receptor D1...
55	ref	NP_000612.1	5-hydroxytryptamine (serotonin) receptor 2...
	ref	NP_004215.1	G protein-coupled receptor 50 [Homo sapiens]
	ref	XP_010228.2	G protein-coupled receptor 50 [Homo sapiens]
	ref	NP_000907.1	oxytocin receptor [Homo sapiens]
	ref	XP_052179.1	oxytocin receptor [Homo sapiens] >gi 14725...
60	ref	NP_005192.1	chemokine (C-C motif) receptor 8; chemokin...
	ref	NP_000670.1	adrenergic, alpha-1B-, receptor; adrenergi...

	ref XP_046588.1	G protein-coupled receptor slt [Homo sapie...
	ref NP_000671.1	adrenergic, alpha-1A-, receptor; adrenergi...
	ref NP_110411.1	brain expressed G-protein-coupled receptor...
5	ref XP_003199.2	growth hormone secretagogue receptor [Homo...
	ref XP_017623.1	G protein-coupled receptor 57 [Homo sapiens]
	ref NP_005949.1	melatonin receptor 1A; melatonin receptor ...
	ref NP_115892.1	G protein-coupled receptor slt; melanin-co...
	ref NP_000903.1	opioid receptor, kappa 1; Opiate receptor,...
10	ref NP_000900.1	neuropeptide Y receptor Y1; Neuropeptide Y...
	ref XP_011716.2	similar to opioid receptor, kappa 1; Opiat...
	ref XP_011707.2	adrenergic, alpha-1A-, receptor [Homo sapi...
	ref XP_048085.1	adrenergic, alpha-1A-, receptor [Homo sapi...
	ref XP_048084.1	adrenergic, alpha-1A-, receptor [Homo sapi...
15	ref NP_000785.1	dopamine receptor D1 [Homo sapiens]
	ref XP_048082.1	adrenergic, alpha-1A-, receptor [Homo sapi...
	ref NP_003292.1	thyrotropin-releasing hormone receptor [Ho...
	ref NP_005305.1	gastrin-releasing peptide receptor [Homo s...
	ref XP_006335.4	dopamine receptor D2 [Homo sapiens]
20	ref XP_006334.3	dopamine receptor D2longer [Homo sapiens]
	ref NP_000786.1	dopamine receptor D2 [Homo sapiens] >gi 14...
	ref XP_036647.1	dopamine receptor D2 [Homo sapiens] >gi 14...
	ref XP_041422.1	similar to dopamine receptor D2 (H. sapien...
	ref NP_057658.1	dopamine receptor D2longer [Homo sapiens]
25	ref NP_000697.1	arginine vasopressin receptor 1A; V1a vaso...
	ref NP_002502.1	neuromedin B receptor [Homo sapiens]
	ref XP_018475.1	neuromedin B receptor [Homo sapiens]
	ref NP_062874.1	5-hydroxytryptamine receptor 7, isoform b;...
	ref NP_000730.1	cholinergic receptor, muscarinic 2; muscar...
30	ref NP_062873.1	5-hydroxytryptamine receptor 7, isoform d;...
	ref NP_000863.1	5-hydroxytryptamine receptor 7, isoform a;...
	ref NP_000667.1	adenosine A2b receptor [Homo sapiens] >gi ...
	ref NP_000675.1	beta-1-adrenergic receptor [Homo sapiens]
	ref NP_005963.1	pancreatic polypeptide receptor 1 [Homo sa...
35	ref NP_000732.1	cholinergic receptor, muscarinic 4; muscar...
	ref XP_039923.1	44527 [Homo sapiens]
	ref NP_000787.1	dopamine receptor D3 [Homo sapiens]
	ref XP_011027.3	dopamine receptor D3 [Homo sapiens]
	ref NP_061822.1	G protein-coupled receptor 14 [Homo sapiens]
40	ref NP_000731.1	cholinergic receptor, muscarinic 3; muscar...
	ref NP_000669.1	adrenergic, alpha-1D-, receptor; adrenergi...
	ref NP_005282.1	G protein-coupled receptor 17 [Homo sapien...
	ref XP_048332.1	similar to purinergic receptor (family A g...
	ref XP_041897.1	similar to G protein-coupled receptor 17 (...)
45	ref NP_005152.1	angiotensin receptor-like 1 [Homo sapiens]...
	ref NP_000016.1	adrenergic, beta-3-, receptor [Homo sapien...
	ref NP_001286.1	chemokine (C-C motif) receptor 1; macropha...
	ref NP_000902.1	opioid receptor, delta 1 [Homo sapiens]
	ref NP_005758.1	purinergic receptor (family A group 5) [Ho...
50	ref XP_006296.1	cholinergic receptor, muscarinic 4 [Homo s...
	ref XP_006058.1	similar to MUSCARINIC ACETYLCHOLINE RECEPT...
	ref NP_000729.1	cholinergic receptor, muscarinic 1; muscar...
	ref NP_000904.1	opiate receptor-like 1; opioid receptor-lik...
	ref NP_006630.1	cysteinyl leukotriene receptor 1 [Homo sap...
55	ref NP_001828.1	chemokine (C-C motif) receptor 3 [Homo sap...
	ref NP_000530.1	rhodopsin; rhodopsin (retinitis pigmentosa...
	ref NP_009154.1	putative G protein coupled receptor [Homo ...]
	ref XP_009882.2	adenosine A2a receptor [Homo sapiens]
	ref XP_045486.1	adrenergic, alpha-1D-, receptor [Homo sapi...
60	ref NP_000397.1	gonadotropin-releasing hormone receptor; g...
	ref NP_000666.2	adenosine A2a receptor; adenosine A2 recep...
	ref XP_002299.1	G protein-coupled receptor 45 [Homo sapiens]

	ref NP_003958.1	putative neurotransmitter receptor [Homo s...
	ref NP_000854.1	5-hydroxytryptamine (serotonin) receptor 1...
	ref XP_001811.2	opioid receptor, delta 1 [Homo sapiens] >g...
	ref NP_005950.1	melatonin receptor 1B; melatonin receptor ...
5	ref NP_001548.1	interleukin 8 receptor, beta [Homo sapiens]
	ref NP_064552.1	neuromedin U receptor 2 [Homo sapiens]
	ref NP_009158.1	G protein-coupled receptor 45 [Homo sapiens]
	ref NP_000625.1	interleukin 8 receptor, alpha; chemokine (...)
	ref XP_017622.1	putative neurotransmitter receptor [Homo s...
10	ref NP_036257.1	cholinergic receptor, muscarinic 5; muscar...
	ref NP_000743.1	leukotriene b4 receptor (chemokine recepto...
	ref NP_000788.1	dopamine receptor D4 [Homo sapiens]
	ref XP_006145.2	dopamine receptor D4 [Homo sapiens]
	ref XP_001543.1	G protein-coupled receptor 52 [Homo sapiens]
15	ref XP_009663.1	G protein-coupled receptor 8 [Homo sapiens]
	ref XP_007212.1	purinergic receptor (family A group 5) [Ho...
	ref NP_006165.1	neuropeptide Y receptor Y5 [Homo sapiens]
	ref NP_002522.1	neurotensin receptor 1 [Homo sapiens]
	ref XP_003692.2	5-hydroxytryptamine (serotonin) receptor 1...
20	ref XP_009612.2	neurotensin receptor 1 [Homo sapiens]
	ref NP_005277.1	G protein-coupled receptor 8 [Homo sapiens]
	ref NP_000515.1	5-hydroxytryptamine (serotonin) receptor 1...
	ref NP_001707.1	Burkitt lymphoma receptor 1, isoform 1; C...
	ref NP_005675.1	G protein-coupled receptor 52 [Homo sapiens]
25	ref NP_000852.1	histamine receptor H1; histamine receptor,...
	ref NP_000639.1	chemokine (C-C motif) receptor 2; chemokin...
	ref NP_116743.1	Burkitt lymphoma receptor 1, isoform 2; C...
	ref NP_006574.1	retinal pigment epithelium-derived rhodops...
	ref NP_031395.1	G-protein coupled receptor [Homo sapiens] ...
30	ref XP_009373.1	formyl peptide receptor-like 2 [Homo sapie...
	ref NP_076917.1	5-hydroxytryptamine (serotonin) receptor 5...
	ref XP_005280.2	G protein-coupled receptor 7 [Homo sapiens]
	ref XP_039818.1	G protein-coupled receptor 91 [Homo sapien...
	ref NP_006164.1	neuropeptide Y receptor Y6 (pseudogene) [H...
35	ref XP_012748.2	galanin receptor 1 [Homo sapiens] >gi 1476...
	ref NP_000862.1	5-hydroxytryptamine (serotonin) receptor 6...
	ref NP_005281.1	G protein-coupled receptor 15 [Homo sapien...
	ref XP_010406.1	angiotensin receptor 2 [Homo sapiens] >gi ...
	ref NP_001328.1	chemokine (C-X3-C) receptor 1; chemokine (...)
40	ref NP_002554.1	purinergic receptor P2Y, G-protein coupled...
	ref NP_004942.1	EBV-induced G protein-coupled receptor 2; ...
	ref NP_000672.1	adrenergic, alpha-2A-, receptor [Homo sapi...
	ref XP_005827.3	beta-1-adrenergic receptor [Homo sapiens]
	ref NP_005288.1	G protein-coupled receptor 24 [Homo sapien...
45	ref NP_002021.2	formyl peptide receptor-like 2 [Homo sapie...
	ref NP_005276.1	G protein-coupled receptor 7 [Homo sapiens]
	ref XP_010009.2	similar to somatostatin receptor-like prot...
	ref NP_003458.1	chemokine (C-X-C motif), receptor 4 (fusin...
	ref NP_000673.1	adrenergic, alpha-2B-, receptor [Homo sapi...
50	ref XP_051229.1	similar to C-X-C CHEMOKINE RECEPTOR TYPE 4...
	ref NP_001453.1	formyl peptide receptor-like 1; lipoxin A4...
	ref NP_006047.1	G protein-coupled receptor 66 [Homo sapien...
	ref NP_000856.1	5-hydroxytryptamine (serotonin) receptor 1...
	ref XP_048737.1	41064 [Homo sapiens]
55	ref NP_005499.1	chemokine (C-C motif) receptor 4; chemokin...
	ref NP_000638.1	chemokine (C-C motif) receptor 2; chemokin...
	ref XP_009664.1	opiate receptor-like 1 [Homo sapiens] >gi ...
	ref NP_005285.1	G protein-coupled receptor 21 [Homo sapien...
	ref XP_009561.2	34426 [Homo sapiens]
60	ref NP_004063.1	chemokine-like receptor 1 [Homo sapiens] >...
	ref XP_035769.1	chemokine-like receptor 1 [Homo sapiens]

	ref NP_000855.1	5-hydroxytryptamine (serotonin) receptor 1...
	ref NP_000045.1	arginine vasopressin receptor 2 [Homo sapi...
	ref NP_064445.1	opsin 1 (cone pigments), long-wave-sensiti...
	ref XP_048964.1	similar to PROBABLE G PROTEIN-COUPLED RECE...
5	ref NP_000665.1	adenosine A1 receptor [Homo sapiens] >gi 1...
	ref XP_011880.1	similar to pancreatic polypeptide receptor...
	ref NP_000676.1	angiotensin receptor 1; angiotensin recept...
	ref NP_000674.1	adrenergic, alpha-2C-, receptor [Homo sapi...
	ref XP_002705.3	G protein-coupled receptor 17 [Homo sapiens]
10	ref NP_000677.1	angiotensin receptor 2 [Homo sapiens]
	ref XP_004279.1	chemokine (C-C motif) receptor 6 [Homo sap...
	ref NP_000504.1	opsin 1 (cone pigments), medium-wave-sensi...
	ref XP_033840.1	similar to chemokine (C-C motif) receptor ...
	ref NP_002555.1	purinergic receptor P2Y, G-protein coupled...
15	ref XP_006367.1	purinergic receptor P2Y, G-protein coupled...
	ref NP_004358.1	chemokine (C-C motif) receptor 6; chemokin...
	ref XP_045851.1	opsin 1 (cone pigments), short-wave-sensit...
	ref NP_057641.1	orphan seven-transmembrane receptor, chemo...
	ref XP_003251.1	chemokine (C-C motif) receptor 9 [Homo sap...
20	ref NP_001699.1	opsin 1 (cone pigments), short-wave-sensit...
	ref XP_002838.5	similar to C-C CHEMOKINE RECEPTOR TYPE 11 ...
	ref NP_000570.1	chemokine (C-C motif) receptor 5; chemokin...
	ref NP_006632.2	chemokine (C-C motif) receptor 9, isoform ...
	ref NP_000859.1	5-hydroxytryptamine (serotonin) receptor 2...
25	ref NP_115942.1	putative purinergic receptor [Homo sapiens...
	ref NP_001497.1	G protein-coupled receptor 32 [Homo sapien...
	ref NP_061843.1	G protein-coupled receptor 85; super conse...
	ref NP_006555.1	G protein-coupled receptor [Homo sapiens] ...
	ref NP_065110.1	cysteinyl leukotriene CysLT2 receptor; cDN...
30	ref NP_004113.1	growth hormone secretagogue receptor [Homo...
	ref NP_055137.1	opsin 3 (encephalopsin) [Homo sapiens]
	ref XP_001515.3	opsin 3 (encephalopsin) [Homo sapiens] >gi...
	ref NP_005274.1	G protein-coupled receptor 5 [Homo sapiens...
	ref NP_061842.1	super conserved receptor expressed in brai...
35	ref NP_005291.1	G protein-coupled receptor 34 [Homo sapien...
	ref NP_037477.1	G protein-coupled receptor [Homo sapiens] ...
	ref XP_003126.1	chemokine binding protein 2 [Homo sapiens]...
	ref XP_007392.1	G protein-coupled receptor 65 [Homo sapiens]
	ref NP_005284.1	G protein-coupled receptor 20 [Homo sapiens]
40	ref NP_005287.1	G protein-coupled receptor 23 [Homo sapien...
	ref NP_009195.1	adrenomedullin receptor; G-protein-coupled...
	ref NP_003941.1	coagulation factor II (thrombin) receptor-...
	ref NP_000701.1	bradykinin receptor B1 [Homo sapiens]
	ref NP_000857.1	5-hydroxytryptamine (serotonin) receptor 1...
45	ref NP_057652.1	G-protein coupled receptor SALPR; somatost...
	ref XP_012745.1	histamine H4 receptor [Homo sapiens] >gi 1...
	ref NP_000858.1	5-hydroxytryptamine (serotonin) receptor 2...
	ref NP_003599.1	G protein-coupled receptor 65; T-cell deat...
	ref NP_001499.1	G protein-coupled receptor 39 [Homo sapien...
50	ref XP_007275.2	bradykinin receptor B1 [Homo sapiens]
	ref XP_006230.3	G protein-coupled receptor 72 [Homo sapien...
	ref XP_037208.1	histamine receptor H3 [Homo sapiens]
	ref XP_010168.2	arginine vasopressin receptor 2 [Homo sapi...
	ref NP_009163.1	histamine receptor H3; G protein-coupled r...
55	ref XP_037209.1	34432 [Homo sapiens] >gi 14786758 ref XP_0...
	ref NP_005273.1	G protein-coupled receptor 4 [Homo sapiens...
	ref NP_000668.1	adenosine A3 receptor [Homo sapiens] >gi 1...
	ref XP_001499.1	endothelial differentiation, sphingolipid ...
	ref NP_114142.1	G protein-coupled receptor 61 [Homo sapiens]
60	ref NP_002020.1	formyl peptide receptor 1 [Homo sapiens] >...
	ref XP_007108.2	endothelin receptor type B, isoform 1 [Hom...

	ref NP_000106.1	endothelin receptor type B, isoform 1 [Homo...
	ref NP_003982.1	endothelin receptor type B isoform 2 [Homo...
	ref XP_007276.2	bradykinin receptor B2 [Homo sapiens] >gi ...
	ref NP_000614.1	bradykinin receptor B2 [Homo sapiens] >gi ...
5	ref XP_001907.1	G protein-coupled receptor 25 [Homo sapiens]
	ref XP_051522.1	G protein-coupled receptor [Homo sapiens]
	ref NP_003476.1	G protein-coupled receptor 68; Ovarian can...
	ref NP_001498.1	G protein-coupled receptor 38 [Homo sapien...
	ref NP_073625.1	platelet ADP receptor [Homo sapiens] >gi 1...
10	ref NP_039226.1	olfactory receptor, family 10, subfamily H...
	ref XP_040869.1	similar to cannabinoid receptor 1 (brain) ...
	ref NP_004145.1	pyrimidinergic receptor P2Y, G-protein cou...
	ref NP_057167.1	central cannabinoid receptor, isoform a; C...
	ref NP_062813.1	seven transmembrane receptor BLTR2; leukot...
15	ref NP_149046.1	olfactory receptor, family 2, subfamily B...
	ref NP_000943.1	platelet-activating factor receptor [Homo ...
	ref XP_008010.2	similar to MELANOCYTE STIMULATING HORMONE ...
	ref NP_063950.1	olfactory receptor, family 2, subfamily S...
	ref NP_001831.1	central cannabinoid receptor, isoform a; C...
20	ref XP_003761.3	48895 [Homo sapiens] >gi 14724697 ref XP_0...
	ref NP_002377.2	melanocortin 1 receptor (alpha melanocyte ...
	ref XP_032638.1	neuromedin U receptor 2 [Homo sapiens]
	ref NP_005270.1	G protein-coupled receptor 1 [Homo sapiens]
	ref NP_005293.1	G protein-coupled receptor 37 (endothelin ...
25	ref NP_004092.1	coagulation factor II (thrombin) receptor...
	ref NP_005289.1	G protein-coupled receptor 25 [Homo sapiens]
	ref NP_001727.1	complement component 5 receptor 1 (C5a lig...
	ref XP_002667.1	41743 [Homo sapiens]
	ref XP_008392.1	CC chemokine receptor 10 [Homo sapiens] >g...
30	ref NP_001496.1	G protein-coupled receptor 30; chemokine r...
	ref NP_057686.1	CC chemokine receptor 10 [Homo sapiens]
	ref NP_001829.1	chemokine (C-C motif) receptor 7; Chemokin...
	ref XP_016412.1	28082 [Homo sapiens] >gi 14721034 ref XP_0...
	ref NP_006009.1	putative chemokine receptor; GTP-binding p...
35	ref NP_000136.1	follicle stimulating hormone receptor; ova...
	ref NP_002541.1	olfactory receptor, family 3, subfamily A...
	ref XP_002212.2	follicle stimulating hormone receptor [Homo...
	ref NP_004769.1	G protein-coupled receptor 44; chemoattrac...
	ref XP_015921.1	similar to putative chemokine receptor; GT...
40	ref XP_015923.1	putative chemokine receptor; GTP-binding p...
	ref NP_055380.1	olfactory receptor, family 1, subfamily A...
	ref NP_065133.1	putative G protein-coupled receptor 92 [Ho...
	ref XP_008716.1	melanocortin 4 receptor [Homo sapiens]
	ref NP_039225.1	olfactory receptor, family 11, subfamily A...
45	ref NP_002556.1	pyrimidinergic receptor P2Y, G-protein cou...
	ref NP_005903.1	melanocortin 4 receptor [Homo sapiens]
	ref NP_005297.1	G protein-coupled receptor 43 [Homo sapien...
	ref XP_012667.1	olfactory receptor, family 1, subfamily A...
	ref NP_112221.1	olfactory receptor, family 12, subfamily D...
50	ref NP_039227.1	olfactory receptor, family 10, subfamily H...
	ref NP_059976.1	olfactory receptor, family 7, subfamily C...
	ref NP_005279.1	G protein-coupled receptor 12 [Homo sapien...
	ref NP_001948.1	endothelin receptor type A [Homo sapiens]
	ref NP_005904.1	melanocortin 5 receptor [Homo sapiens] >gi...
55	ref NP_110401.1	prostate specific G-protein coupled recept...
	ref NP_004758.1	endothelin type b receptor-like protein 2 ...
	ref NP_003544.1	olfactory receptor, family 1, subfamily E...
	ref NP_000520.1	melanocortin 2 receptor; Melanocortin-2 re...
	ref NP_036492.1	olfactory receptor, family 1, subfamily F...
60	ref NP_002557.1	purinergic receptor P2Y, G-protein coupled...
	ref XP_030219.1	gonadotropin-releasing hormone receptor [H...

	ref XP_009029.4	purinergic receptor P2Y, G-protein coupled...
	ref NP_036284.1	endothelial cell differentiation gene 7; c...
	ref NP_039228.1	olfactory receptor, family 10, subfamily H...
5	ref NP_036484.1	olfactory receptor, family 1, subfamily A...
	ref XP_012668.1	olfactory receptor, family 1, subfamily A...
	ref NP_001983.1	coagulation factor II receptor precursor; ...
	ref NP_036501.1	olfactory receptor, family 2, subfamily F...
	ref NP_055694.1	putative G-protein-coupled receptor; G pro...
10	ref XP_004852.1	olfactory receptor, family 2, subfamily F...
	ref NP_000949.1	prostaglandin E receptor 4 (subtype EP4) (...)
	ref NP_072093.1	putative leukocyte platelet-activating fac...
	ref NP_110503.1	olfactory receptor, family 5, subfamily V ...
	ref XP_003907.1	coagulation factor II receptor precursor [...]
15	ref XP_004216.1	similar to olfactory receptor 89 (H. sapie...
	ref NP_112167.1	olfactory receptor, family 2, subfamily J...
	ref NP_002542.1	olfactory receptor, family 3, subfamily A...
	ref NP_036505.1	olfactory receptor, family 3, subfamily A...
	ref NP_005292.1	G protein-coupled receptor 35 [Homo sapiens]
20	ref NP_004045.1	complement component 3a receptor 1; comple...
	ref NP_005290.1	G protein-coupled receptor 31 [Homo sapiens]
	ref NP_005233.2	coagulation factor II (thrombin) receptor...
	ref NP_003546.1	olfactory receptor, family 1, subfamily G...
	ref XP_003671.3	coagulation factor II (thrombin) receptor...
25	ref NP_112163.1	olfactory receptor, family 7, subfamily A...
	ref NP_039229.1	olfactory receptor, family 10, subfamily C...
	ref NP_002539.1	olfactory receptor, family 1, subfamily D...
	ref XP_037263.1	similar to coagulation factor II (thrombin...
	ref NP_067647.1	leucine-rich repeat-containing G protein-c...
30	ref NP_065103.1	inflammation-related G protein-coupled rec...
	ref NP_055314.1	putative purinergic receptor [Homo sapiens...
	ref NP_036509.1	olfactory receptor, family 7, subfamily C...
	ref NP_036507.1	olfactory receptor, family 52, subfamily A...
	ref XP_035507.1	similar to NONE_RETURNED (H. sapiens) [Hom...
35	ref XP_004280.1	G protein-coupled receptor 31 [Homo sapiens]
	ref NP_003545.1	olfactory receptor, family 1, subfamily E...
	ref XP_036497.1	olfactory receptor, family 1, subfamily F...
	ref NP_061844.1	super conserved receptor expressed in brai...
	ref NP_063941.1	melanocortin 3 receptor [Homo sapiens]
40	ref NP_009091.1	olfactory receptor, family 2, subfamily H...
	ref XP_008678.2	olfactory receptor, family 1, subfamily D...
	ref NP_003543.1	olfactory receptor, family 1, subfamily D...
	ref XP_011731.3	similar to adrenergic, beta-3-, receptor (...)
	ref XP_009545.1	melanocortin 3 receptor [Homo sapiens]
45	ref NP_076403.1	G protein-coupled receptor 86 [Homo sapiens]
	ref XP_042200.1	G protein-coupled receptor 86 [Homo sapiens]
	ref NP_004711.2	endothelial differentiation, lysophosphati...
	ref NP_037440.1	platelet activating receptor homolog [Homo...
	ref NP_005217.1	endothelial differentiation, sphingolipid ...
50	ref NP_005295.1	G protein-coupled receptor 41 [Homo sapien...
	ref NP_005296.1	G protein-coupled receptor 42 [Homo sapiens]
	ref NP_115943.1	putative chemokine receptor [Homo sapiens]...
	ref NP_055381.1	olfactory receptor, family 1, subfamily D...
	ref XP_042826.1	luteinizing hormone/choriogonadotropin rec...
55	ref XP_010797.3	luteinizing hormone/choriogonadotropin rec...
	ref NP_000224.1	luteinizing hormone/choriogonadotropin rec...

Table 9

&gt;G(S)-1

mgclgnskte dqrneekaqr eankkiekql qkdkqvyrat hrllllgage sgkstivkqm



rilhvngfng eggeedpqaa rsnsdgekat kvqdiknnlk ealetivaam snlvppvela  
 npenqfrvdy ilsvmnvpdf dfppefyeha kalwedegvr acyersneyq lidcaqyfld  
 kidvikgady vpsdqdllrc rvltsgetet kfqvdkvnfh mfdvvggrde rrkwiqcfnd  
 vtaaiifvvas ssynmvired nqtnrlqeal nlfksiwnnr wlrtisvilf lnkqdllaek  
 5 vlagkskied yfpefarytt pedatpepge dprvtrakyf irdeflrist asgdgrhycy  
 phftcavdte nirrvmfndcr diiqrmhlrq yell (SEQ ID NO:13).

## &gt;G(S)-2

mgclgnskte dqrneekaqr eankkiekql qkdkqvyrat hrlllllgage sgkstivkqm  
 10 rilhvngfng eggeedpqaa rsnsdseka tkvqdiknnl keaetivaa msnlvppvel  
 anpenqfrvd yilsvmnvpd fdfppefyeh akalwedegv racyersney qlidcaqyfl  
 dkidvikgad yvpsdqdllr crvltsgife tkfqvdkvnf hmfdivgggrd errkwiqcfnd  
 dvtaiifvva sssynmvire dnqtnrlqea lnlfksiwnn rwlrtisvil flnkqdllaek  
 kvlagkskie dyfpefaryt tpedatpepg edprvtraky firdeflris tasgdgrhyc  
 15 yphftcavdt enirrvmfndc rdiqrmhlr qyell (SEQ ID NO:14).

## &gt;G(S)-3

mgclgnskte dqrneekaqr eankkiekql qkdkqvyrat hrlllllgage sgkstivkqm  
 20 rilhvngfng dekatkvqdi knnlkealet ivaamsnlvp pvelanpenq frvdyilsv  
 nvpdfdfppe fyehakalwe degvracyer sneyqlidca qyfldkidvi kqadyvpsdq  
 dllrcrvlts gifetkfvq kvnfhmfdiv ggqrderrki qcfndvtaii fvvasssynm  
 virednqtn lqealnlfs iwnnrwlrti svilflnkq dllaekvlag skiedyfpef  
 aryttpedat pepgedprvt rakyfirdef lristasgdg rhycyphftc avdtenirr  
 25 fndcrdiiqr mhlrqyell (SEQ ID NO:15).

## &gt;G(S)-4

mgclgnskte dqrneekaqr eankkiekql qkdkqvyrat hrlllllgage sgkstivkqm  
 rilhvngfng dsekatkvqd iknnlkeaie tivaamsnlv ppvelanpen qfrvdyilsv  
 30 mnvpdfdfppe efeyhakalw edegvracye rsneyqlidc aqyfldkidv ikqadyvpsd  
 qdllrcrvlt sgifetkfvq kvnfhmfdiv ggqrderrkw iqcfndvtai ifvvasssyn  
 mvirednqtn rlqealnlfs siwnnrwlrt isvilflnkq dllaekvlag kskiedyfpe  
 faryttpeda tpepgedprv trakyrirde flristasgd grhycyphftc cavdtenirr  
 35 vndcrdiiq rmhlrqyell (SEQ ID NO:16).

## &gt;G(s)-xl

meisgppfeie gsapagvddt pvnmdspia ldgppikvsg apdkreraer ppveeeaaem  
 egaadaaegg kvpspgygs aagaasadta araapaapad pdsgatpedp dsgtapadpd  
 sgafaadpds gaapaapadp dsgaapdap dpgsgaapda padpdagaap eapaapaae  
 40 traahvapaa pdagaptapa asatraaqvr raasaapasg arrkihlrpp speigaadpp  
 tprptrasaw rgksessrg rvydegvas sdddsgdes ddgtsgclrw fqhrnrrrr  
 kpqrnlrlnf lvqafggcfcg rsesppqkas rslkvkvpl aekrrqmrke alekraqkra  
 ekkrsklidk qlqdekmgym cthrlill (SEQ ID NO:17).

## &gt;g-olf

45 mgclgnskt tedqgvdeke rreankkiek qlqkerlayk athrlillga gesgstivk  
 qmrilhvngf npeekkqkil dirknvkdai vtivsamsti ippvplanpe nqfrsdyiks  
 iapitdfeyz qeffdhvkkk wdegvkcac ersneyqlid caqyflerid svslvdytpt  
 dqdllrcrvl tsigifetrfg vdkvnfhmfdiv vggqrderrk wiqcfndvta iiyvaacsy  
 nmvirednnt nrlresldlf esiwnnrwlrt tisiilflnk qdmlaekvla gkskiedyfp  
 50 eyanytvpd atpdagedpk vtrakffird lflristatg dgkhycyphf tcavdtenir  
 rvfndcrdii qrmhlkqyel l (SEQ ID NO:18).

## &gt;II

55 mgctlsaedk aaverskmid nrlredgeka arevklillg agesgstiv kqmkiheag  
 yseeckqyk avvysntiqs iiaiiiramgr lkidfgdsar addarqlfvl agaaegfmt  
 aelagvikrl wkdsqvgacf nrsreyqlnd saayylndld riagpnnyipt qgdvlrtrvk  
 ttgivethft fkdllhfkmdf vggqrserkk wihcfegvta iifcvalsdy dlvlaedeem  
 nrmhesmklf dsicnnkwft dtisiilflnk kdlfeekikk splticypey agsntyeeaa

ayiqcqfedl nkrkdtkeiy thftcatdtk nvqfvfdavt dviiknnlkd cglf (SEQ ID NO:19).

> I2

5 mgctvsaedk aaaerskmid knlredgeka arevkl111lg agesgkstiv kqmkiihedg  
yseeecrqyr avvysntiqs imaiivkamgn lqidfadpsr addarqlfal sctaeeggv1  
pddlsqvrr lwadhgvqac fgrsreyqln dsaayylndl eriaqsdyip tqgdv1rtrv  
kttgivethf tfkdlhfkmf dvvgqrserk kwihefegvt aiifcvalsa ydlvlaedee  
nmrmhesmkl fdsicnnkwf tdtssiilfln kkd1feekit hsp1ticfpe ytgankydea  
10 asyiqskfed lnrkrdtkei ythftcatdt knvqfvfdav tdviiknnlk dcglf (SEQ ID NO:20).

> I3

mgctlsaedk aaverskmid rnlredgeka akevkl111lg agesgkstiv kqmkiihedg  
ysedeckqyk vvvysntiqs iiairamgr lkidfgeaar addarqlfvl agsaeegvmt  
15 pelagvikrl wrdggvqac fgrsreyqlnd sasyylnldl risqsnyipt qgdv1rtrvk  
ttgivethft fkdlyfkmfd vggqrserkk wihcfegvta iifcvalsdyl dlvlaedee  
nmrmhesmkl dsicnnkwft etsiilflnk kdlfeekikr spl1ticypey tgsntyeeaa  
ayiqcqfedl nrrkdtkeiy thftcatdtk nvqfvfdavt dviiknnlke cgly (SEQ ID NO:21).

>G01

20 mgctlsaer aalerskaie knlkedgisa akdvkl111lg agesgkstiv kqmkiihedg  
fsgedvkqyk pvvysntiqs laaivramdt lgieygdker kadakmvdv vsmredtepf  
saellsammr lwgdsgiqec fnrsreyqln dsakyyldsl drigaadyqp teqdilrtrv  
kttgivethf tfknlhfrlf dvvgqrserk kwihefedvt aiifcvals g ydqv1hedet  
tnrmhesm1l fdsicnnkff idtsiilfln kkd1fgekik ksp1ticfpe ytgpn1yeda  
25 aayiqafes knrspnkeiy chmtcatdtn niqvvfdavt diiannlrg cgly (SEQ ID  
NO:22).

>g02

30 mgctvsaedk aaaerskmid knlredgeka arevkl111lg agesgkstiv kqmkiihedg  
yseeecrqyr avvysntiqs imaiivkamgn lqidfadpsr addarqlfal sctaeeggv1  
pddlsqvrr lwadhgvqac fgrsreyqln dsaayylndl eriaqsdyip tqgdv1rtrv  
kttgivethf tfkdlhfkmf dvvgqrserk kwihefegvt aiifcvalsa ydlvlaedee  
nmrmhesmkl fdsicnnkwf tdtssiilfln kkd1feekit hsp1ticfpe ytgankydea  
35 asyiqskfed lnrkrdtkei ythftcatdt knvqfvfdav tdviiknnlk dcglf (SEQ ID  
NO:23).

>G(T-1)

40 mgagasaek hsrelekk1k edaekdartv kl11lgages gkstivkqmk iihqdgysle  
eclefiiaiy gntlqsilai vramttniq ygdarqdda rklmhmadi eegtmpkems  
diigr1wkds g1qacferas eyqlndsagy ylsdlerlvt pgyvptegdv lrsrvk1tgi  
ietqfsfkd1 nfrmf1vvgg rserkkwihc fegvtciifi aalsaydmvl veddevnmh  
eslh1fnsic nhryfatts1 vlflnkkdvf fekikkah1s icfpdydgpn tyedagnyik  
vqflelnmrr dvkeiysht catdtqnvkf vfdavtdiii kenlkdcglf (SEQ ID NO:24).

45 >G(T-2)

mgsgasaedk elakrskele kklqedadke aktvkl111lg agesgkstiv kqmkiihqdg  
yspeeclef1k aiiygnv1qs ilaiiramtt lgidyaepsc addgrqlnnl adsieegt1mp  
pelvevirr1 wkdg1vqacf eraaeyqlnd sasyylnqle ritdpeylps eqdv1rsvk  
ttgi1etkfs vkdl1nrmfd vggqrserkk wihcfegvtc iifcaalsay dmvlveddev  
50 nmrmhesl1hf nsicnhkffa atsiv1flnk kdlfeekikk vhsicfpey dgnnsy1dag  
nyikeqf1dl nmrkdvkeiy shmtcatdtq nvkf1vfdavt diiikenlkdcglf (SEQ ID NO:25).

>G(Z)

55 mvflsgnasd ssnctqppap vniskail1g vilgg1l1fg vlg1nilvils vachrh1hsv  
thyyivn1lav adll1tstvl pfsaifev1g ywafgrvfc1n iwaavdv1cc tasimg1cii  
sidryigvsh plryptiv1tq rrglmallcv wals1visig plfgwrq1pap edeticq1ne  
epgyv1fsal gsfylplai1 lvm1crvyvv akresrg1ks glktdksdse qvtlrih1kn  
apag1sgmas aktkthfs1v l1kfsrekka aktlgiv1vgc fvlcw1pffl vmpigsff1pd

fkpsetvfki vfwlgylncs inpiiyccss qefkkafqnv lriqclcrkq sskhalgytl  
 hppsqavegq hkdmvripvg sreafygisr tdgvcewkff ssmprgsari tvskdqssct  
 tarvrksfsl qvcccvpst psldknhqvp tikvhtisls engeev (SEQ ID NO:26).

## 5 &gt;G(Q)

macclseeak earrindeie rqlrrdkrda rrelkl11lg tgesgkstfi kqmriihgsg  
 ysdedkrqft klvyqnifta mqamiramdt lkipykyehn kahaqlvrev dvekvsafen  
 pyvdaikslw ndpgigecyd rrreyqlsds tkyylnldlr vadpaylptq qdvlrvrvpt  
 tgiieypfdl qsvifrmvdv ggqrserrrk ihcfenvtsi mflvalseyd qvlvesdnen  
 10 rmeeskalfr tiitypwwfn ssvilflnkk dlleekimys hlvdyfpeyd gpqrdaqaar  
 efilkmfvdl npdsdkiiys hftcatdten irfvfaavkd tilqlnlkey nav (SEQ ID NO:27).

## &gt;G(Y-11)

mtlesmmacc lsdevkeskr inaeiekqlr rdkrdarrel kl111gtges gkstfikqmr  
 15 iihgagysse dkrqftklvy qniftamgam irametlkil ykyeqnkana llirevdvek  
 vttfehgysv aikt1wedpg iqecydrre yqlsdsakyy ltdvdriatl gylptqqdvl  
 rvrvpttgii eypfdlenii frmvdvggqr serrkwihcf envtsimflv alseydqvlv  
 esdnenrmee skalfrtiit ypwfqnssvi lflnkkdlle dkilyshlvd yfpefdgppqr  
 daqaarefil kmfvdlnpds dkiyyshftc atdtenirfv faavkdtilq lnlnkeynlv (SEQ ID NO:28)

20

## &gt;G(Y-12)

msgvvrtlsr cllpaeagga rerragsgar daerearrs rdidallare rravrrlvki  
 25 lllgagesgk stflkqmrii hgrefdqkal lefrdtifdn ilkgsrvlvd ardklgipwq  
 ysenekhgms lmafenkagl pvepatfqly vpalsalwrd sgireafsrr sefqlgesvk  
 yfldnldrig qlnyfyskqd illarkatkg ivehdfvikk ipfkmdvvgg qrsqrqkwf  
 cfdgitsilf mvssseydgv lmedrrtnrl vesmnifeti vnnklffnvs iilflnkmld  
 lvekvktvsi kkhfpdfrgd phqledvqry lvqcfdrkrr nrskplfhhf ttaidtenvr  
 fvfhavkdti lqenlkdiml q (SEQ ID NO:29).

## 30 &gt;G(Y-13)

madflpsrsv lsvcfpgc11 tsgeaeqqrk skeidkclsr ektyvkr1vk illlgagesg  
 kstflkqmrii ihgqdfdgrra reeifrptiys nvikgmrvlv dareklhipw gdnsngqhg  
 kmmsfdtrap maaqgmvetr vflqylpair alwadsgiqn aydrrefql gesvkyfldn  
 35 ldklgepdyi psqqdillar rptkgiheyd feiknvpfkm ldvggqrser krwfecfsv  
 tsilflysss efdqvlmedr ltnrltesln ifetivnnrv fsnvsilfl nktdlleeqv  
 qivsikdyfl efegdpheclr dvqkflvecf rnkrddqqqk plyhhfttai ntenirlvfr  
 dvkdtihlhdn lkqlmlq (SEQ ID NO:30).

## &gt;G(Y-14)

magccclsae ekesgrisae ierqlrrdkk darrelkl11 lgtgesgkst fikqmriihg  
 40 sgysdedrkg ftklvyqnif tamqamiram dtlrigyvce qnkenaqiir evevdksml  
 sreqveaikq lwqdpigiec ydrreyqls dsakyy1tdi driatpsfvp tqgdvlrvrv  
 pttgieypf dleniifrmv dvggqrser kwhcfesvt siiflvalse ydqvlaecd  
 enrmeskal fktiitypwf lnssvilfln kkdllleekim yshlisypfe ytgpkqdvra  
 45 ardfilklyq dqnpdkekvi yshftcatdt dnirfvfaav kdtlqlnlr efnlv (SEQ ID NO:31).

## &gt;G(Y-15)

marsltwrcc pwc1tedeka aarvdqeinr illeqkkqdr gelkl111gp gesgkstfik  
 50 qmriihgagy seeerkgrfp lvyqnifvsm ramieamerl qipfsrpesk hhaslvmsqd  
 pykvttfekar yaaamqlwr dagiracyer rrefhllda vyy1shleri teegyvp1taq  
 dvlrsmptt gineycfsvg ktnlrvdvvg gqkserrkwi hcfenviali ylaslseydq  
 cleennqenr mkeslalfgt ilelpwfkst svilflnktl ileekiptsh latyfpsfqq  
 pkqdaaakr fildmytrmy tgcvdgpegg kkgarsrrlf shytcattatq nirkvfkdv  
 55 dsvlarylde in11 (SEQ ID NO:32).

## &gt;Gusducin

mfvdvggqrse rkkwihcfeg vtciifcaal saydmvlved eevnrmhesl hlfnsicnhk  
 yfsttsivlf lnkkdifqek vtkvhlscif pey1tgpn1te dagnyiknqf ldlnlkkedk

eiyshmtcat dtgnvkfvfd avtdiiken lkdcglf (SEQ ID NO:33).

1. Selection of nucleic acid sensor molecules which are activated by GPCRs not bound to ligand:

- 5        A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes is allowed to react with purified GPCRs at a final concentration of 1  $\mu$ M GPCR. Selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in prior examples. Selections are carried out in parallel fashion. Selections can also be carried out in mixed pools of anywhere from 5-10 GPCRs. In the final rounds of nucleic acid sensor molecule selection, the RNA pools may separated into aliquots which may then be used to carry out *in vitro* selection against single GPCR proteins to yield unique nucleic acid sensor molecules selective for all 400 plus GPCRs.

2. Selection of nucleic acid sensor molecules which are activated by ligand bound GPCRs.

- 15        Stable complexes of each GPCR and corresponding -ligand are formed with from 1-10 equivalents of ligand. Selection of catalytic nucleic acid sensor molecules is carried out by procedures outlined in the Detailed Description. Selections are carried out in parallel fashion. Selections can also be carried out in mixed pools of anywhere from 5-10 GPCR-ligand complexes. In the final rounds of nucleic acid sensor molecule selection, the RNA pools may separated into aliquots which may then be used to carry out *in vitro* selection against single GPCR-ligand complexes to yield unique nucleic acid sensor molecules selective for all GPCR-ligand complexes.

- 25        Selection of nucleic acid sensor molecules using peptide fragments of GPCRs. Nucleic acid sensor molecules that specifically recognize conformational isoforms of GPCRs that are revealed upon ligand binding can also be selected for using the methods described herein.

- 30        Molecular cloning studies have identified over 400 human GPCRs, and have identified the ligands for 120. GPCRs consist of three domains: an extracellular N-terminus, a central domain of seven trans-membrane helices, and a cytoplasmic C-terminus. Activation of GPCRs is induced by ligand binding, which causes a conformational change in the receptor transmitting a signal across the plasma membrane to intracellular members of a signaling pathway. This method provides for generation of unique biosensors for each GPCR.

A library of up to  $10^{17}$  variants of *in vitro* synthesized ribozymes is allows to react with peptide fragments of the GPCRs comprising regions of the GPCR that are exposed upon activation. Nucleic acid sensor molecules which recognize these domains are then capable of recognizing them within the context of the full length protein and hence  
5 recognize the activated state of the GPCR. Examples of the use of peptide fragments to generate nucleic acid sensor molecules which recognize the full length protein are known in the art and are incorporated herein. See, for example, data on nucleic acid sensor molecule selection and recognition of HIV rev peptide and full length protein [Michael Robertson, 2001, University of Texas, Austin, Ph.D. Dissertation]. In the case of HIV rev, unique  
10 peptide sequences are recognized both as free peptides and in the context of the full protein.

Nucleic acid sensor molecules specific for GPCRs are generated by *in vitro* selection for recognition of peptide fragments of the GPCRs comprising regions of the GPCR that are exposed to the inside face of the plasma membrane when ligand binds to the GPCR. Exemplary suitable GPCR peptide fragments are presented in Table 9. Nucleic acid sensor  
15 molecules which recognize these peptides are then capable of recognizing them within the context of the full length protein and hence recognize the activated state of the GPCR.

**Example 6. Nucleic acid sensor molecule specific to phosphodiesterase (PDE) for target validation**

20 Nucleic acid sensor molecules are generated to be specific to various subclasses of PDEs are used for understanding the role of PDE subclasses in the molecular pathology of disease, and as PDE target validation. For example, nucleic acid sensor molecules which are modulated by each of four PDE4 subtypes have specific utility in understanding the role  
25 of PDE4 in human disease. The four subclasses of PDE4 are differentially localized between cell type and also the PDE4 isozymes differ with respect to their intracellular localization. This differential localization, together with the transcriptional regulation and post-translational modification, controls the cAMP level in cells in response to the cells' environment (Muller, Engels et al. 1996).

30 The cDNAs for four PDE4 subtypes are cloned from human blood leukocyte cDNA library as described (Wang, Myers et al. 1997). Each subclass of PDE4 is expressed as recombinant protein fused with a His-tag in *E. coli* or SF9 insect cells (Richter, Hermsdorf et al. 2000) (Wang, Myers et al. 1997). The expressed proteins are purified through  $\text{Ni}^{++}$  columns according to established procedures. Catalytic nucleic acid sensor molecules  
35 modulated by the four subclasses of PDE4 are then selected as described above. The

nucleic acid sensor molecules are tested for their subclass specificity, by determining the switch factor.

Tissue samples from different organs can be prepared, and the cell extract can be tested against a panel of PDE4 subclass-specific nucleic acid sensor molecules to determine the protein level of each PDE in the organ. Thus, one can obtain more precise information about PDE4 levels relative to methods based on measuring the mRNA level (Bloom and Beavo 1996) (Oberholte, Ratzliff et al. 1997) (Nagaoka, Shirakawa et al. 1998).

Likewise, different classes of PDE (PDE1-11) are expressed in a tissue-specific manner and play different physiological roles (Conti 2000), and the subcellular localization of PDE regulates their activity. Accordingly, the nucleic acid sensor molecule can be used to determine the subcellular localization of each PDE from fractionated cell extracts (Bolger, Erdogan et al. 1997), or in situ hybridization technique (Sirinarumitr, Paul et al. 1997).

The nucleotide sequences of a cAMP-dependent PDE nucleic acid sensor molecule and cGMP-dependent nucleic acid sensor molecule are presented in Table 10. Allosteric domains are shown in bold font and the cleavage site nucleotide is underlined. CGMP modulated NASMs that are configured for homogeneous, solution based fluorescence assays (FRET) are shown the Figure 62. Multiplexed camp and cGMP-modulated FRET-sensor NASM-based assays are shown in Figure 65B. The optical NASMs modulated by cAMP and cGMP are used in PDE assays as described in detail below.

Table 10

cAMP-Hammerhead RNA seq:

5' - GGGC GAC CC UGA UGA GCC UGU GGA AAC AGA CGU GGC ACA UGA CUA CGU CGA  
AAC GGU GAA AGC CGU AGG UUG CCC -3' (SEQ ID NO:34)

cGMP-Hammerhead RNA seq:

5' - GGGC GAC CC UGA UGA GCC CUG CGA UGC AGA AAG GUG CUG ACG ACA CAU CGA  
AAC GGU GAA AGC CGU AGG UUG CCC -3' (SEQ ID NO:35)

The cAMP and cGMP-dependent nucleic acid sensor molecules were added to a solution containing various amounts of PDE and the corresponding cyclic nucleotide (cAMP or cGMP). A decreasing amount of the cyclic nucleotide was found to correspond to the increasing amount of PDE. These results demonstrate that the cyclic nucleotide-dependent nucleic acid sensor molecules can be used to measure PDE activity.

The calmodulin activated PDE specific for cGMP was used under the following enzyme assay conditions: Assay conditions (10 mM Tris, 20 mM MgCl<sub>2</sub>, 100 μM CaCl<sub>2</sub>) 200 nM cGMP-NASM; the assay is quenched with 0.1% SDS and then 200 nM cGMP-NASM is added to the mixture. Remaining cGMP was determined by the amount of  
5 conversion of nucleic acid sensor molecule to product using a gel based radioactive product-release (gel-shift) assay format. The nucleic acid sensor molecule is active in a variety of formats and is not inhibited by GMP produced by the components of the PDE assay. Similarly, the cAMP nucleic acid sensor molecule is active in a variety of assay formats and is not inhibited by GMP produced by the components of the PDE assay. The conversion of  
10 cGMP to GMP is followed by the optical NASM formats described in Figures 62-72.

*High Throughput Screening (HTS) assays using cAMP-dependent PDE nucleic acid sensor molecules:*

A cAMP-dependent nucleic acid sensor molecule is used in HTS assays for PDEs  
15 (PDE1, PDE2, PDE3, PDE4, PDE7, PDE8, PDE10, and PDE11). Similarly, cGMP-dependent-nucleic acid sensor molecules can be used in HTS assays for PDEs (PDE5, PDE9, PDE10, and PDE11). Representative cAMP-dependent and cGMP-dependent PDE nucleic acid sensor molecules are shown above, and in Figures 62-72 for all solution and chip-based NASM assay configurations.

20 Each class of PDE can be isolated from human tissue (Ballard, Gingell et al. 1998), or expressed as recombinant proteins in various system (*e.g.* *E coli*, SF9 cells). Thus, the nucleic acid sensor molecule monitors the PDE activity in the presence and the absence of candidate drugs. For example, PDE and its substrate (*i.e.*, cAMP and/or cGMP) are incubated at predetermined durations in a multiwell chamber (*e.g.*, 96, 384 well) with  
25 various concentration of compounds for screening, and the reaction is terminated by changing the buffer conditions (*e.g.*, addition of sufficient amount of EGTA, shifting buffer pH), or by separating enzyme and substrate (*e.g.*, filtration). Next, the nucleic acid sensor molecules are added to measure the altered concentration of the substrate, cAMP and cGMP. Alternatively, optical nucleic acid sensor molecules can be added without  
30 terminating the PDE activity. CAMP and cGMP modulated these HTS assays, as described in Figure 62 and Figure 63.

cAMP- or cGMP-dependent nucleic acid sensor molecules can also be used to characterize the IC<sub>50</sub> of the drug *in vitro*. A PDE assay is performed with serial dilutions of a compound of interest. Purified PDE or, alternatively, soluble extract from cells

(Moreland, Goldstein et al. 1998) can be used for the assay. The assay can be performed as described herein.

Alternatively, cAMP- or cGMP-dependent nucleic acid sensor molecules (Figures 62, 33, and 64) are used to characterize the IC<sub>50</sub> values of drug candidate *in vitro* by analyzing cAMP- or cGMP synthesized by adenylyl and guanylyl cyclases. Adenylyl and guanylyl cyclase assays are set up with serial dilution of a compound of interest. Membrane fractions containing adenylyl and guanylyl cyclases are used for the assay. The assay can be setup as described in the literature using ATP or GTP as the substrate.

*Competitive assays using PDE nucleic acid sensor molecules:*

Nucleic acid sensor molecules are generated that interact with the active sites of PDEs. PDE4 proteins are obtained as described above. The nucleic acid sensor molecules are selected against PDE4 with negative selection in the presence of PDE4 complexed with subnanomolar inhibitor (*e.g.*, Rolipram). Thus, the nucleic acid sensor molecule is modulated by free, uncomplexed PDE4, the PDE nucleic acid sensor molecules compete for PDE binding with inhibitors.

The direct inhibition by the nucleic acid sensor molecules can be tested using commercially available PDE assay kits (Amersham SPA assay kit for cAMP, Molecular Devices HEPP cAMP assay kit). In drug screening, the competition is performed by monitoring the signal from the nucleic acid sensor molecules in the presence of various inhibitors. Purified PDE or soluble cell extract from appropriate source (*e.g.*, Wistar rat brain (Andersson, Gemalmaz et al. 1999) ) is incubated with nucleic acid sensor molecules (100 nM) in the presence and the absence of compounds in 10 mM Tris buffer pH 7.5 containing 10 mM MgCl<sub>2</sub>. The changes in the initial rate of each nucleic acid sensor molecule response in the presence and the absence of the drug can be monitored in homologous system. Multiple PDEs can be tested against a same compound in the same well. This assay is expanded if desired to determine the tissue specific interaction of each class of PDE and any compounds.

*Cell-based assays using cyclic nucleotide-dependent PDE nucleic acid sensor molecules*

Nucleic acid sensor molecules are used to monitor the cellular cAMP and cGMP level in response to the injection of drugs in tissue or rat cell lines. For example, strips of human corpus collasum (HCC) tissue or rat HCC cell lines (N1S1 and McA-RH7777 cells) can be incubated in the presence and absence of a drug against PDE5 (Min, Kim et al. 2000) (Arora, de Groen et al. 1996), and the cGMP specific nucleic acid sensor molecule can be



used to measure the amount of cGMP in soluble extract from the tissue or cell sample as described above.

Alternatively, the cAMP, and cGMP-dependent nucleic acid sensor molecules are incorporated into a reporter-gene plasmid as described above. This construct is introduced  
5 in cell lines by standard transfection (*e.g.* lipid-mediated transfection, calcium-phosphate co-precipitation, microinjection, electroporation, retroviral infection). The level of cGMP or camp in the cell is measured by the expression of the reporter gene.

*Class specific PDE assay:*

10 Nucleic acid sensor molecules are selected for the catalytic domains of each class of PDE1-11 are prepared. These nucleic acid sensor molecules are then used for target validation as described above.

Alternatively, the nucleic acid sensor molecules are used in competitive inhibition assays. Competitive nucleic acid sensor molecules are used in *in vitro* assays to screen  
15 compounds against multiple PDEs in multiplex assays, as described above.

**Example 7. Pharmacokinetics studies using nucleic acid sensor molecules**

Nucleic acid sensor molecules modulated by drug leads or drug compounds used in preclinical testing and clinical trial for pharmacokinetics studies are selected and identified  
20 as described in the Detailed Description. A human serum sample with or without the administration of a drug or other therapeutic agent is prepared (Berzas Nevado, Rodriguez Flores et al. 2001). The nucleic acid sensor molecule is added to the sample. The nucleic acid sensor molecule is then used in optical or PCR based detection methods as described in later examples, thereby quantifying the drug concentration in the whole serum or extract  
25 from the serum.

The nucleic acid sensor molecule modulated by various drugs or leads can also be used to determine the drug distribution in an animal model system. For example, a drug can be administrated in animals (*e.g.*, Sprague Dawley rats, New Zealand white rabbit) by IV or orally (Andersson, Gemalmaz et al. 1999) (Jeremy, Ballard et al. 1997). At various time  
30 intervals after drug administration, the animal is sacrificed. Various organs are tested for the drug distribution by in situ hybridization using the drug-dependent nucleic acid sensor molecule. Alternatively, each organs/serum is prepared for pharmacokinetic studies.

**Example 8. Cell-permeability studies using nucleic acid sensor molecules**

Nucleic acid sensor molecules against a test compound are used to test cell permeability of the compound. These nucleic acid sensor molecules can be incorporated into a reporter gene construct, if desired, to make a drug-sensitive reporter gene system as described above. This construct is introduced in established cell lines (*e.g.* HELA cells, 293 cell, CHO cell). The cells are cultured in various concentrations of drug in media, and the expression of the reporter gene is monitored.

**Example 9. Expression and purification of recombinant MAPKs**

All protein ORFs were obtained from Upstate Biotech, and cloned into pRSET (Invitrogen), except for ERK which was cloned directly downstream from a hexahistidine tag into pRSET from which the leader sequence had been removed. Constructs were sequence verified before transformation into Rosetta *plys*(S) cells (Novagen). After growth and induction as described below, cells were washed in 10 ml PBS per liter of culture and resuspended in 7.5 ml Lysis Buffer (500 mM KCl, 20 mM Tris-Cl, pH 8.0, 10% glycerol, 0.5% NP-40, supplemented with 1 Complete EDTA-Free Protease Inhibitor tablet (Roche) per 50 ml) per liter of culture. Cells were frozen in liquid nitrogen and stored at -80 °C. Lysis and clarification was accomplished by rapid thawing in a 40 °C bath for 10 minutes, incubation at 4 °C for thirty minutes, and centrifugation for 60 minutes at 100,000 x g. As a standard first step, metal chelate affinity chromatography (MCAC) was performed as follows: cell lysate was diluted 1:2 with Buffer A (500 mM KCl, 20 mM Tris-Cl, pH 8.0, 10% glycerol) and applied to a 5 ml HiTrap Chelating column (Amersham Biosciences) charged with nickel. The column was washed with 10 column volumes of Buffer A, and eluted stepwise with 10 column volumes of Buffer A plus 16, 56, 167 and 500 mM imidazol, pH 8.0 at a flow rate of 1.5 ml/min and collecting 5 ml fractions. MAPKs eluted in either the 167 or 500 mM imidazol fractions. After the purification described below, proteins were essentially pure as determined by silver staining of SDS-PAGE gels.

For purification of ERK, cells were grown to an OD<sub>600</sub> of 0.7 at 37 °C and induced with 1 mM IPTG for 2 hours at 37 °C. The MCAC fraction was diluted tenfold with Buffer B (20 mM HEPES 8.0, 1 mM DTT, 1 mM EDTA, 10% glycerol) and loaded onto a 5 ml HiTrap Q column (Amersham Biosciences) previously equilibrated with Buffer B plus 50 mM KCl. ERK was eluted in a 10 column gradient to Buffer B plus 500 mM KCl, at a flow rate of 1.5 ml/min, taking 2.5 ml fractions.

For purification of Jnk1, stationary phase culture was diluted to OD<sub>600</sub> of 0.7 and grown at 37 °C for one hour, and induced with 0.5 mM IPTG for 2 hours at 37 °C. MCAC

fractions containing Jnk1 were pooled and desalted into Buffer C (20 mM HEPES 7.0, 1 mM DTT, 1 mM EDTA, 10% glycerol) plus 100 mM NaCl by passage over a PD-10 column (Amersham Biosciences). PD-10 eluate was diluted 1:5 with buffer C immediately before loading on a 1 ml HiTrap SP column (Amersham Biosciences) equilibrated in Buffer C. The column was washed with 5 column volumes Buffer C, 5 column volumes Buffer C plus 50 mM NaCl, eluted with a 7 column volume gradient to Buffer C plus 300 mM NaCl, and finished with 5 column volumes of Buffer C plus 300 mM NaCl. The SP column was run at 0.5 ml/min collecting 0.5 ml fractions.

Ion-exchange chromatography conditions for p38gamma, h-Ras, and RhoA were identical to ERK, except KCl was substituted for NaCl in Buffer D additions. For purification of p38gamma, cultures were grown to an OD600 of 0.8 and induced with 0.5 mM IPTG for 3 hours at 30 °C. For purification of RhoA and h-Ras, cells were grown at 37 °C to an OD600 of 0.8 at 37 °C, and induced with 0.5 mM IPTG for 3 hours. MCAC fractions were diluted two fold with ddH<sub>2</sub>O, and dialyzed into Buffer D (20 mM HEPES 7.4, 1 mM DTT, 1 mM EDTA, 10% glycerol) plus 50 mM NaCl. Dialysate was applied to a 1 ml HiTrap Q column (Amersham Biosciences), washed with 5 column volumes Buffer D plus 50 mM NaCl, eluted with 8 column volumes to Buffer D plus 500 mM NaCl, and flushed with 3 column volumes Buffer D plus 500 mM NaCl. Column flow rate and fractionation parameters were identical to Jnk1 purification.

For purification of p38delta, cells were grown at 37 °C to OD600 1.0 and induced with 0.5 mM IPTG for 2 hours. MCAC fractions were dialysed against Buffer B plus 100 mM KCl, then diluted threefold with Buffer B immediately before loading on a 1 ml HiTrap SP column. The column was washed with 5 column volumes of Buffer B plus 50 mM KCl, and eluted with 10 column volumes to Buffer B plus 500 mM KCl, at a flow rate of 0.5 ml/min taking 0.5 ml fractions.

For purification of Mek1, a fresh transformation was grown for 12 hours at 30 °C, split 1:100 into fresh media, and grown to OD600 0.7 at 37 °C, then induced with 1 mM IPTG for 3 hours. MCAC fractions were desalted on a PD-10 column into Buffer E (40 mM HEPES, pH 7.4, 1 mM EDTA, 1 mM DTT, 10% glycerol) and applied to a 1 ml HiTrap Q column. The column was washed with 10 CV Buffer E and Mek1 was eluted in a single step of Buffer E plus 85 mM NaCl. Q column run parameters were as for Jnk1; fractions containing Mek1 were collected and dialyzed into Buffer F (150 mM NaCl, 10 mM HEPES, pH 7.5, 1 mM DTT, 1 mM EDTA) and applied to a Superdex 75 column (Amersham Biosciences). The Superdex 75 column was run at 1 ml/min and 1 ml fractions were

collected. Fractions containing Mek1 were pooled and concentrated tenfold on a Centricon YM-10 (Amicon), and glycerol was added to 10% v/v before proteins were frozen in liquid nitrogen.

- For purification of Mek1-DD (Huand and Erikson, 1994), cells were grown to  
5 OD600 of 0.7 at 37C and induced with 0.5 mM IPTG for 3 hours. Purification conditions were identical to RhoA and H-ras.

**Example 10. Activation of ERK and generation of ppERK.**

- Recombinant ERK and Mek1-DD were mixed in a 20:1 molar ratio in 1X Kinase  
10 Activation Buffer (100 mM NaCl, 10 mM MgCl<sub>2</sub>, 10 mM HEPES, pH 8.0, 10% glycerol). ATP was added to 1 mM and the mixture incubated for 30 minutes at 23 °C. Excess ATP was removed from the reaction by desalting on a PD-10 column into Buffer G (20 mM HEPES, pH 8.0, 2 mM DTT, 10% glycerol) plus 25 mM NaCl. The eluate was loaded on a  
15 1 ml HiTrap Q column and washed with 20 column volumes of Buffer G plus 100 mM NaCl. ppERK was eluted with a 60 column volume gradient to Buffer G plus 300 mM NaCl at 0.5 ml/min, collecting 0.5 ml fractions. Activated ERK eluted at 176 mM NaCl, while inactivated ERK eluted at 228 mM NaCl. Fractions containing ppERK were diluted six fold with buffer G and reappplied to a 1 ml HiTrapQ column. The column was washed with 50 mM NaCl, and the ppERK was eluted with a single step of Buffer G plus 500 mM NaCl.

20

**Example 11. *In vitro* selection of cis-hammerhead derived ERK and ppERK NASMs**

- Library design.** Figure 1B, illustrates an RNA ribozyme library derived from a hammerhead sequence pool consisting of up to 10<sup>17</sup> variants of randomized sequences appended to the hammerhead ribozyme motif (Figure 19). The ribozyme library is prepared  
25 on DNA synthesizer. Random nucleotides are incorporated during the synthesis to generate pools of roughly 10<sup>17</sup> molecules. Figure 19 illustrates linker scanning library designed to identify cis-hammerhead NASMs that are modulated by the protein kinase ERK. The linker library was generated by appending an ERK target modulation domain to the randomized linker domain to create a library of potential ERK-modulated cis-hammerhead NASMs.  
30 The linker library of ERK-modulated cis-hammerhead NASMs consists of up to 65,000 variants. Most molecules in the randomized NASM pools are non-functional NASMs.

**Selection of NASMs.** Sorting among the billions of NASMs to find the desired molecules starts from the complex sequence pool, whereby desired target-modulated NASMs are isolated through an iterative *in vitro* selection process: in addition to the target-

activated ribozymes that one desires, the starting pool is usually dominated by either constitutively active or completely inactive ribozymes. The selection process removes both types of contaminants. In a following amplification stage, thousands of copies of the surviving sequences are generated to enable the next round of selection. During  
5 amplification, random mutations can be introduced into the copied molecules — this 'genetic noise' allows functional nucleic acid sensor molecules to continuously evolve and become even better adapted as target-activated enzymes. The entire experiment reduces the pool complexity from  $10^{17}$  molecules down to around 100 NASMs that require detailed characterization.

10 ***In vitro* Selection Method.** Negative selections were set up in which 100  $\mu$ l reactions containing the library (c.f. Figure 19, where the linker library of ERK NASMs consists of up to 65,000 variants, alternatively the library consists of up to  $10^{17}$  variants of randomized sequences appended to the hammerhead ribozyme motif) of target modulated ribozymes containing randomized target modulation domains and randomized linker regions  
15 (1  $\mu$ M), were incubated at 72 °C, 5 seconds, followed by 25°C for 1hr in a reaction buffer containing KCl (0-150 mM), HEPES pH 7.5 (20 mM),  $MgCl_2$  (20 mM), EDTA (0.5 mM). Cleaved RNA molecules (radiolabeled internally) were then removed from the mixture by electrophoresis through a 7M urea, 8% PAGE gel, followed by cutting the uncleaved band from the gel. Detection of cleaved and uncleaved RNAs were carried out using a phosphor-  
20 imager system. The uncleaved hammerhead-derived RNAs were then subjected to a round of positive selection for ERK, or ppERK, target modulation, by incubating the pool of RNA in selection to which the protein target (50nM to 1  $\mu$ M ERK, or 50nM to 1  $\mu$ M ppERK) were added, and the mixture was further incubated at 37 °C for 30 minutes. Cleaved RNA molecules (radiolabeled internally) were then isolated and removed from the mixture by  
25 electrophoresis through a 7M urea, 8% PAGE gel, followed by cutting the cleaved band from the gel. incubated with reverse transcription mix at 65 °C for 1 hour using a thermostable reverse transcriptase. The pool was subsequently separated from the reverse transcription mixture by filtration and was then amplified using PCR first with the substrate 3-specific 5'-PCR primer and the library-specific 3'-PCR primer, and secondly with the  
30 regeneration 5'-PCR primer and the library-specific 3'-PCR primer in order to add the T7 promoter. Transcriptions were performed with these PCR products directly and after denaturing PAGE purification the entire process was repeated for several rounds with assays run after every round.

#### Cleavage Assays

Cleavage assays were performed using radiolabeled RNA and analytical denaturing polyacrylamide gel electrophoresis (PAGE) (gel-based assays). Assays were performed upon both the library and clonal sequences. In a representative gel-based assay transcription was performed in the presence of alpha-<sup>32</sup>P-labelled UTP, and the resultant transcripts were gel-purified using denaturing PAGE. Assay mixtures were then made in which 10 µl reactions containing RNA (1 µM), and protein target (1 µM) were incubated at 37 °C for between 15min and 16h in a reaction buffer containing buffer containing KCl (0-150 mM), HEPES pH 7.5 (20 mM), MgCl<sub>2</sub> (20 mM), EDTA (0.5 mM). The resultant samples are quenched by the addition of EDTA and the relative extents of cleavage measured by comparison of the intensity of the corresponding bands on a denaturing PAGE observed by reading a phosphorimager plate that had been exposed to the gel.

See general Example 1A for a discussion of the cloning, sequencing and characterization of individual NASMs.

**Example 12. Nucleic acid sensor molecules modulated by ERK and phosphorylated ERK generated by engineering target modulation domains into hammerhead catalytic domains.**

ERK and ppERK modulated nucleic acid sensor molecules were generated by a strategy combining both engineering and *in vitro* selection, Figure 19. Target modulation domains selected for binding to ERK and to the phosphorylated form of ERK (ppERK) are the starting point of the engineering efforts (Seiwert *et al.* 2000). These target modulation domains when isolated as discrete aptamers specifically recognize ERK but do not detectably interact with other mitogen-activated protein kinases such as Jun N-terminal kinase or p38 (as monitored by the ability to inhibit kinase activity). An aptamer (target modulation domain) selected for ppERK binding efficiently discriminates between phosphorylated and non-phosphorylated forms of the protein, binding ppERK with a  $K_D$  of 4.7 nM and ERK with a  $K_D$  of 50 nM (Seiwert *et al.* 2000). Simply engineering an ERK aptamer sequence onto a stably duplexed linker domain appended to the hammerhead ribozyme motif as shown in Figure 21, results in non-functional NASMs, *e.g.*, certain ribozyme NASMs, derived through direct engineering, when assayed as described above, exhibit little or no cleavage activity in the absence of ERK and no detectable ERK-modulation of cleavage activity in the presence of ERK. Inactive NASMs remained unresponsive to added ERK protein in concentrations ranges tested from 50 nM to 5 µM. Hence, ERK concentrations greater than 1000 fold over the aptamer  $K_D$  values does not

result in target modulation of the NASM, suggesting that functional NASM activity will require more than a functional target modulation domain appended to a ribozyme motif.

*In vitro* selection of cis-hammerhead NASMs based on the 65,000 member ERK target modulated ribozyme library shown in Figure 19, were performed as described in.

- 5 Example 1A. *In vitro* selection of hammerhead derived ERK and ppERK NASMs, and the subsequently identified and characterized a series of ERK-modulated NASMs are shown in Figure 19, and in Table 11. The detailed methods for the analysis ERK-modulated NASM clones is given in Example 1A Cloning, sequencing and characterization of individual NASMs.

10

**Example 13. *In vitro* selection of L1-ligase derived ERK and ppERK NASMs**

- Starting from a random linker domain library of L1-ligase derived NASMs as shown in Figure 31, we carried out *in vitro* selection to identify ERK-modulated and or ppERK-modulated L1-ligase NASMs. Negative selections were set up in which 100  $\mu$ l reactions
- 15 containing the library of ribozymes (1  $\mu$ M), effector oligonucleotide (1.5  $\mu$ M) and substrate 3 (5  $\mu$ M) were incubated at 25 °C for between 16 and 160 h in a reaction buffer containing KCl (150 mM), HEPES (20 mM), MgCl<sub>2</sub> (10 mM), EDTA (1 mM), DTT (1 mM), tRNA (0.1 mg/ml) and glycerol (10% w/v). Ligated RNA molecules were then removed from the mixture by incubation with immobilized neutravidin. The flow through was collected by
- 20 filtration and to this was added more substrate 3 (2  $\mu$ M), protein target (1  $\mu$ M ERK or ppERK) and RNase inhibitor were added and the mixture was further incubated at 25 °C for between 15 minutes and 1 hour. Ligated RNA molecules were then captured by incubation with immobilized neutravidin and, after washing, the matrix was incubated with reverse transcription mix at 65 °C for 1 hour using a thermostable reverse transcriptase. The matrix
- 25 was subsequently separated from the reverse transcription mixture by filtration and was then amplified using PCR first with the substrate 3-specific 5'-PCR primer and the library-specific 3'-PCR primer, and secondly with the regeneration 5'-PCR primer and the library-specific 3'-PCR primer in order to add the T7 promoter. Transcriptions were performed with these PCR products directly and after denaturing PAGE purification the entire process
- 30 was repeated for several rounds with assays run after every round.

**Ligation Assays**

Ligation assays were performed using radiolabeled RNA and analytical denaturing polyacrylamide gel electrophoresis (PAGE) (gel-based assays), or by PCR utilizing the substrate 3-specific 5'-PCR primer (PCR-based assay). Assays were performed upon both

the library and clonal sequences. In a representative gel-based assay transcription was performed in the presence of alpha-<sup>32</sup>P-labelled UTP, and the resultant transcripts were gel-purified using denaturing PAGE. Assay mixtures were then made in which 10 µl reactions containing RNA (1 µM), effector oligonucleotide (1.5 µM), substrate 3 (5 µM) and optionally RNase-inhibitor and protein target (1 µM) were incubated at 25 °C for between 15min and 16h in a reaction buffer containing protein target (1 µM ERK or ppERK), KCl (150 mM), HEPES (20 mM), MgCl<sub>2</sub> (10 mM), EDTA (1 mM), DTT (1 mM), tRNA (0.1 mg/ml) and glycerol (10% w/v). The resultant samples were quenched by the addition of EDTA and the relative extents of ligation were measured by comparison of the intensity of the corresponding bands on a denaturing PAGE observed by reading a phosphorimager plate that had been exposed to the gel. PCR assays were performed using the same reaction mixture except that the relative extents of ligation in different samples were compared by observing the relative rates of appearance of PCR products using the substrate 3-specific 5'-PCR primer and the library-specific 3'-PCR primers. The detailed methods for the analysis of ERK-modulated NASM clones is given in Example 1A, Cloning, sequencing and characterization of individual NASMs.

**Example 14. Rational design and Engineering of ERK dependent ligases.**

ERK-dependent ligase ribozymes were built on the catalytic core of the L1 ligase ribozyme of Robertson and Ellington (2000; NAR 28, 1751-1759) in which the non-conserved, stem C element was replaced with the ERK-interacting domain and joined to the catalytic core by a 2-4 base pair helical element, or "communication module". Ten different ribozymes were designed, identical in sequence except for the bases in the linker domain (Figure 32). (SEQ ID NOs:109-116)

Templates for transcription by T7 RNA polymerase were prepared for each ribozyme by PCR amplification using a set of three overlapping primer oligonucleotides. The forward, or 5' primer  
TAATACGACTCACTATAGGACTTCGGCGAAAGCCGTTCTGACC (SEQ ID NO:315),  
included the T7 RNA polymerase promoter and sequence corresponding to the 5'-proximal region of the L1 ligase catalytic core. The other two primers corresponded to sequences spanning the 3'-proximal portion of the L1 ligase core  
ATTCGAGATGTCCTTGGACCAAAGCCGCACCTAACCTCCTGTCTAAG (SEQ ID NO:316) and the ERK interacting domain (including sequences), respectively. RNA was synthesized by *in vitro* transcription (Milligan & Uhlenbeck (1989) Methods Enzymol 180,



- 51-62) using the T7-MEGAscript™ transcription kit from Ambion, and purified by gel-filtration (to remove transcription buffer components) and/or polyacrylamide gel electrophoresis (PAGE). The purified RNAs were then quantified by their absorbance at 260 nm, and stored in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) at -20 °C.
- 5 Radiolabeled RNAs were prepared exactly as described above, except that  $\alpha$ -<sup>32</sup>P-UTP was included in the *in vitro* transcription reaction.

**Example 15. Rational design and Engineering of ppERK dependent ligases.**

- ppERK-dependent ligase ribozymes were built on the catalytic core of the L1 ligase ribozyme of Robertson and Ellington (2000; NAR 28, 1751-1759) in which the non-conserved, stem C element was replaced with the ERK-interacting domain and joined to the catalytic core by a 2-4 base pair helical element, or "communication module". Fourteen different ribozymes were designed, identical in sequence except for the bases in the linker domain (Figure 41) (SEQ ID NO:352).

- 15 Templates for transcription by T7 RNA polymerase were prepared for each ribozyme by PCR amplification using a set of three overlapping primer oligonucleotides. The forward, or 5' primer  
TAATACGACTCACTATAGGACTTCGGCGAAAGCCGTTTCGACC (SEQ ID NO:315), included the T7 RNA polymerase promoter and sequence corresponding to the 5'-proximal  
20 region of the L1 ligase catalytic core. The other two primers corresponded to sequences spanning the 3'-proximal portion of the L1 ligase core  
ATTCGAGATGTCTTGGACCAAAGCCGCACCTAACCTCCTGTCTAAG (SEQ ID NO:316) and the ppERK target modulation domain (including sequences), respectively. RNA was synthesized by *in vitro* transcription (Milligan & Uhlenbeck (1989) Methods  
25 Enzymol 180, 51-62) using the T7-MEGAscript™ transcription kit from Ambion, and purified by gel-filtration (to remove transcription buffer components) and/or polyacrylamide gel electrophoresis (PAGE). The purified RNAs were then quantified by their absorbance at 260 nm, and stored in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) at -20 °C. Radiolabeled RNAs were prepared exactly as described above, except that  $\alpha$ -<sup>32</sup>P-UTP was  
30 included in the *in vitro* transcription reaction.

**Example 16. Linker-region selection of effector-dependent ligases**

**Oligonucleotides**

Synthetic DNA and the DNA-RNA chimeric substrates were synthesized using standard solid-phase methodology and purified by denaturing (8M urea) polyacrylamide gel electrophoresis (PAGE). Random-sequence regions of synthetic DNA were generated using a single mixed "N" bottle on the DNA synthesizer that contains all four nucleotide phosphoramidites mixed in proportions. Variable-length regions of synthetic DNA were generated by interrupting the synthesis and removing part of the reaction mixture, re-starting the synthesis and then replacing the removed material after a subsequent interruption. Templates for transcription were generated by PCR, and the transcripts thus generated were also purified by denaturing (8M urea) polyacrylamide gel electrophoresis (PAGE). Gel-purification is followed by the localization of nucleic acids within the gel by UV-shadowing, nucleic-acid containing gel pieces are excised and the purified nucleic acids are recovered by electroelution. The following synthetic oligonucleotides were utilized:

**Oligonucleotides common to all selections:**

Substrate 3 (DNA, RNA underlined)

15 Biotin-TEG-CATGCGACCTTACGATCAGATGACCTUGCACU (SEQ ID NO:321)

Substrate 3-specific 5'-PCR primer

20 CATGCGACCTTACGATCAGAT (SEQ ID NO:322)

Regeneration 5'-PCR primer

TCTAATACGACTCACTATAGGACTTCGGCGAAAGC (SEQ ID NO:323)

**ERK selection-specific oligonucleotides:**

25 Library (DNA) (random regions 3-5 nucleotides in length)

GGACTTCGGCGAAAGCCGTTTCGACNNN(N)(N)AAGGAGGATTTCCGAAAGCGG  
CTACGGTCCGCCNNN(N)(N)CTTAGACAGGAGGTTAGGTGCGTAGGTAACCGAT  
AGTTCCG (SEQ ID NO:324)

30 Effector (2'-O-methyl DNA), 3'-PCR primer (DNA) and RT primer (DNA)  
CGGAACTATCGGTTACCTAC (SEQ ID NO:325)

**ppERK selection-specific oligonucleotides:**

Library

GGACTTCGGCGAAAGCCGTTTCGACNNNN(N)(N)CAGACGCTAGCGAATTGGTT  
 CCTCGAAAGGGGAAAGCGTTATTAAGAAACCAAAATGNNNN(N)(N)CTTAGACA  
 GGAGGTTAGGTGCGTCAATGCTGCAAGTTACTG (SEQ ID NO:326)

- 5        Effector (2'-O-methyl DNA), 3'-PCR primer (DNA) and RT primer (DNA)  
           CAGTAACTTGCAGCATTGAC (SEQ ID NO:327)

**bFGF selection-specific oligonucleotides:**

- 10       Library (random regions 5-7 nucleotides in length)  
           GGACTTCGGCGAAAGCCGTTTCGACNNNNN(N)(N)GCAACGCTACAGACA  
           AGTGCANNNNN(N)(N)CTTAGACAGGAGGTTAGGTGCCCCGAGTTGTTGCAACGA  
           GAC (SEQ ID NO:328)

- 15       Effector (2'-O-methyl DNA), 3'-PCR primer (DNA) and RT primer (DNA)  
           GTCTCGTTCGAACAACCTCGG (SEQ ID NO:329)

**Thrombin selection-specific oligonucleotides:**

**Library**

- 20       GGACTTCGGCGAAAGCCGTTTCGACNNNN(N)(N)ATCGAAGTTAGTAGGNNN(N)  
           (N)CTTAGACAGGAGGTTAGGTGCGTCAATCGATTGCAGATCCG (SEQ ID  
           NO:330)

- 25       Effector (2'-O-methyl DNA), 3'-PCR primer (DNA) and RT primer (DNA)  
           CGGATCTGCAATCGATTGAC (SEQ ID NO:331)

**Example 17. Nucleic acid sensor molecules modulated by native ERK enzyme.**

- Hammerhead-derived nucleic acid sensor molecules (88 nucleotides in length) were  
 selected from populations of nucleic acid molecules with randomized linker domain in stem  
 30    II, as shown in Figure 19. The Table in Figure 19 depicts illustrative linker-domain  
 sequences of several of the ERK-modulated nucleic acid sensor molecules isolated from *in*  
*vitro* selection linker randomized clones (described in detail in Example 11). Each NASM  
 displays a varying degree of modulation driven by addition of equal amounts (1  $\mu$ M) native  
 ERK. Individual ERK-modulated cis-hammerhead nucleic acid sensor molecules are shown

in Figure 19. Clones 1-14, 1-13, 1-2, 1-6, 2-7, 2-2, 2-3, 2-13, 2-14, and 2-20 were tested in target modulation assays as described previously. The NASM's relative dependence on ERK is denoted in the Table in Figure 19 by the extent of activity of the ribozyme in the presence of ERK protein.

5       The time course of signal generation in the presence of nonphosphorylated ERK, phosphorylated ERK, and in the absence of protein is determined by measuring signal released over time by a radiolabeled nucleic acid sensor molecule. Significant amounts signal, corresponding to cleavage of the nucleic acid sensor molecule is observed over time only with the nonphosphorylated ERK. Clones 1-2, 1-13 and 1-14 all display sensitivity to  
10   ERK concentration, as shown in Figures 20 A, B and C. However, of these three clones, clone 1-14 displays the greatest enhancement in activity upon addition of ERK. Clone 1-14 is able to differentiate between varying concentrations of ERK, as indicated by the dose-dependent change in the activity of clone-14 upon the addition of 5, 10, 20, 50 and 100 nM ERK (Figure 21).

15

**Example 18. ERK-modulated NASM-based competitive inhibition assays, & target-protein specific profiling biosensors.**

The specificity of the interaction between clone 1-14 and ERK was assessed by measuring the activity of clone 1-14 in the presence of 50 nM ERK and increasing  
20   concentrations of known protein kinase inhibitors. The kinase inhibitors staurosporine and 5-iodo-tubercidin are compounds that are known bind to in an ATP-substrate competitive manner to ERK and to modulate ERK kinase catalytic activity, decreasing its ability to phosphorylate other proteins. We tested whether kinase inhibitors would compete directly for the ATP-binding site in ERK, and thereby block NASM activation by ERK. As shown  
25   in Figure 22A, increasing concentrations of the compounds staurosporin and or ITU disrupt the interaction between clone 1-14 and ERK protein, suppressing the activation of the nucleic acid sensor molecule. We have used NASM-based assay to confirm the known kinase-inhibitory IC<sub>50</sub> values of staurosporin and ITU (5 and 1  $\mu$ M, respectively). This result indicates that clone 1-14 recognizes free unliganded ERK and is not modulated by the  
30   ERK-staurosporine or ERK-ITU complex. Since, staurosporin or ITU (ATP analogs) are general kinase inhibitors, and bind to the active/ATP binding sites of kinases, these data suggest that the NASMs either bind directly to the kinase active site, or are exquisitely sensitive to the conformational state of the enzyme. Hence, when saturating levels of the kinase inhibitors are present (10  $\mu$ M) in the NASM-assay, along with  $\mu$ M levels of ERK, the

nucleic acid sensor molecule does not signal the presence of ERK protein. Thus, the ERK-modulated NASMs are used as general HTS screening reagents for the discovery of small molecule drug leads.

Similar results to those described above for the hammerhead-derived NASMs of Figure 19 have been obtained with ERK-modulated L1-ligase NASMs described in detail in Example 13. Figure 58, shows additional staurosporine competitive NASM assay results using an L1-ligase-derived NASM which is also modulated by ERK protein. Additional modifications of the 3-piece L1-ligase NASM (c.f., Figure 36, constructs 27 and constructs 28) render these biosensors suitable for cellular assays as intracellular biosensors (c.f., Figure 37 and the detailed description of the 3-piece, and 1-piece ligases described in detail in Example 22). Thus, ERK-modulated nucleic acid sensor molecules are used for cell-based drug discovery and drug candidate screening.

The data in Figures 22B and Figures 57 A and 57B indicates that ERK NASMs (clone 1-14 and constructs 27 and 28) are modulated specifically by ERK only in its fully active, and not by ppERK, or by related homologues and MAP Kinase pathway molecules, For example clone 1-14 recognizes only ERK, and not related kinase MEK. Hence, NASMs and more specifically, the ERK and ppERK NASMs, are shown here to be useful specific protein profiling biosensors.

A compilation of novel hammerhead ERK-modulated nucleic acid sensor molecule sequences are disclosed in Table 11.

**Table 11. Cis-hammerhead nucleic acid sensor molecules**

Sequences of 157 ERK activated cis-hammerhead clones.

Sequence	SEQ ID
NO:	
GGGCGACCCUGAUGAGUCGGGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAAUACGAAACGGUGAAAGCCGUAGGUUGCC	140
GGGCGACCCUGAUGAGGGAGGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGCGCGAAACGGUGAAAGCCGUAGGUUGCC	141
GGGCGACCCUGAUGAGGGUUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGCGCAAACAGNGAAANNNGNANGUGGNC	142
GGGCGACCCUGAUGAGUACGGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAGAACGAAACGGUGAAAGCCGUAGGUUGCC	143
GGGCGACCCUGAUGAGNCAUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAAACNCGAAACGGUGAAAGCCGUAGGUUGCC	144
GGGCGACCCUGAUGAGUUGCGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAAACGAAACGGUGAAAGCCGUAGGUUGCC	145
GGGCGACCCUGAUGAGUUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAUACGAAACGGUGAAAGCCGUAGGUUGCC	146
GGGCGACCCUGAUGAGUAAUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAGGCCAAACGGNGAAAGCCCGUANGNUGCC	147
GGGCGACCCUGAUGAGCUCUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGGCCGAAACGGUGAAAGCCGUAGGUUGCC	148
GGGCGACCCUGAUGAGGCAUGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUGCGCGAAACGGUGAAAGCCGUAGGUUGCC	149

GGGCGACCCUGAUGAGCCCCGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUUGAGCGAAACGGUGAAAGCCGUAGGUUGCC  
150  
GGGCGACCCUGAUGAGUACGGCUNAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUNANACGAAACNGNGAAAGCCGUAGGUUGCC  
151  
5 GGGCGACCCUGAUGAGCAUGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUACUACAGAAACGGUGAAAGCCGUAGGUUGCC  
152  
GGGCGACCCUGAUGAGUGACGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUUGCGCGAAACGGUGAAAGCCGUAGGUUGCC  
153  
10 GGGCGACCCUGAUGAGUCUGGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUUCGCGAAACGGUGAAAGCCGUAGGUUGCC  
154  
GGGCGACCCUGAUGAGNANCGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
155  
GGGNGACCCUGANGANNACGCUAAGGAGGAUUCGAAAGCGGCUACUAGUCCNCAUUGNACCGAAACGGCCUAAAGCCGGAGGUUGCC  
156  
15 GGGCGACCCUGAUGAGAUACGCUAAGGAGGAUUCGAAAGCGGCUACUGGUCGCCAGUUGCGAAACGGUGAAAGCCGUAGGUUGCC  
157  
GGGCGACCCUGAUGAGUGGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUCACGAAACGGUGAAAGCCGUAGGUUGCC  
158  
20 GGGCGACCCUGAUGAGCUGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUCGACGAAACGGUGAAAGCCGUAGGUUGCC  
159  
GGGCGACCCUGAUGAGGCGUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
160  
GGGCGACCCUGAUGAGAACUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGUCAGNUACUCGAAACGGUGAAAGCCGUAGGUUGCC  
161  
25 GGGCGACCCUGAUGAGNGCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGCGCGAAACGGUGAAAGCCGUAGGUUGCC  
162  
GGGCGACCCUGAUGAGNANGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGCGAAACGGUGAAAGCCGUAGGUUGCC  
163  
30 GGGCGACCCUGAUGAGNUNUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGCGAAACGGUGAAAGCCGUAGGUUGCC  
164  
GGGCGACCCUGAUGAGGAGUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
165  
GGGCGACCCUGAUGAGAGCCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUCNACGAAACGGUGAAAGCCGUAGGUUGCC  
166  
35 GGGCGACCCUGAUGAGGCGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
167  
GGGCGACCCUGAUGAGUAAACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
168  
40 GGGCGACCCUGAUGAGNUACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
169  
GGGCGACCCUGAUGAGUUGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
170  
GGGCGACCCUGAUGAGUUNUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUNUACGAAACGGUGAAAGCCGUAGGUUGCC  
171  
45 GGGCGACCCUGAUGAGACUG - CUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
172  
GGGCGACCCUGAUGAGGUGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
173  
50 GGGCGACCCUGAUGAGUUAUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
174  
GGGCGACCCUGAUGAGUGGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
175  
GGGCGACCCUGAUGAGUCUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGGCGAAACGGUGAAAGCCGUAGGUUGCC  
176  
55 GGGCGACCCUGAUGAGCCUNGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
177  
GGGCGACCCUGAUGAGUCAUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGGCGAAACGGUGAAAGCCGUAGGUUGCC  
178  
60 GGGCGACCCUGAUGAGAACCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGGCGAAACGGUGAAAGCCGUAGGUUGCC  
179  
GGGCGACCCUGAUGAGGGNGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
180  
GGGCGACCCUGAUGAGUACGCUAANGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
181  
65 GGGCGACCCUGAGAGUACNNGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
182  
GGGCGACCCUGAUGAGUCUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
183  
GGGCGACCCUGAGAGGAUA - GCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
184  
70 GGGCGACCCUGAUGAGUAGUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
185  
GGGCGACCCUGAUGAGUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUAGGUUGCC  
186  
75 GGGCGACCCUGNNGAGAACUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUGACGAAACGGUGAAAGCCGUANGUUGCC  
187

GGGCGACCCUGANGAGNUCNGCUNAGGAGGAUUCGAAAGCGGCUACGGAGCCGUCAGUAUUGCGAAACGGCGAAAGCCGUAGGUUGNC  
188  
GGGCGACCCUGAUGAGUAACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUAUUGCGAAACGGUGAAAGCCGUAGGUUGCC  
189  
5 GGGCGACCCUGAUGAGCUUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUAAACGAAACGGUGAAAGCCGUAGGUUGCC  
190  
GGGCGACCCUGAUGAGNCCAGCUAAGGAGGAUUCGAAAGCGGCUACGGNCCGCCAGUUAACCGAAACGGNGAAAGCCGUANGUUGCC  
191  
10 GGGCGACCCUGAUGAGNCCAGCUAAGGAGGAUUCGAAAGCGGCUACGGNCCGCCAGUUAACCGAAACGGNGAAAGCCGUAGGUUGCC  
192  
GGGCGACCCUGAUGAGCUUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGUCGAAACGGUGAAAGCCGUAGGUUGCC  
193  
GGGCGACCCUGAUGAGCUNNGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUNUNCGAAACGGUGAAAGCCGUAGGUUGCCU  
194  
15 GGGCGACCCUGAUGAGACCUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
195  
GGGCGACCCUGAUGAGCAUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUCGCGGAAACGGUGAAAGCCGUAGGUUGCC  
196  
20 GGGCGACCCUGAUGAGAUUCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUNACGCGAAACGGUGAAAGCCGUAGGUUGCC  
197  
GGGCGACCCUGA - GAGUUUGCUAAGGAGGAUUCGAA - GCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
198  
GGGCGACCCUGN - GAGUUUAGCUAAGGAGGAUUCGAA - GCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
199  
25 GGGCGACCCUGNNGAGGCGUGCUAAGGAGGAUUCGAA - GCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
200  
GGGCGACCCUGAUGAUAUUGCUAAGGAGGAUUCGAA - GCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
201  
GGGCGACCCUGAUGAGCAGAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
202  
30 GGGCGACCCUGAUGAGCAUAGCUAAGGAGGAUUCGAA - AGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
203  
-----UGNAGAGCGUUGCUAAGGAGGAUUCGAAAGC - GCUACGUCGCGCCAGUAACACGAAACGGUGAAAGCCGUAGGUUGCCU  
204  
35 GGGCGACCCUGAUGAGCAACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
205  
GGGCGACCCUGAUGAGGAGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
206  
40 GGGCGACCNUGAUGAGCUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
207  
GGGCGACCCUGAUGAGACCGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
208  
GGGCGACCNUGAUGAGUAUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
209  
45 GGGCGACCNUGAUGAGUAUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
210  
GGGCGACCNUGAUGAGUNGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
211  
50 GGGCGACCCUGAUGAGCAACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
212  
GGGCGACCCUGAUGAGUCGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
213  
GGGCGACCCUGAUGAGUCACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
214  
55 GGGCNACCUCUGAGUAGCAGGCUAAGNAGNAUUCGNAANCNGNUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
215  
GGGCGACCCUGAUGAGCUGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
216  
GGGCGACCCUGAUGAGCANNGCUAAGNAGUAUUCGNAANCNGNUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
217  
60 GGGCGACCCUGAUGAGCAUAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
218  
GGGCGACCNUGAUGAGCUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
219  
65 GGGCGACCCUGAUGAGACACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
220  
GGGCGACCNUGAUGAGCNUGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
221  
70 GGGCGACCNUGAUGAGUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
222  
GGGCGACCNUGAUGAGGUGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
223  
GGGNGACCCUGAUGAGUAGGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
224  
75 GGGNAGACUUGAUGAGUAGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACCGAAACGGUGAAAGCCGUAGGUUGCC  
225

GGGCGACCCUGAUGAGCUCUGCUAAGGAGGATUUCGAAAGCGGCUACGGUCCGCCAGUUGGUCGAAACGGUGAAAGCCGUAGGUUGCC  
226  
GGGCGACCCUGAUGAGGGCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
227  
5 GGGNGACCCUGAUGAGCCAUGCUAAGGAGNAUUCUNAAAGCGGC-ACGGUCCGCCAGUUCGCGAAACGGUGAAAGCCGUAGGUUGCC  
228  
GGGCGACCCUGAUGAGUACGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUCACGAAACGGUGAAAGCCGUAGGUUGCC  
229  
10 GGGCGACCCUGAUGAGACTUGGUAAGGAGGAUUCGAAANCGNNUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
230  
GGGCGACCCUGAUGAGNUUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAUCGAAACGGUGAAAGCCGUAGGUUGCC  
231  
GGGCGACCCUGAUGAGCUCGCUAAGGAGGAUUCGAAAGCGGNNACGGUCCGCCAGUUCACGAAACGGUGAAAGCCGUAGGUUGCC  
232  
15 GGGCGACCCUGAUGAGNUNCNUAAGGAGNAUUCGAAAGCGGANACGGUCCGCCAGUNANCCGNAACGGUGAAAGCCGUAGGUUGCC  
233  
GGGCGACCCUGAUGAGGAUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUACAGCGAAACGGUGAAAGCCGUAGGUUGCC  
234  
20 GGGCGACCCUGAUGAGUCCUGCUAAGGAGGAUUCGAAAGCGGNNACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
235  
GGGCGACCCUGAUGAGUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
236  
GGGCGACCCUGAUGAGNAUNGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAANCGAAACGGUGAAAGCCGUAGGUUGCC  
237  
25 GGGCGACCCUGAUGAGGAAUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
238  
GGGCGACCCUGAUGAGCAGUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUGGAGCGAAACGGUGAAAGCCGUAGGUUGCC  
239  
30 GGGCGACCCUGAUGAGNUAUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUNAGNCGAAACGGUGAAAGCCGUAGGUUGCC  
240  
GGGCGACCCUGAUGAGCACUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUGAGACGAAACGGUGAAAGCCGUAGGUUGCC  
241  
GGGCGACCCUGAUGAGCUCAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
242  
35 GGGCGACCCUGAUGAGGCUUGCUAAGGAGGAUUCGAAAGCGGCNACGGUCCGCCAGUCCGUGCGAAACGGUGAAAGCCGUAGGUUGCC  
243  
GGGCGACCCUGAUGAGCUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUCGCGAAACGGUGAAAGCCGUAGGUUGCC  
244  
40 GGGCGACCCUGAUGAGGCAAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
245  
GGGCGACCCUGAUGAGCGUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAUCGAAACGGUGAAAGCCGUAGGUUGCC  
246  
GGGCGACCCUGAUGAGCUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAUCGAAACGGUGAAAGCCGUAGGUUGCC  
247  
45 GGGCGACCCUGAUGAGUGGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAA  
248  
GGGCGACCCUGAUGAGACUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUGGUAACGAAACGGUGAAAGCCGUAGGUUGCC  
249  
50 GGGCGACCCUGAUGAGUAUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
250  
GGGCGACCCUGAUGAGCNUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
251  
GGGCGACCCUGAUGAGCUNNGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
252  
55 GGGCGACCCUGAUGAGUUCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
253  
GGGCGACCCUGAUGAGAGCUCUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
254  
60 GGGCGACCCUGAUGAGAAUCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAACGAAACGGUGAAAGCCGUAGGUUGCC  
255  
GGGCGACCCUGAUGAGCAGCUCUAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
256  
GGGCGACCCUGAUGAGCAUCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
257  
65 GGGCGACCCUGAUGAGGUCAGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
258  
GGGCGACCCUGAUGAGAGCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
259  
70 GGGCGACCCUGAUGAGGCUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
260  
GGGCGACCCUGAUGAGGCUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
261  
GGGCGACCCUGAUGAGGCUUGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
262  
75 GGGCGACCCUGAUGAGCUCGCUAAGGAGGAUUCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
263



GGGCGACCCUGAUGAGAAACUGCTAAGGAGGAUUUACCGAAAGCGGCUACGGUCCGCCAGUUGCGCGAAACGGUGAAAGCCGUAGGUUGCC  
264  
GGGCGACCCUGAUGAGCGACGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGGGCGAAACGGUGAAAGCCGUAGG  
265  
5 GGGCGACCCUGAUGAGUAAACGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUACAGACGAAACGGUGAAAGCCGUAGGU  
266  
GGGCGACCCUGAUGAGGACUGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAUCGAAACGGUGAAAGCCGUAGGU  
267  
10 GGGCGACCCUGAUGAGCCGGGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGAGCGAAACGGUGAAAGCCGUAGGUUGCC  
268  
GGGCGACCCUGAUGAGCANAGCTAAGGAGGAUUUCCGAAAG  
269  
GGGCGACCCUGAUGAGUUGCGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGU  
270  
15 GGGCGACCCUGAUGAGCAGCGCTAAGGAGGAUUUCCGAAAGCGGCUAGGUCCGCCAGUUCUACGAAACGGUGAAAGCCGUAGGUUG  
271  
GGGCGACCCUGAUGAGUAAAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGUGCGAAACGGUGAA  
272  
20 GGGCGACCCUGAUGAGCACCGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAACGCGAAACGGUGAAAGCCGUAGGUUGCC  
273  
GGGCGACCCUGAUGAGGCAUGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAG  
274  
GGGCGACCCUGAUGAGUUUAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAGACGAAACGGUGAAAG  
275  
25 GGGCGACCCUGAUGAGAAAGCGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAG  
276  
GGGUGACCTCUGAUGAGUUGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUAGAGCGAAACGGUGAAAGCCGUAGGUUGCC  
277  
30 GGGCGACCCUGAUGAGCANGCGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
278  
GGGCGACCCUGAUGAGCGCTAAGGAGGAUUUCCGAAAGCGGCTACACGGUCCGCCAGUUCACGANNCGGUNAAG  
279  
GGGNGACCCUGAUGAGUUNAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
280  
35 GGGCGACCCUGAUGAGGNCAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGGACGAAACGGUGAAAGCCGUAGGUUGCC  
281  
GGGCGACCCUGAUGAGCAUGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
282  
40 GGGCGACCCUGAUGANNCTGNTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUAAGCGAAACGGUGAAAGCCGUAGGUUGCC  
283  
GGGCGACCCUGAUGAGUAGGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUGGCGAA  
284  
GGGCGACCCUGAGAGCTACGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGACGAAACGGUGAAAGCCGUAGGUUGCC  
285  
45 Construct 1-14  
GGGCGACCCUGAUGAGGGGCTGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
286  
Construct 1-13  
GGGCGACCCUGAUGAGUCTCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
287  
50 Construct 1-2  
GGGCGACCCUGAUGAGCTCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
288  
Construct 1-6  
GGGCGACCCUGAUGAGUCTCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
289  
Construct 2-7  
GGGCGACCCUGAUGAGCTCTGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
290  
60 Construct 2-2  
GGGCGACCCUGAUGAGCTCTGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
291  
Construct 2-3  
GGGCGACCCUGAUGAGAACTGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
292  
65 Construct 2-13  
GGGCGACCCUGAUGAGUAAAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
293  
Construct 2-14  
GGGCGACCCUGAUGAGUUUUGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
294  
70 Construct 2-20  
GGGCGACCCUGAUGAGGCGAGCTAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUUCGAAACGGUGAAAGCCGUAGGUUGCC  
295  
75

These results demonstrate that the nucleic acid sensor molecule is specific for the non-phosphorylated form of the ERK.

**Example 19. Cis-hammerhead-derived nucleic acid sensor molecules activated by ppERK**

A stem-selection library of hammerhead-derived nucleic acid sensor molecules modulated by ppERK is shown in Figure 23A. ppERK-modulated nucleic acid sensor molecules were designed to by joining ppERK a target modulation domain (TMD) to a hammerhead core sequence via a linker region. Figure 23A shows the general structure of ppERK-modulated hammerhead ribozyme with the linker region depicted as a randomized sequence (N6). Individual ppERK dependent *cis* hammerhead RNA sequences are shown in Figure 23B.

Figure 24 is a table displaying different properties of the *cis* hammerhead ERK sensors shown in Figure 23B. Each construct displays variability in its sequence, its activity and its stability.

Figure 25 shows data for the ppERK-NASMs that have been tested in target modulation assays using *in situ* generated ppERK (using constitutively activated MEK activate to generate ppERK from native ERK). ppERK has been generated *in situ* by direct phosphorylation of ERK with constitutively active MEK mutant, MEK-EE in the presence of  $\text{MgCl}_2$  and ATP. Figure 25 also shows data for radiolabeled nucleic acid sensor molecules for ppERK-hammerhead NASM constructs 6, 10, and that were incubated at 37° C for 30 minutes in the absence of protein; in the presence of 2  $\mu\text{M}$  ERK, 0.14  $\mu\text{M}$  constitutively active MEK (MEK-EE); in the presence of 2  $\mu\text{M}$  ERK, 0.14  $\mu\text{M}$  constitutively active MEK-EE, 1 mM ATP; in the presence of 1 mM ATP only. All these reactions were performed in the following buffer of 20 mM HEPES (pH 7.5), 10 mM  $\text{MgCl}_2$ , and 0.1 mM EDTA. The mixtures of ERK, ERK + MEK-EE, and ERK+MEK+EE with ATP were incubated in the presence of 10mM  $\text{MgCl}_2$  prior to mixing with radiolabeled nucleic acid sensor molecule. After the incubation, the nucleic acid sensor molecule reaction mixtures were subjected to 5% acrylamide gel containing 8 M Urea as shown in panel B of Figure 25., in order to examine the NASM endonuclease activity. Three constructs displayed an increase in endonuclease activity in the presence of ERK + MEK + EE with ATP as shown in Figure 25. These data indicate general methods for a NASM-based ERK activation assay: as shown in Figure 25, NASMs are used to directly

detect MEK conversion of ERK to ppERK as detected by ppERK-modulation of nucleic acid sensor molecules.

**Example 20. Characterization of the ERK and ppERK modulated L1-Ligase derived NASMs**

A template for an ERK dependent ligase is shown in Figure 31. Figure 32 discloses the different clones of ERK dependent ligases based on this template. The ERK dependence of these clones is shown in Figure 33. Constructs 17 and 19 are particularly sensitive to 2 $\mu$ M ERK. Construct 19, also known as Clone E, was tested to see if it could be used to detect increasing concentrations of ERK. Construct 19 was found to have a distinct signal at 1  $\mu$ M, 100 nM, and 10nM ERK. Time dependency was also tested for Construct 19 as shown in Figure 34. Increasing signal from Construct 19 could be seen at 1, 2, 3, and 4 hours. Concentration dependence of Construct 19 (Clone E) is also shown in the graph in Figure 35. Construct 19 can differentiate between different concentrations of ERK. Two formats that can be used with nucleic acid sensor molecule ligases are the 3-piece ligase platform, and the 1-piece ligase platform, both shown in Figure 36. The different platforms allow the nucleic acid sensor molecules to be used in different applications. For example, the 3-piece could be used *in vitro*, while the 1-piece could be used *in vivo*, even intracellularly. The performance of these two formats is shown in Figure 37. There is also a 2-piece platform shown generally in Figure 38 and shown as ERK regulated 2-piece ligases as shown in Figure 39A. The 2-piece platform of Construct 19 is also ERK dependent as shown in Figure 39B.

Ligase nucleic acid sensor molecules can be used in the presence of cell extracts. Construct 19 shows ERK activation dependence in the presence of 50% HeLa cell extract as shown in Figure 40.

Ligase based nucleic acid sensor molecules were also found that were specifically activated by ppERK. Figure 41 shows a template for the ppERK sensitive ligase. Figure 42 show the specific sequences left generalized in Figure 41. ppERK sensitivity was shown in the clones described Figure 57 shows activation of 3-piece Construct 19 (DG.20.58E) (ERK2-dependent) (SEQ ID NO:111) and TK.16.118.M (ppERK2-dependent) (SEQ ID NO:123) ligases in the presence of purified proteins. Ligation assays (10 mM MgCl<sub>2</sub>, 100 mM NaCl, 30 mM Tris, pH 7.5) were performed at room temperature for 1 hour, using 1  $\mu$ M ligase RNA, 1  $\mu$ M effector oligonucleotide, 2  $\mu$ M substrate oligonucleotide and 1  $\mu$ M

protein. RNAs were incubated in the presence of: (A) purified ERK, in either its non-phosphorylated or diphosphorylated form; or (B) other members of the MAP kinase pathway (Ras, MEK1 and p38), and ricin. Ligase activity was measured in solution, using real-time RT-PCR detection.

5 It has been shown that Construct 19 still has activity in low magnesium ion. In one embodiment of the invention, these nucleic acid sensor molecules can be used in cell extracts and intracellularly, to detect ppERK. There is very low magnesium ion concentrations in these environments, so it is important that these nucleic acid sensor molecules maintain function even in low concentrations of magnesium ion.

10 Figure 42 shows the full sequences of the different clones of ppERK sensitive nucleic acid sensor molecules in the 3-piece format.

#### Example 21. Amplicon Detection

Amplicon dependent nucleic amplification is a high-throughput method for cell or  
15 tissue profiling of proteins and metabolites. The method can include using real-time PCR to follow the activity of nucleic acid sensor molecules over time to follow the changing concentrations of a protein or metabolite.

The method can also comprise the use of multiplexed chips. A chip can be exposed to a sample solution containing proteins or metabolites to be analyzed. Using multiple  
20 nucleic acid sensor molecules on one chip, the concentrations of many analytes can be shown simultaneously. This can also be extended to cellular assays as shown in Figure 72.

TaqMan can also be used in Protein-PCR, as shown in Figure 58. Figure 58 shows solution-based detection of ligase activity using real-time RT-PCR. Ligation reactions (10 mM MgCl<sub>2</sub>, 100 mM NaCl, 30 mM Tris, pH 7.5) were performed for 0.5-2 hours at room  
25 temperature using 1  $\mu$ M 3-piece construct 19 (DG.20.58E), (SEQ ID NO:111) ligase, 2  $\mu$ M effector oligonucleotide, 2  $\mu$ M substrate oligonucleotide and purified ERK protein. Part A shows the effect of 293 cell extracts on the ERK concentration dependence of ligase activity. The increase in ligation product observed with increasing concentrations of ERK (0-3.2  $\mu$ M) is shown in the presence of buffer only (circles) or buffer containing 10% 293  
30 cell extract (squares). (inset) The ERK2 concentration dependence of ligation is shown as the decrease in cycles of amplification required for detection of ligase product RNA, as measured by real-time RT-PCR. In Panel B ligation in the presence of 1  $\mu$ M ERK and increasing concentrations of the kinase inhibitor staurosporine is shown. ERK nucleic acid

sensor molecules can be used to detect ERK in a concentration dependent manner even in the presence of 293 cell extract. This detection is sensitive to the presence of staurosporine, a kinase inhibitor, which indicates that the activity of ERK is important to the detection of ERK by the nucleic acid sensor molecule.

5        Amplicon dependent nucleic acid amplification technique can be used to show the downstream effects of EGF and MEK inhibitor administration in a cell. The diphenosphorylated ERK is detected in the presence of EGF and not in the presence of the MEK inhibitor as would be expected. This technique can be used to screen other substances for their ability to have ERK phosphorylated. In one embodiment, this technique could be  
10        used to screen for carcinogens in certain cell types.

**Example 22. Analysis of radiolabeled ligation products in 3, 2, 1-piece systems.**

The ERK-modulated 3-piece ligase is converted to the 1-piece ligase NASM by displacement of the effector oligonucleotide binding domain by a stable tetraloop, as  
15        depicted in Figure 36. The deletion of the effector oligonucleotide binding domain of construct 27 (in general), and construct 27, specifically, to generate construct 28 has no effect on the switch factor of the 1-piece ligase NASM (in comparison to the 3-piece ligase NASM), nor does it have an effect on the ERK-modulation KD value for the 1-piece ligase NASM (in comparison to the 3-piece ligase NASM), Figure 37. The exact sequence  
20        constraints for the displacement of the effector oligonucleotide binding domain in construct 27 to generate construct 28, were in fact delineated by the analysis obtained from the lysozyme-modulated 1-piece ligase NASMs shown in Figure 26. The data in Figure 26 also indicates that the length of the stem region proximal to the tetraloop has a significant effect on ligase (e.g. self-circularization) efficiency. The self-circularizing ligase NASMs are then  
25        used as biosensors in intracellular target modulation assays, Figure 30. Detailed assay conditions are described in subsequent Examples.

Additionally, the ERK-modulated 3-piece ligase is converted to the 2-piece ligase NASM by linking the effector oligonucleotide substrate to the oligonucleotide substrate (Figure 38), effectively creating the oligonucleotide supersubstrate, which base pairs with  
30        both the effector oligonucleotide binding domain and the oligonucleotide substrate binding domain. The 2-piece ligase NASM configuration has particular utility in chip applications, described in detail in subsequent Examples (cf., Figures 75 and 79).

Ligation reactions containing <sup>32</sup>P-labeled ligase RNA were analyzed by 8-12% PAGE to separate unligated and ligated products, and quantified using a PhosphorImager

(Molecular Dynamics). Initial rates of ligation were calculated by linear regression from plots of the increase in ligation product over time.

Analysis of ligation products can also be performed by amplicon dependent nucleic acid amplification (ADNA). Amplicon dependent nucleic acid amplification (ADNA) was performed using the TaqMan® One-Step RT-PCR protocol from the same Applied Biosystems. In a typical assay, approximately 0.7 nM ligase RNA (diluted from the EDTA-quenched ligation reaction) was mixed in 25 µL of TaqMan® One-Step master mix with 300 nM each of both the forward (5'-GCGACCTTACGATCAGATGAC (SEQ ID NO:317)) and reverse (5'-CCGCACCTAACCTCCTGTCTAA (SEQ ID NO:318)) PCR primers (Integrated DNA Technologies), and 350 nM TaqMan® fluorescent probe oligonucleotide (ERK-ligase: 5'-6FAM-AAGGAGGATTTCCGAAAGCGGCTACG-TAMRA (SEQ ID NO:319); ppERK-ligase: 5'-6FAM-CGCTAGCGAATTGGTTCCTCGAAAGG-TAMRA (SEQ ID NO:320)). RT-PCR amplification was performed in the ABI Prism™ 7000 Sequence Detection System by incubating the reaction mixes at 48 °C for 30 min to generate the cDNA complement of the ligated RNA product, followed by 40 cycles of PCR amplification. The increase in fluorescence upon amplification is quantified in units of C<sub>T</sub> (threshold cycles), equal to the number of cycles of amplification required for sample fluorescence to exceed background levels. The molar quantity of ligation product can be calculated from these data by comparison with a standard curve generated using a series of known product (ligated biosensor) concentrations.

**Example. 23 Measurement of affinity between ERK protein and ERK-dependent nucleic acid sensor molecules.**

The affinity of the ERK protein target for the ERK-dependent nucleic acid sensor molecule can be measured directly, by a filter-binding assay, or indirectly, by an activity-based assay. In the direct method, increasing concentrations of ERK protein target (0-10 µM) are equilibrated with trace (<1 nM) concentrations of <sup>32</sup>P-labeled, ERK-dependent nucleic acid sensor. The samples are drawn through a sandwich of Protran BA85 nitrocellulose (Schleicher and Schuell) and Hybond P nylon (Amersham-Pharmacia) filter papers, using a 96-well vacuum manifold. Protein and protein-bound RNA species adhere to the nitrocellulose, and free RNA to the nylon. The concentration of ERK protein/sensor complex is plotted as the fraction of the total radioactivity per sample bound to the

nitrocellulose versus total ERK protein concentration. The dissociation constant ( $K_D$ ; Figure 59) is determined as the ERK protein concentration at which 50% of the nucleic acid sensor RNA is bound to protein.

In the indirect method, the activity of the ERK-dependent nucleic acid sensor molecule is measured as a function of increasing ERK protein concentration. In a typical measurement, 50 nM nucleic acid biosensor is incubated with ERK protein (0-10  $\mu$ M) for one hour at room temperature. The extent of ligation at each concentration of ERK protein is determined by real-time RT-PCR (Figure 60, Panels C and D). The ligation product (nM) is plotted against ERK protein concentration, and the apparent dissociation constant ( $K_{act}$ ) is determined as the protein concentration at which the extent of ligation is half-maximal. Panels A and B show ERK-dependent ligation is determined by RT-PCR on an agarose gel containing ethidium bromide.

#### Example 24. High throughput screening (HTS) assays using ppERK nucleic acid sensor molecules

The competitive inhibitor nucleic acid sensor molecule can be used in a HTS assay. A nucleic acid sensor molecule specific for phosphorylated ERK is used as a competitive inhibitor for ATP binding (Seiwert, Stines Nahreini et al. 2000). A competitive assay for compounds is established by incubating ERK with 10 nM nucleic acid sensor molecule in the presence of various concentrations of the inhibitors in 10 mM Tris buffer pH 7.5 containing 0.5  $\mu$ g/ $\mu$ l tRNA and 10 mM  $MgCl_2$ . The reactions are terminated by addition of EDTA and the amount of reacted nucleic acid sensor molecule has been determined.

In other embodiment, the nucleic acid sensor molecules that are sensitive for the phosphorylation state of peptides (or protein substrates) can be used in HTS assays for kinase activity. The nucleic acid sensor molecules can be generated to be specific for the phosphorylation state of substrates or its peptides. HTS assay can be performed using these substrate (Mansour, Candia et al. 1996). For example, MEK can be expressed as GST-tagged protein and purified by a standard method. The activity are measured at 30°C under standard reaction conditions of 20 mM HEPES (pH 7.4), 2 mM dithiothreitol, 0.01% Triton X-100, 10 mM  $MgCl_2$ , 0.1 mM ATP, and 1M His-tagged ERK, at concentrations of MEK, 0.5 nM, in the presence of various concentration of compounds. After incubation, the reaction is terminated by absorbing MEK with GST-column. Phosphorylation of ERK is

quantified by nucleic acid sensor molecule. Alternatively, the nucleic acid sensor molecule can be used in western-blotting format (Bianchini, Radrizzani et al. 2001).

The nucleic acid sensor molecule specific for phosphoERK (ppERK) described above binds to ppERK, presumably in a competitive manner at the active site of the kinase enzyme, and inhibits the phosphorylation of the ppERK substrate by ppERK enzyme. Up to 200 nM nucleic acid sensor molecule inhibits up to 80% of the ppERK phosphorylation of ERK substrate.

**Example 25. *In vivo* assays using phosphorylation state-sensitive nucleic acid sensor molecules**

The phosphorylation state sensitive ppERK nucleic acid sensor molecule is used to determine drug efficacy *in vivo* (e.g., tissue and cell culture). For example, T84 cells on glass coverslips are incubated in the presence and the absence of the MEK kinase inhibitor, and cells are fixed by 4% paraformaldehyde and permeabilized using 0.3% Triton X-100. The slides are incubated with FRET nucleic acid sensor molecule (Bianchini, Radrizzani et al. 2001). The localization of phosphorylated substrate can be observed using a confocal microscope.

Alternatively, phosphorylation-state-sensitive nucleic acid sensor molecule are incorporated into reporter-gene constructs. These constructs are introduced into cells, and phosphorylation of the substrates is monitored.

Nucleic acid sensor molecules made of nuclease resistant forms of hammerhead, -ligase or -hairpin ribozymes are transfected into mammalian cells using standard lipofectamine reagents or liposomal solutions known to effect internalization an cellular uptake of polynucleotides. If desired, the nucleic acid sensor molecules are attached to polypeptides such as tat or antennapodia or the like and are transported into mammalian cells by methods known in the art.

The activity of nucleic acid sensor molecule systems is followed by changes in fluorescence if the nucleic acid sensor molecule is fluorescently tagged, i.e., is an optical nucleic acid sensor molecule or by changes in size as determined by RT-PCR.



**Example 26. Detection of analytes in crude biological samples.**

Nucleic acid sensor molecules can also be used to detect the presence of and measure the concentration of non-nucleic acid analytes in crude biological samples as shown in Figure 56.

As noted above, nucleic acid sensor molecules can be created whose activity is activated in the presence of a specific target molecule. These nucleic acid sensor molecules together with any required substrates, can be combined with a complex mixture of biological origin and allowed to undergo reaction. Subsequent quantification of the product of the reaction can be used to infer the concentration of the analyte in the sample.

In a preferred embodiment, the crude biological sample is human serum or mammalian cell extract. The nucleic acid sensor molecule catalyzes self-ligation with an oligonucleotide substrate. RNase-inhibitors such as ribonucleoside vanadyl complex can be used to prevent degradation of the product. The product of catalysis can be measured by a variety of means (e.g., a gel mobility shift assay as shown in Figure 50).

Ribonucleoside vanadyl complex (RVC) can be used as a general purpose inhibitor of endogenous RNases to enable the use of allosteric ribozymes in assays of samples containing RNase activity (e.g. human serum, mammalian cell extracts, bacterial extracts). RVC inhibits RNA degradation mediated by generic RNases as well as RNase H-like activities that specifically target RNA-DNA hybrids. Inhibition of RNase activity prevents degradation of the reaction product generated by ribozyme action that ultimately forms the basis for readout in a ribozyme-based assay. In a specific example, a lysozyme-dependent nucleic acid sensor molecule RNA with the sequence

5'GGACCUCGGCGAAAGCUAACGUCUCAUGGCUAAAUUGCCAUGUUGCUACAA  
AUGAU AUGACUAGAGAGGUUAGGUGCCUCGUGAUGUCCAGUCGC (SEQ ID  
NO:312) is prepared through transcription off a double-stranded DNA template using T7 polymerase (Ambion MegaShortscript kit, Ambion, Austin, TX), including trace amounts of [ $\alpha$ - $^{32}$ P]UTP. For the ligation reaction, 1  $\mu$ M nucleic acid sensor molecule RNA, 2  $\mu$ M DNA effector oligo 5' GCGACTGGACATCACGAG (SEQ ID NO:313) and 2  $\mu$ M oligonucleotide substrate 5' GTACGATGCGATGCTAGCGATTGTTgugcacu (SEQ ID NO:314) (DNA: upper case; RNA: lower case) were incubated in the presence or absence of 1  $\mu$ M hen egg white lysozyme in reaction buffer (50mM Tris pH 7.4, 100 mM KCl, 10 mM MgCl<sub>2</sub>) as previously described (Robertson MP & Ellington AD, Nature Biotechnol. 2001, 19, 650-655). HeLa extract is added to some of the samples (10% v/v), plus one of various

5 RNase inhibitors (human placenta RNase inhibitor (Sigma, St. Louis, MO), RNasin (Ambion, Austin, TX) and vanadyl ribonucleoside complex, RVC, (Sigma, St. Louis, MO)). After 60 min at room temperature, the reaction is stopped by addition of loading buffer containing formamide, and the samples are analyzed by electrophoresis on a 6% denaturing gel, followed by Phosphorimager scanning (Figure 51, left). The HeLa extract samples that contain protein-based RNase inhibitors show significant broadening of the precursor RNA band compared to a buffer-only control, suggesting RNase H degradation where the RNA is bound to the DNA effector oligonucleotide. The sample containing the VRC RNase inhibitor does not show any significant degradation. All of the samples show lysozyme-dependent product formation. To further test this hypothesis, additional test are done with an analog of the nucleic acid sensor molecules where the oligonucleotide effector domain is deleted

10 (5' GGACCUCGGCGAAAGCUAACGUCUCAUGGCUAAAUUGCCAUGUUGCUACA AAUGAUAUGACUAGAGAGGUUAGGUGC) (SEQ ID NO:332) under otherwise identical conditions (Figure 51, right). No broadening of the RNA bands is observed, suggesting that the RNase activity was indeed specific to the RNA/DNA hybrid region. In a series of related experiments the above described reactions with the lysozyme-dependent nucleic acid sensor molecule are done in buffer to which 10% v/v human serum is added (Figure 53). Without the addition of RVC complete degradation of the RNA is observed. In the presence of VRC, however, no difference in RNA stability and lysozyme-dependent product formation is observed.

20 By inactivating endogenous RNase H-like activity, it is possible to carry out allosteric ribozyme assays in which the ribozymes have been immobilized on a chip surface through hybridization with a tethered DNA probe.

#### 25 **Example 27. Formatting an ERK Nucleic acid sensor molecule chip**

Immobilization of nucleic acid sensor molecules on a substrate provides a straightforward mechanism for carrying out multiple arrays in parallel. Initially, the optimal attachment chemistries were determined to be used to immobilize these molecules on a solid substrate. These molecules are were further configured such that their activity and allosteric behavior is maintained following immobilization. Generally, the chip is configured such that it may be placed at the bottom of a sample holder and overlaid with sample solution, target (ERK and ppERK) (and substrate oligonucleotide). Following an incubation to allow

30

ERK and pERK present within the sample to activate catalysis, the sample is washed away and the extent of ribozyme catalysis quantified.

**Detection.** Immobilized nucleic acid sensor molecules for ERK and pERK are prepared as described in the following section and are assayed for activity by monitoring either retention of end-labeled oligonucleotide substrate (for L1 ligase-based ribozymes) or release of end-labeled ribozyme (for endonucleases as originally described by (Seetharaman et al. 2001). Radioactive tracers are used for labeling RNAs and substrates. To the extent that different attachment chemistries do not interfere, both fluorescent and biotin labels are used in end-labeling the oligo substrate.

**Attachment chemistries.** One advantage of using nucleic acid sensor molecule arrays as opposed to protein arrays is the relative ease with which nucleic acid sensor molecules can be attached to chip surfaces. Several different chemistries for attaching nucleic acid sensor molecules to solid supports were tested and optimized, such as:

1. Conventional DNA arrays using aldehyde coated slides and 5'-amino modified oligonucleotides. The attached oligonucleotide serves as a capture tag that specifically hybridizes to a 3'-end extension on the ribozyme.

2. Nucleic acid sensor molecule RNA treated with periodate to specifically introduce an aldehyde modification at the 3'-end. Modified RNA can be used in two different ways:

- Subsequent reaction with biotin hydrazide enables RNA capture on commercially-available streptavidin coated slides.

- Subsequent reaction with adipic acid dihydrazide enables RNA capture on commercially-available aldehyde coated slides.

3. (Endonuclease) Nucleic acid sensor molecules are generated by transcription in the presence of  $\gamma$ -thio-GTP (introducing a unique thiol at their 5'-end) and subsequently attached to a thiol-reactive surface (e.g. gold-coated polystyrene as described by Seetharaman et al.).

Attachment methodologies are evaluated on the basis of the following criteria:

- **efficiency** – what is the yield of nucleic acid sensor molecule capture?
- **capacity** – what is the maximum concentration of nucleic acid sensor molecules that can be localized in a given spot size?

- **stability** – are ribozymes efficiently retained under a variety of solution conditions and during long-term storage?
- **detection** – do immobilization chemistries interfere with the ability to generate a detectable signal?

5       **Reconfiguring ribozymes for activity in solid phase applications.** To the extent, that allosteric activity for immobilized ERK and pERK nucleic acid sensor molecules is diminished, three different strategies are possible:

1.       **Immobilization chemistries.** A variety of different immobilization chemistries are compared on the basis of their ability to maintain allosteric behavior. To the extent that they leave different surfaces available for protein effectors to interact with, that they tether different ends of the nucleic acid sensor molecules, and that they position the NASM either directly at the surface or displaced from the surface (in the case of streptavidin capture), different behaviors are observed depending upon the immobilization method. Protein-target activated NASMs have been shown to function in both direct and indirect attachment scenarios.
2.       **Blocking chemistries.** Blocking agents (*e.g.*, carrier proteins) are tested to determine whether losses in allosteric responsiveness are due to non-specific interactions between the allosteric activators and the chip surface.
3.       **Tethers.** Steric effects may cause decreased catalytic activity upon direct end attachment to a solid support. Arbitrary sequence tethers are added as needed to increase the spacing between the attachment end and the core of the ribozyme.

25       While endogenous activators and nucleases do not appear to be an issue for the lysozyme-responsive nucleic acid sensor molecules, the following strategies are used in the eventuality that they are a problem for the ERK/pERK (or other) nucleic acid sensor molecules.

30       **Nucleic acid sensor molecule pre-screening.** As noted above, individual clones isolated from the selection experiment are tested early for allosteric activation in the presence of ERK-depleted extracts. Molecules that respond to endogenous non-specific activators are eliminated from further consideration.

**Negative selection with depleted extracts.** To the extent that all isolated nucleic acid sensor molecules are activated by ERK-depleted extracts, depleted extracts are included in the negative selection step of the selection process.

**Nuclease inhibition.** Commercially available RNase inhibitors and competing  
5 RNase substrates (*e.g.* tRNA) are added to test samples.

**Kinetic considerations.** RNase-mediated degradation of the nucleic acid sensor molecule proceeds at a rate in competition with the rate of nucleic acid sensor molecule catalysis. As such, nucleic acid sensor molecules with fast turnover rates can be assayed for shorter times and are thus less susceptible to RNase problems. Nucleic acid sensor  
10 molecules with fast turnover can be obtained by (1) reducing the length of the incubation during the positive selection step, and/or (2) choosing fast nucleic acid sensor molecules (potentially with less favorable allosteric activation ratios) when screening individual clones emerging from the selection experiment.

**Nuclease-Resistant Nucleic acid sensor molecules.** By carrying out selection in  
15 the presence of nucleases (*e.g.* by including depleted extracts during the negative selection step), the experiment intrinsically favors those molecules that are resistant to degradation.

**Modified RNA.** Covalent modifications to RNA that can render it highly nuclease-resistant are performed. Several of these modifications, including for example 2'-*O*-methylation, are compatible with hammerhead cleavage activity and are used to minimize  
20 non-specific cleavage in the presence of biological samples (Usman and Blatt 2000).

#### **Example 28. Rolling circle amplification.**

A circularized nucleic acid product generated by the target-activated ligase nucleic acid sensor molecule serves as a template to promote rolling circle replication. In most  
25 cases, rolling circle replication initially requires that an oligonucleotide primer be annealed to the circular nucleic acid template. This oligonucleotide primer may be provided subsequent to the nucleic acid sensor molecule reaction that generates the circular template. Alternatively, the primer may be present during the nucleic acid sensor molecule reaction as either a separate oligonucleotide or as part of the nucleic acid sensor molecule itself (Figure  
30 53). Also, the primer may exist free in solution or be tethered either covalently (*e.g.*, via 5'-end modification) or non-covalently (*e.g.*, association of biotinylated primer with immobilized streptavidin or through base-pairing with complementary immobilized oligonucleotide) to a solid support. In a preferred embodiment for solid-state (*e.g.*, chip) applications, an immobilized nucleic acid sensor molecule serves directly as the primer for

rolling circle amplification as illustrated in Figure 53. Immobilization makes it possible to confine the signal generated through the rolling circle reaction to a specific spatial location, enabling the development of arrays in which each element in the array is composed of a different target-activated nucleic acid sensor molecules. In another embodiment, an  
5 immobilized primer is used for rolling circle extension, also leading to a spatially localized signal. Preferably, the primer is at least 10 nucleotides long and it hybridizes to a unique site on the circular template.

Rolling circle amplification may proceed using either a DNA-directed or RNA-directed polymerase, depending upon the nature of the circular template generated by the  
10 target-activated nucleic acid sensor molecules. Concatamers generated from primarily DNA templates can be generated by an enzyme preferably from the following group: Taq DNA polymerase, Tth DNA polymerase, Klenow fragment of *E. coli* DNA polymerase, bacteriophage O29 DNA Polymerase, or Bst DNA polymerase. More preferably, the polymerase enzyme is Bst DNA Polymerase. Concatamers generated from primarily RNA  
15 templates can be generated by an enzyme preferably from the following group: AMV reverse transcriptase, MMLV reverse transcriptase, or Q-beta replicase. More preferably, the polymerase enzyme is MMLV reverse transcriptase.

The rolling circle amplification reaction requires a supply of nucleotide triphosphates for incorporation to generate the nucleic acid product. Quantification of the  
20 rolling circle amplification reaction may be facilitated by using nucleotides containing fluorescent or radioactive labels or affinity tags as described below.

The rolling circle amplification reaction may be made to proceed exponentially by the addition of a linear oligonucleotide primer containing all or part of the circular template sequence (Figure 54). The uncircularized form of the ligase substrate can perform this  
25 function. As such, in the preferred embodiment, the detection reaction is carried out with excess oligonucleotide substrate.

#### Detection

The rolling circle amplification reaction generates single- and double-stranded DNA. The  
30 amount of the rolling circle amplification product generated can be determined by the variety of methods outlined below. The signal measured by each method can be directly related to a corresponding analyte concentration by constructing a standard curve in which a range of known concentrations of the analyte are analyzed in parallel.

*Fluorescent detection.* Products of the rolling circle amplification reaction may be detected fluorescently by the following methods:

- 5 a. *Intercalating dyes.* The fluorescence intensity of several dyes is known to increase upon their binding to DNA. The yield from the rolling circle amplification reaction may be measured by including one of these dyes during the reaction and using a fluorimeter to monitor the change in fluorescence intensity as the reaction proceeds. For this, a variety of nucleic acid stains is readily available (see Handbook of Fluorescent Probes and Research Products, Chapter 8, Molecular Probes, OR).
- 10 b. *Labeled nucleotides.* Fluorescent labels may be attached to the nucleotides that serve as substrates for the rolling circle amplification reaction. Fluorescent nucleotides incorporated into the newly synthesized DNA can be detected by two different mechanisms:

15 Homogeneous assays: nucleotides with appropriate fluorescence donor and acceptor pairs may be combined such that their sequential incorporation into a polynucleotide increases their spatial proximity and, correspondingly, the efficiency of fluorescence resonance energy transfer (FRET) (Figure 55). The resulting FRET signal may be read using a fluorimeter. The strength of this signal may be maximized by appropriate design of the ligase substrate such that  
20 its sequence contains many acceptor-donor dinucleotides. A similar process using a fluorescent UTP and a biotin-CTP derivative, followed by addition of an fluorescence-acceptor streptavidin conjugate has been described for labeling of RNA (Alpha-Bazin et al. Anal. Biochem. (2000) 286, 17-25).

25 Heterogeneous assays: unincorporated nucleotides may be physically removed by covalently or non-covalently capturing the rolling circle amplification reaction product (e.g. by pre-immobilization of the rolling circle amplification primer or by hybridization to an immobilized capture probe) and followed by washing. Fluorescent nucleotides retained with the immobilized nucleic acid product can be detected and quantified (e.g. using a confocal scanner) to obtain  
30 an estimate of the extent of the rolling circle amplification reaction.

*Sequence-specific fluorescent hybridization probes.* Hybridization of a complementary, fluorescently labeled probe with the rolling circle amplification reaction product is used in a variety of ways to generate a fluorescent signal. Strategies include (1) physical retention of a fluorescent probe by captured/immobilized RCA product, (2) co-immobilization of a

fluorophore acceptor-donor oligonucleotide pair (FRET), (3) 5'-nuclease induced cleavage of a fluorophore-quencher oligonucleotide (Taqman<sup>TM</sup>) and (4) conformational changes in a fluorescent probe induced by hybridization which change its fluorescent properties. In a related aspect, the fluorescent probe may be attached to the primer that is used, leading to a conformational change during transcription (e.g. Amplifluor<sup>TM</sup>, Intergen, NY)

5 *Proximity Assays:* In addition to the aforementioned methods that rely on inducing physical proximity of two fluorophores through hybridization of probes with rolling circle amplification products (e.g., FRET, molecular beacons, Taqman<sup>TM</sup> etc.), other proximity-based assays may be used. This includes, but is not limited to, luminescence-based assays such as Amplified Luminescent Proximity Homogeneous Assay (AlphaScreen, Packard BioScience, CT) and Bioluminescence Resonance Energy Transfer (BRET, Packard BioScience, CT).

10 *Radiometric read-out.* By using appropriate radioactively labeled nucleotides in the rolling circle amplification reaction (e.g. alpha-P<sup>32</sup>-ATP), the resulting rolling circle amplification product becomes radioactive and may be quantified using instrumentation that detects radioisotope decay (e.g. phosphorimagers, x-ray film, scintillation counters). In the preferred embodiment, nucleotides provided for the rolling circle amplification reaction are labeled with tritium and the primer for the rolling circle amplification reaction is immobilized on a solid support that contains a scintillant. Co-localization of the

15 radioactively-labeled RCA product on the scintillant surface generates the signal readily detected by a scintillation counter (scintillation proximity assay, SPA).

20 *Surface Plasmon Resonance (SPR).* The primer used to initiate the rolling circle amplification reaction is immobilized on the gold surface of a detection element in an SPR detector (e.g. Biacore). As the rolling circle amplification reaction proceeds, the effective mass of the primer increases, generating a signal that can be detected due to the change in surface plasmon resonance.

25 *Coagulation:* Microspheres bearing a hybridization probe complementary to the rolling circle amplification product or the primers for the rolling circle amplification reaction can lead to coagulation as the reaction progresses. Coagulation can be readily detected by means that include measurement of changes in refractive index or turbidity.

30 *Others.* Rolling circle amplification is capable of generating very long polynucleotides that are very likely to lead to significant changes in the sample's physical properties. Examples include, but are not limited to, changes in viscosity, refractive index, turbidity, and



electrophysical properties such as conductivity. In general, any kind of method that can detect these changes can likely be used to monitor the rolling circle amplification reaction.

**Example 29. High Throughput Screening (HTS) Assay using G protein nucleic acid sensor molecules**

Activated G $\alpha$ -protein dependent nucleic acid sensor molecules are used *in vitro* to test the efficacy of the agonists and inverse agonists for any GPCR. G $\alpha$ -protein cDNA can be obtained (Guthrie cDNA resource Center) and expressed in *E. coli* as His-tagged protein (Lee, Linder et al. 1994). GDP or GTP- $\gamma$ S are added during the purification to avoid degradation. GDP-complex G $\alpha$ -protein and GTP-complex G $\alpha$ -protein are used to generate nucleic acid sensor molecules.

GTP-complex G $\alpha$ -protein (activated-G $\alpha$ -protein)-dependent nucleic acid sensor molecules are tested in a target modulation assay to screen the selected nucleic acid sensor molecules with desirable properties. For example, membrane fraction of C6 glioma cells in 12-well plates are loaded for 16 h in the presence and the absence of adrenaline. Cell lysate can be tested for activated G $\alpha$ -protein-dependent nucleic acid sensor molecule response. Alternatively, the competitive modulation by for adenylyl cyclase is tested following the inhibition of adenylyl cyclase activity by activated G $\alpha$ -protein in the presence of nucleic acid sensor molecules by a modified assay based on a previously described assay (Burt, Sauté et al. 1998).

Adenylyl cyclase activity is measured as described in the presence and the absence of nucleic acid sensor molecules (Kozasa and Gilman 1995). Cell membranes from HeLa cells transfected with human cloned 5-HT<sub>1A</sub> receptors resuspended in buffer are used to screen compounds. The membranes are incubated with 30 M GDP and decreasing concentrations of test drugs (from 100 pM to 0.1 nM) or decreasing concentrations of 5-HT, from 100 M to 0.1 nM (reference curve) for 20 min at 30°C in a final volume of about 0.5 mL (Stanton and Beer 1997). Then samples are added with GTPS and the nucleic acid sensor molecules and then incubated for a further 30 min at 30°C.

Alternatively, nucleic acid sensor molecules are generated that depend on the presence of G $\beta$ -protein uncomplexed with G $\alpha$ -protein. There are four known classes of G $\beta$ -protein. G $\beta$ -protein complex with G $\gamma$ -protein can also play important roles in the signal transduction. After dissociation from alpha subunit, this complex is known to regulate various target protein, such as GRKs, Raf kinase, adenylyl cyclase, PLCs and ion channels.

Producing nucleic acid sensor molecules that block the complex keep the GPCR's from activating their effectors.

**Example 30. Cell-based assays using GPCR nucleic acid sensor molecules**

5 GPCR nucleic acid sensor molecules are used in cell-based assay using modifications of previously described GPCR assays (Hun, Ellington et al. 2001). For example,  $\beta$ -adrenergic receptor can be expressed in CHO cells. CHO cells are grown in multiwell tissue culture plates in Dulbecco's modified Eagle media (DMEM) with 10% fetal bovine serum. On the day of assay, medium is replaced with 0.2 mL treatment medium  
10 containing DMEM media containing 250 M IBMX (isobutyl-1-methylxanthine) plus 1 mM ascorbic acid with test compound dissolved in DMSO. Test compounds are added at a desired concentration range (e.g.  $10^{-9}$  to  $10^{-4}$  M). Isoproterenol ( $10^{-5}$  M) is used as an internal standard for comparison of activity. Cells were incubated at 37°C on a rocker for 15-30 min. Then cells are lysed and the level of the activated G $\alpha$ -protein is measured by the  
15 nucleic acid sensor molecule. The antagonist is screened in the same format in the presence of the known agonist by detecting the decreasing amount of the activated G $\alpha$ -protein.

Alternatively, the agonism and the antagonism of compounds for specific or general GPCR is measured using the nucleic acid sensor molecule. For example, evaluation of compounds for antagonism can be performed using Sprague Dawley rats. The aorta from  
20 animal is isolated and freed of adhering connective tissue. Desmethylinipramine (0.1 M) and corticosterone (1 M) to block neuronal and extraneuronal uptake of noradrenaline, ( $\pm$ )-propranolol (1M) to block  $\beta$ -receptors, and yohimbine (0.1 M) to block  $\gamma$ -receptors are added. Aortic strips are incubated with various concentration of compounds in the presence and the absence of 10 M noradrenaline. Then the cell extract is prepared and the activated  
25 G $\alpha$ -protein level in the cell is measured with the nucleic acid sensor molecules (Barlocco, Cignarella et al. 2001).

**Example 31. *In vitro* assay for GPCR activation by following liberation of G $\alpha$ -protein with G $\alpha$ -protein dependent nucleic acid sensor molecules.**

30 Nucleic acid sensor molecules generated against G $\alpha$ -protein subunits are used to test the efficacy of agonists and inverse agonists for any GPCR *in vitro* using activated G-protein. For example, G $\alpha$ -protein cDNA can be obtained (Guthrie cDNA resource Center) and expressed in E coli as His-tagged protein (Lee, Linder et al. 1994). GDP or GTP- $\gamma$ S are

added during the purification to avoid degradation. GDP-complex  $G_{i\alpha}$ -protein and GTP-complex  $G_{i\alpha}$ -protein are used to generate nucleic acid sensor molecules. GTP-complex  $G_{i\alpha}$ -protein (activated- $G_{i\alpha}$ -protein)-dependent nucleic acid sensor molecule can be tested for the binding assay to screen the selected nucleic acid sensor molecules with desirable properties. For example, membrane fraction of C6 glioma cells in 12-well plates were loaded for 16 h in the presence and the absence of adrenaline. Cell lysate can be tested for activated  $G_{i\alpha}$ -protein-dependent nucleic acid sensor molecule response. Alternatively, the competitive binding for adenylyl cyclase can be tested following the inhibition of adenylyl cyclase activity by activated  $G_{i\alpha}$ -protein in the presence of nucleic acid sensor molecules by a modified assay based on a previously described assay (Burt, Sautel et al. 1998).

Adenylyl cyclase activity can be measured as described in the presence and the absence of nucleic acid sensor molecules (Kozasa and Gilman 1995). Cell membranes from HeLa cells transfected with human cloned 5-hydroxytryptamine 1A ( $5-HT_{1A}$ ) receptors resuspended in buffer can be used to screen compounds. The membranes are incubated with 30 mM GDP and decreasing concentrations of test drugs or decreasing concentrations of 5-HT, from 100 M to 0.1 nM (reference curve) for 20 min at 30 °C in a final volume of about 0.5 mL (Stanton and Beer 1997). Then samples are added with GTPS and the nucleic acid sensor molecules and then incubated for a further 30 min at 30°C.

Nucleic acid sensor molecules can also be generated that depend on the presence of  $G\beta$ -protein uncomplexed with  $G\alpha$ -protein. There are four known class of  $G\beta$ -protein.  $G\beta$ -protein complex with  $G\gamma$ -protein can also play important roles in the signal transduction. After dissociation from alpha subunit, this complex is known to regulate various target protein, such as GRKs, Raf kinase, adenylyl cyclase, PLCs and ion channels. Producing nucleic acid sensor molecules that block the complex keep the GPCR's from activating their effectors.

### Example 32 Multiplex assay for $G\alpha$ proteins

Nucleic acid sensor molecules are also generated whose activity is dependent on multiple activated G-proteins. Human G-protein cDNAs are obtained from publicly available databases or are cloned by RT-PCR from polka-RNA pool of appropriate source. They can be expressed as described above and use to select nucleic acid sensor molecules. The readouts for multiplex assay system are discussed above.

**Example 33. Nucleic acid sensor molecules activated by native lysozyme.**

Lysozyme activated nucleic acid sensor molecules lys.L1.A, B, and C are disclosed in Figure 26. Stems above the ligation junction varied in length from 2, 5 and 10 base pairs. The unimolecular NASMs were generated by PCR of the original lysozyme NASM with 5' primer (TCTAATACGACTCACTATAGGACCTCGGCGAAAGC) (SEQ ID NO:296) and the following 3' primers: A ACTCTCGCTAACCTCTCTAGTCATA (SEQ ID NO:297), (B) AGTGCTCTCGCACCTAACCTCTCTAGT (SEQ ID NO:298) or (C) AGTGCGAGCCTCTCGGCTCGCACCTAACCTCTCTAGT (SEQ ID NO:299) using Taq polymerase (Invitrogen) following the manufacturer's protocol. RNAs were generated with Ampliscribe T7 transcription kits following the manufacturer's protocol. RNA was heated in water for 3 min at 70 °C and then brought to room temperature before adding 5X buffer (5X buffer: 150 mM Tris, 50 mM MgCl<sub>2</sub>, pH 7.5) and lysozyme (1 uM or absent). Ligations were performed at room temperature for one hour then quenched with stop dye [formamide, 0.1 % bromophenol blue, xylene cyanol] and analyzed on a 6% acrylamide gel. Self-ligation was increased in the presence of lysozyme for constructs B and C. No ligation was observed for construct A, suggesting that the stem length above the ligation junction is critical.

This circularization can be detected using PCR across the ligation site as shown in Figure 27. Activation of (clone B is shown in Figure 28. Protocols from the supplier were followed for Superscript RT and recombinant Taq Polymerase (Invitrogen). All RNA samples were reverse transcribed and PCR amplified separately with two different sets of primers. The first set was designed to amplify the total NASM RNA that was used as input into *in vitro* (TCTAATACGACTCACTATAGGACCTCGGCGAAAGC (SEQ ID NO:300) and AGTGCTCTCGCACCTAACCTCTCTAGT (SEQ ID NO:301)). The second set of primers was designed to selectively amplify the RNAs that circularized (GTTGCTACAAATGATATGAC (SEQ ID NO:302) and ATGGCAATTTAGCCATGAGA (SEQ ID NO:303)). Following amplification with RT/PCR the circularized species display a ladder like pattern as a result of strand displacement. The ladder can be avoided by the use of a reverse transcriptase with RNase H activity. The gel shows in Figure 28 that after 25 cycles of PCR, a circularization signal appears. This signal allows for quantitation of the degree of activation of ligase in response to their target molecules.

Ligase nucleic acid sensor molecules are also able to detect their targets in cell extracts. In Figure 29, (clone B is combined with lysozyme, and increasing amounts of HeLa cell extracts. Experiments were carried at as described for Figure 28 with HeLa cell

extract added in place of water. Even at the highest HeLa cell extract concentration, activation of clone B by lysozyme is still detectable.

Lysosome-dependent ligase (lys11-2 from Nature Biotechnology 19, 650-655 (2001)) incubated in the presence of effector oligonucleotide 18.90a (ibid.), substrate S28A (ibid.), optionally 10% cellular lysate (HeLa cells or reticulocyte lysate) and various concentrations of lysozyme at 25 °C for 1 hour, as shown in Figure 50. Ligation is observable at 100 nM and is ~30% at 1 µM. Ligation yield is unaffected by the presence of 10% reticulocyte lysate or 10% HeLa lysate.

Ligase nucleic acid sensor molecules are also able to detect their targets *in vivo*.

10 Mammalian cells were electroporated with 1, 5, 10 and 25µg RNA. The RNA was brought to a total volume of 30 ul in the presence of 20 ug salmon sperm DNA. Cells were trypsinized, counted, pelleted and resuspended in 250 ul of media. The ligase biosensor/Salmon sperm DNA mixture was added to the cells, mixed gently and incubated at room temperature for 5 minutes. The cells were electroporated and diluted in pre-

15 warmed media and incubated at ambient temperature for 5 minutes. Subsequently, the cells were plated and transferred to a 37 °C incubator and allowed to recover for 6 hours. Following recovery, the cells were trypsinized, pelleted, washed with PBS and pelleted again. Cells were resuspended in an ice cold mixture of 150 mM NaCl, 1 µl of 10% NP-40 and 6 µl 10% SDS, 28 µl 10 mM Tris-HCl, pH 8.0 solution and added to a mixture of 40

20 Proteinase K (20 mg/ml). The lysate was passed through a 1ml 26G 5/8 syringe 20 times. Lysates were incubated at 37 °C for 2 hours. Following the incubation the slurry was phenol/chloroform/isoamyl alcohol extracted two times, the aqueous phase was removed and ethanol precipitated RT-PCR reactions were carried as described for figure 30. Extent of ligation and circularization was determined by RT-PCR as shown in Figure 30B, the cells

25 that were cotransfected with lysozyme showed an increased circularization.

**Example 34. RT-PCR analysis of ligase nucleic acid sensor molecules/amplicon-dependent nucleic acid amplification.**

RT-PCR is a quantitative method that can be used to measure the activity of a ligase

30 nucleic acid sensor molecule. As shown in Figure 46, PCR primers are used that hybridize to sequences of the ligase nucleic acid sensor molecule that are only joined after self-ligation. After ligation occurs the nucleic acid sensor molecule can be amplified, and the PCR product can be detected on an agarose gel, using ethidium bromide or other nucleic



acid staining techniques. Alternatively, the progress of the PCR reaction can be monitored using real-time detection techniques including, but not limited to, Taqman (Applied Biosystems, Foster City, CA), SYBR Green (Molecular Probes, Eugene, OR and Applied Biosystems, Foster City, CA), Scorpion (Eurogentec, Liege, Belgium), Amplifluor (Serologicals, Norcross, GA) or related systems.

A specific example of a lysozyme-dependent ligase nucleic acid sensor molecule is as follows:

The ligase RNA has the sequence

5'GGACCUCGGCGAAAGCUAACGUCUCAUGGCUAAAUUGCCAUGUUGCACAA  
10 AUGAUAUGACUAGAGAGGUUAGGUGCAUCUUCAUGUCCAGUCGCUUGCAAU  
GCCC (SEQ ID NO:304) and is prepared through transcription off a double-stranded DNA template using T7 polymerase (Ambion MegaShortscript kit, Ambion, Austin, TX). For the ligation reaction, 200 nM nucleic acid sensor molecule RNA, 1  $\mu$ M DNA RT primer 5'GGGCATTGCAAGCGACTGGACAT (SEQ ID NO:305) and 1  $\mu$ M substrate oligo  
15 5'ACTGAACCTGACCGTACAAAGATgcac (SEQ ID NO:306) (DNA: upper case; RNA: lower case) are incubated with 1nM - 10  $\mu$ M hen egg white lysozyme in reaction buffer (50 mM Tris pH 7.4, 100 mM KCl, 10 mM MgCl<sub>2</sub>) as previously described (Robertson MP & Ellington AD, Nature Biotechnol. 2001, 19, 650-655). After 15 min at room temperature, the reaction is stopped by diluting 300-fold with 1 mM EDTA. Of this, 5  
20  $\mu$ L are transferred into 25  $\mu$ L of a RT-PCR reaction mix (Applied Biosystems, Foster City, CA) that contains 300 nM DNA forward primer 5'ACTGAACCTGACCGTACAAAGA (SEQ ID NO:307) and DNA reverse primer 5'TTTGTAGCAACATGGCAATTTA (SEQ ID NO:308) plus 350 nM Taqman probe 5'6FAM-CGGCGAAAGCTAACGTCTCATGG-TAMRA (SEQ ID NO:309) (Applied Biosystems, Foster City, CA). Reverse transcription is  
25 performed through incubation for 30 min at 48 °C. The sample is then denatured for 10 min at 95 °C followed by 40 PCR cycles (15 sec at 95 °C, 1 min at 60 °C). The amplification process is monitored using a ABI 7000 instrument (Applied Biosystems, Foster City, CA), and threshold cycle values are calculated using the instrument software. Plotting the Ct against the logarithm of the lysozyme concentration reveals a largely linear relationship  
30 (Figure 48). The linear dynamic range spans about three orders of magnitude from ~1 nM to ~1  $\mu$ M. A control reaction without the addition of lysozyme gives a threshold cycle values of ~24.5. This value defines the upper limit for threshold cycle values, and is due to the small amount of product that is inevitably generated through the background ligation. On the opposite end, the lower plateau of threshold cycle values is reached with lysozyme > ~1

$\mu\text{M}$ . This concentration is close to the apparent  $K_d$  ( $1.5 \mu\text{M}$ ), and the nucleic acid sensor molecule target modulation domain is beginning to be saturated by lysozyme, triggering a reaction at maximum catalytic rate. Another specific example describes the use of the SYBR Green RT-PCR amplification detection system (Applied Biosystems, Foster City, CA). Ligation reactions are run as described above, and analysis is done through transferring  $5 \mu\text{L}$  of the reaction into  $25 \mu\text{L}$  of a SYBR-Green RT-PCR reaction mix (Applied Biosystems, Foster City, CA) that contains  $100 \text{ nM}$  DNA forward primer 5'ACTGAACCTGACCGTACAAAGA (SEQ ID NO:310) and DNA reverse primer 5'TTTGTAGCAACATGGCAATTGA (SEQ ID NO:311). Reverse transcription is performed through incubation for 30 min at  $48^\circ\text{C}$ . The sample is then denatured for 10 min at  $95^\circ\text{C}$  followed by 40 PCR cycles (15 sec at  $95^\circ\text{C}$ , 1 min at  $60^\circ\text{C}$ ). The amplification process is monitored using a ABI 7000 instrument (Applied Biosystems, Foster City, CA), and threshold cycle values are calculated using the instrument software as above (Figure 49). The threshold cycle values are directly proportional to the logarithm of the lysozyme concentration between  $\sim 1 \text{ nM}$  to  $\sim 1 \mu\text{M}$ . A melting profile confirms that a uniform amplification product was obtained, without the appearance of artifacts such as primer-dimers.

**Example 35. Ligation reactions in 3 or 2 piece systems.**

Ligation reactions with varying volumes depending on the application type were performed in buffer 1, [30 mM Tris-HCl, pH 7.5, 10 mM  $\text{MgCl}_2$ , 50  $\mu\text{g/mL}$  tRNA type X (Sigma)], or buffer 2 [20 mM HEPES, pH 7.4, 10% glycerol, 150 mM KCl, 10 mM  $\text{MgCl}_2$ , 1 mM EDTA, 1 mM DTT, 0.5 mg/ml tRNA, 2 U/ $\mu\text{L}$  RNaseOUT™ (Invitrogen)]. Ligation reactions performed in the presence or absence of crude cell lysates contained 200 nM or 50 nM ligase RNA, respectively. Ligase RNA was pre-annealed with a 2-fold molar excess of effector oligonucleotide by incubation for 3 min at  $70^\circ\text{C}$ , followed by the addition of 5 $\times$  reaction buffer at  $25^\circ\text{C}$ . The appropriate concentration of protein and/or cell lysate was added to the RNA and incubated for an additional 15 min at room temperature. Ligation reactions were then initiated by the addition of this mixture to a 10 $\times$  solution of the substrate oligonucleotide TK.04.82.C ACGTAGCATAGCATCGATAGCTGTTGugcacu (SEQ ID NO:357) (small letters correspond to RNA nucleotides) final concentration =  $1 \mu\text{M}$ ). The reactions were incubated at  $25^\circ\text{C}$ , and small aliquots (2–10  $\mu\text{L}$ ) of reaction mix were quenched by dilution in EDTA.



**Example 36. Transfection of ligase biosensors into mammalian cells by electroporation.**

HEK 293, 3T3 or HeLa cells were grown in DMEM medium containing 10% fetal bovine serum to 85-90% confluency, and harvested by trypsinization. Aliquots of  
5 approximately 250,000 cells were pelleted by centrifugation (3 min at 2000 rpm), resuspended in 250  $\mu$ L of growth medium, and gently mixed with 30  $\mu$ L of nucleic acid mix containing 20  $\mu$ g of salmon sperm DNA (carrier) and 1, 5, 10 or 25  $\mu$ g nucleic acid sensor molecule ligase RNA. This mixture was incubated at room temperature for exactly 5 minutes without agitation, and transferred to electroporator cuvettes. An electric field pulse  
10 was applied to the cells at ambient temperature using a Gene Pulser II power supply (BioRad; settings: 250V, 0.975  $\mu$ F high capacitance). The cells were plated in pre-warmed growth medium, incubated at 37 °C for varying periods, and harvested. Cells were washed with PBS, pelleted, and immediately subjected to total, cytoplasmic, or nuclear RNA isolation procedures or flash-frozen in liquid nitrogen and stored at -80 °C for further  
15 analysis.

Purification of total RNA from mammalian cells Pelleted cells were resuspended in ice old buffer (150 mM NaCl, 10mM Tris HCl, pH 8.0) and added to 40  $\mu$ L of 10% SDS, 28  
20  $\mu$ L of 10% NP40 and 6  $\mu$ L of Proteinase K (20  $\mu$ g/ ml). The mixture was drawn into a 1 ml syringe and passed through a 26 gauge needle at least 20 times, and then incubated at 37 °C for 2 hours. Following incubation, the slurry phenol/chloroform/isoamyl alcohol extracted two-times, the aqueous phase was removed and ethanol precipitated. The sample was 70% ethanol washed, and dried. The RNA pellets were resuspended and the yield was quantified based on OD 260. The cleanliness of the prep was assessed based on 260/280 ratio. All  
25 RNA samples were stored at -80 °C.

**Example 37. Analysis of ligation and/or circularization by RT/PCR:**

Protocols from the supplier were followed for reverse transcription and PCR. Either Superscript RT or AMV RT, and recombinant Taq Polymerase (Invitrogen) were used.

In order to avoid any occurrence of ligation in the detection phase of experiments,  
30 all reverse transcriptions were performed at 48 °C. All RNA samples were reverse transcribed and PCR amplified separately with two different sets of primers. First set was designed to amplify the total RNA that was used as input into *in vitro* or *ex vivo* assays. The second set of primers were designed to selectively amplify the RNAs that were ligated

to the substrate oligo (resulting in a higher molecular weight) or were ligated to their 5' end and became circularized without molecular weight change. two and three-piece ligase platforms give rise to higher molecular weight products as a result of target dependent ligation. One piece ligase platform circularize as a result of target dependent ligation.

- 5 When observed directly with gel based assays the circularized form has a different mobility than the linear RNA. Following amplification with RT/PCR the circularized species display a ladder like pattern as a result of strand displacement that happens in the reverse transcription phase. The ladder like pattern can be avoided by the use of a reverse transcriptase with RNase H+ activity.

10

**Example 38. Preparation of extracts from mammalian cell cultures:**

- After stimulation with the reagent that regulates target levels, mammalian cells were rinsed briefly in cold TBS and then lysed in TBS lysis buffer (137 mM NaCl, 20 mM Tris, pH 8.0, 1% vol/vol NP-40, and 10% vol/vol glycerol; Knusel *et al.* 1994) or Brij Lysis  
15 buffer (150 mM NaCl, 10 mM Tris, pH 7.5, 2mM EDTA, 0.125% vol/vol NP-40, 0.875% vol/vol Brij and 10% vol/vol glycerol). Lysis buffers were supplemented with Mini Complete protease inhibitor cocktail (Boehringer Mannheim Corp.) and 1.5 mM sodium vanadate. Lysates were scraped into Eppendorf tubes and rocked for 30 min at 4 °C. Samples were then cleared by centrifugation at 10,000 x g for 10 min at 4 °C. Protein  
20 concentration was determined by the BCA assay (Pierce Chemical Co.) using BSA as a standard.

**Example 39. Intron-based catalytic NASMs**

- Figure 61 illustrates an optical NASM derived from an intron-based catalytic  
25 NASM. The optical nucleic acid sensor molecule derived from the group I intron splices *in vitro* such that it excises itself from a larger nucleic acid sequence and subsequently ligates, or joins together the two terminal portions of the RNA molecule (*i.e.*, the 5' and 3' exons). The distal termini of the 5' and 3' exons are modified to incorporate the donor and acceptor components of a FRET signaling pair. In the absence of target modulation, fluorescence  
30 emission will be observed only from the donor fluorophore which is being directly excited optically. Splicing and ligation of the 5' and 3' exons brings the donor and acceptor molecules on the exons into close proximity, resulting in efficient fluorescence energy transfer (FRET) and thus a measurable increase in the fluorescence emission signal from the acceptor fluorophore (with a corresponding decrease in fluorescence emission from the

donor fluorophore). In an alternative embodiment (not shown), the acceptor molecule may be replaced with a quencher molecule (*i.e.*, an acceptor species which absorbs but does not re-radiate fluorescent energy). In this case, the splicing and ligation event will only result in a measurable decrease in the fluorescence emission signal from the donor fluorophore.

- 5       The thymidylate synthase ("td") intron (Construct 41)(SEQ ID NO:358) -  
 TAATTGAGGCCTGAGTATAAGGTGACTTATACTTGTAATCTATCTAAACGGGGA  
 ACCTCTCTAGTAGACAATCCCGTGCTAAATTGTAGGACTGCCCGGGTTCTACAT  
 AAATGCCTAACGACTATCCCTTTGGGGAGTAGGGTCAAGTGACTCGAAACGAT  
 AGACAACTTGCTTTAACAAGTTGGAGATATAGTCTGCTCTGCATGGTGACATGC  
 10 AGCTGGATATAATTCCGGGGTAAGATTAACGACCTTATCTGAACATAATG - is a  
 group I intron sequence of 265 nucleotides in length. Other lengths and their sequences are  
 known in the art and can be used as described herein in lieu of SEQ ID NO:358 above.

- An example of an intron-derived optical NASM that is modulated by target  
 recognition is the theophylline dependent td group I intron which splices itself out of a  
 15 thymidylate synthase gene. The intron sequence of this optical NASM is as follows:

- TAATTGAGGCCTGAGTATAAGGTGACTTATACTTGTAATCTATCTAAACG  
 GGGAACCTCTCTAGTAGACAATCCCGTGCTAAATTGATACCAGCATCGTCTTG  
 ATGCCCTTGGCAGCATAAATGCCTAACGACTATCCCTTTGGGGAGTAGGGTCA  
 AGTGACTCGAAACGATAGACAACTTGCTTTAACAAGTTGGAGATATAGTCTGCT  
 20 CTGCATGGTGACATGCAGCTGGATATAATTCCGGGGTAAGATTAACGACCTTAT  
 CTGAACATAATG (construct 42)(SEQ ID NO:359) wherein the bolded nucleotides  
 represent a target modulation domain that recognizes theophylline.

- The complete optical NASM comprises the intron sequence described above and the  
 5' and 3' exons attached thereto. Exon 1, the 5'-exon - TTTCTTGGGT (construct 43)(SEQ  
 25 ID NO:360) - is designed to contain a first fluorescent donor molecule, such as Cy3  
 (Amersham Biosciences) coupled to a nucleotide in the vicinity (*i.e.*, within no more than  
 about 15 nucleotides) of the 3' terminus of the 5' exon. The 5'-exon length can vary with a  
 minimum of 6 nucleotides of complementarity to P10. Exon 2, the 3'-exon -  
 CTACCGTTTA (construct 43)(SEQ ID NO:361) - is designed to contain a second  
 30 fluorescent molecule, such as Cy5 (Amersham Biosciences) coupled to a nucleotide in the  
 vicinity (*i.e.*, no more than about 15 nucleotides) of the 5' terminus of the 3' exon. The 3'-  
 exon length can vary with a minimum of 2 nucleotides of complementarity to P10.

Intron Splicing conditions are as follows: the intron (500 nM) is heated in H<sub>2</sub>O to 70  
 °C for 3 minutes then put on ice for 1 minute. Splicing buffer (20 mM Tris-HCl, pH 7.5,

100 mM KCl, 3 mM MgCl<sub>2</sub>), is added and the reaction is incubated on ice for an additional 15 minutes. At this time a 4.5 µl aliquot is removed for a time zero point and quenched with 5 µl stop dye (95% formamide, 20 mM EDTA, 0.5% xylene cyanol, and 0.5% bromophenol blue). GTP (50 µM) is added to the remaining reaction (4.5 µl) to start the splicing  
5 reaction. The reaction is incubated at 37 °C for 1 hour and then terminated with stop dye (5 µl). The reactions are heated to 70° C for 3 minutes. Splicing of full length RNA transcripts is detected by gel analysis of labeled RNA.

The theophylline-dependent optical NASM splices *in vitro* under the same conditions as described for the intron splicing reaction, but with theophylline added to a  
10 final concentration 1.5 mM in the splicing buffer. The theophylline-dependent optical NASM (500 nM) is heated in H<sub>2</sub>O to 70 °C for 3 minutes then put on ice for 1 minute. Splicing buffer (20 mM Tris-HCl, pH 7.5, 100 mM KCl, 3 mM MgCl<sub>2</sub>) and theophylline (1.5 mM) are added and the reaction is incubated on ice for an additional 15 minutes. At this  
15 time a 4.5 µl aliquot is removed for a time zero point and quenched with 5 µl stop dye (95% formamide, 20 mM EDTA, 0.5% xylene cyanol, and 0.5% bromophenol blue) (95% formamide, 20 mM EDTA, 0.5% xylene cyanol, and 0.5% bromophenol blue). GTP (50 µM) is added to the remaining reaction (4.5 µl) to start the splicing reaction. The reaction is incubated at 37 °C for 1 hour.

Target dependent modulation of the intron-derived optical NASM is followed by  
20 fluorescence measurement as described herein.

**Example 40. Solid-phase nucleic acid sensor array for assays with fluorescent detection—FRET chip**

This example describes a general method for implementing a FRET-based  
25 (fluorescence resonance energy transfer) assay utilizing nucleic acid sensor molecules. (in this case, cGMP-dependent hammerhead nucleic acid sensor molecule) wherein the nucleic acid sensor molecule is immobilized on a solid substrate, *e.g.*, within a microtiter plate well, on a membrane, on a glass or plastic microscope slide, etc.

In the specific embodiment described here, a first oligonucleotide of the nucleotide  
30 sensor molecule is 3'-labeled with an acceptor or quencher fluorophore, such as TAMRA, AlexaFluor 568, or DABCYL, via specific periodate oxidation (see Example 34. for specific protocol). A second oligonucleotide of the nucleic acid sensor molecule, complementary to at least part of the first oligo portion of the NASM, is labeled with a 3' biotin and a 5' donor fluorophore, such as fluorescein (FAM, FITC, etc.). These two nucleic oligonucleotides are

heat-denatured in solution and allowed to anneal/hybridize during cooling to room temperature. After hybridization, the NASM solution is applied to a surface which has been coated with some type of avidin (streptavidin, neutravidin, avidin, etc.). This surface could include a microtiter plate well, a streptavidin-impregnated membrane, a glass or plastic  
5 microscope slide, etc. In any case, the ribozyme-oligo complex is specifically immobilized via the 3' biotin on the donor oligo, leaving the binding domain free to interact with the target effector molecule.

Figure 62 shows the solution-phase FRET construct (upper panel) from which the solid-phase nucleic acid sensor molecule construct (lower panel) was derived. In the figure,  
10 the species labeled 'A' and 'D' represent the acceptor and donor fluorophores, respectively; similarly, the species labeled 'F' and 'Q' represent the donor and quencher fluorophores, respectively.

As shown in Figure 62 (lower panel), the donor and acceptor fluorophores form an efficient FRET-pair; that is, upon excitation of the donor fluorophore near its spectral  
15 absorption maxima, the incident electromagnetic energy is efficiently transferred (nonradiatively) via resonant electric dipole coupling from the donor fluorophore to the acceptor fluorophore. The efficiency of this resonant energy transfer is strongly dependent on the separation between the donor and acceptor fluorophores, the transfer rate being proportional to  $1/R^6$ , where R is the intermolecular separation. Therefore, when the donor  
20 and acceptor are in close proximity, *i.e.*, a few bond-lengths or roughly 10-50 Angstroms, the fluorescent emission from donor species will be reduced relative to its output in an isolated configuration, while the emission from the acceptor species, through indirect excitation by the donor, will be detectable. Upon separation of the donor and acceptor, the donor fluorescence emission signal will increase strongly, while the acceptor emission  
25 signal will show a commensurate decrease in intensity. These effects are shown in Figure 63 (upper panel) for the cGMP-dependent NASM system (SEQ ID: NO. 102).

The black arrow shown in the uppermost panel of Figure 63 indicates the site of phosphodiester bond cleavage, which occurs upon modulation of the target modulation domain (TMD) by a cGMP, to the TMD. The melting temperature,  $T_m$ , of the remaining 5-  
30 base pair sequence (GGUCU in the figure) is  $\sim 25^\circ\text{C}$ . Thus, after effector-mediated cleavage at room temperature, the 5-base pair cleavage fragment will rapidly dissociate from the ribozyme body and diffuse away into solution, as shown in the upper panel of Figure 63.

This target-activated nucleic acid sensor molecule system constitutes a highly sensitive real-time sensor for detecting and quantitating the concentration of the target molecule present in an unknown sample solution. The ultimate limit of detection (LOD) for this system is determined by the switch factor, defined as the ratio of the catalytic rate (in this example, the rate of cleavage) of the ribozyme sensor in the presence of its target to that of the ribozyme in the absence of its target. The dynamic range of the ribozyme sensor will be determined by the dissociation constant,  $K_d$ , for the interaction of the ribozyme binding domain with the target molecule. In theory, the effective dynamic range over which the rate-response of the NASM is linear in the target concentration has  $K_d$  as an upper bound. The lower panel of Figure 63 shows experimental data from the surface-immobilized cGMP-activated NASM. The data shown in the figure represents the FRET signal from the donor fluorophore with the sensor exposed to concentrations of 0, 100  $\mu\text{M}$ , and 600  $\mu\text{M}$  cGMP. The upper panel of Figure 64 shows both donor and acceptor fluorescence signals for the FRET system in the presence of 200  $\mu\text{M}$  cGMP. Note that the experimental data exhibits the behavior expected, as shown in the lower panel of Figure 63. The lower panel of Figure 64 shows the donor signal from the plot in the upper panel fitted to a pseudo-first order rate equation. As shown by the closeness of the data fit, the kinetic response of this sensor system closely approximates a pseudo-first order reaction.

The measured dissociation constant for this cGMP-activated NASM with cGMP is approximately 200-500  $\mu\text{M}$ . In practice, concentration measurements up to 1 mM are possible with this sensor in solution-phase measurements. The absolute precision of measurements made with this NASM will depend on the amount of background catalytic activity (*i.e.*, in the absence of target) and baseline drift of the fluorescence signals from both sample and controls due to physical factors, such as liquid handling errors, reagent adhesion, evaporation, or mixing. After some optimization, run-to-run CVs of a few percent are possible with similar FRET-based NASMs measured in solution. Immobilization of the NASM does not degrade its catalytic activity, although it may limit the effective availability of the target-binding domain for interaction with target molecules. The locally high concentration of surface-immobilized NASM will tend to offset this effect by driving the equilibrium for the association (and subsequent catalytic) reactions toward formation of ribozyme-target complex. The net result is a reduction in the observed catalytic rate of approximately 4-fold at  $K_d$  (200-500  $\mu\text{M}$ ) for the surface-immobilized cGMP-dependent hammerhead NASM, relative to the observed rate for the same sensor system in solution. This effect is shown in the experimental data presented in Figure 65A, which shows linear

(Panel A) and logarithmic (Panel B) plots of the observed catalytic rate constant versus cGMP concentration for the cGMP-dependent hammerhead NASM in both solution and surface-immobilized ('solid') configurations. The upper panel plots the observed rate as a function of target (cGMP) concentration on linear axes, while the lower plot plots the same rate data on log-log axes. The observed rate constant (at a given concentration) for the immobilized solid-phase sensor is roughly 10-fold lower than that for the solution-phase sensor. The practical effect of this is that, while the solution-phase sensor's linear dynamic range is limited to about 0-600  $\mu\text{M}$ , the solid-phase sensor's linear dynamic range is at least 2 mM. Figures 65C, 65D, and 65E shows how this general NASM strategy could be extended to accomplish multiplexed detection of multiple analytes in a sample simultaneously, by using NASMs labeled with fluorophores having different emission wavelengths. Squares are NASMs immobilized on a solid. Circles are NASMs in solution. In this case, the sensors were immobilized in neutravidin-coated microtiter plate wells (96 well plate, volume of 50  $\mu\text{L}$ ). Detection of the fluorescent signals was accomplished in this case by a microplate fluorescence reader equipped with the appropriate lamps, optics, filters, and optical detectors (PMT) manufactured by Packard Instrument Co.

This cGMP-dependent hammerhead NASM system was immobilized on streptavidin-impregnated membranes, and target-activated FRET activity observed. The generalization of this application of surface-immobilized NASM with FRET detection to a micro- or macro-arrayed format on an extended substrate such as glass or plastic is easily envisioned. Such a sensor array could be used to detect and quantify the presence of an arbitrary target molecule in a complex solution, *e.g.*, crude cell extract or biological fluid, in real time. In addition, this general NASM strategy could be extended to accomplish multiplexed detection of multiple analytes in a sample simultaneously, by using NASMs labeled with fluorophores having different emission wavelengths. Experimental data for multiplexed detection using solution-phase cGMP and cAMP FRET sensors is shown in Figure 66B. This NASM, and further extensions of it to include large numbers of unique analyte-sensors, could be used for high throughput screening (HTS) in drug discovery or clonal analysis. In all of these scenarios, optical detection of the FRET signals could be accomplished using a commercially available microarray imager or scanning fluorescence microscope.

The following example contains a sample protocol for the complete process of performing the solid-phase FRET assay described above, including reagent preparation,

DNA amplification, RNA transcription, RNA labeling, substrate and instrument set-up, and, finally, assay data acquisition.

**Example 41. Protocol for micotiter plate-based solid-phase FRET assay:**

5 The nucleic acid sensor molecules used in the assays described in Example 33. for the case of the cGMP NASM system were prepared as follows:

The appropriate sequence construct was synthesized and amplified via PCR. This was accomplished using the appropriate DNA template molecule(s) and the corresponding primers:

10 cGMP PCR template: MK.08.92.A (SEQ ID NO:333)  
 5'GCTTGCAAGCCCTTAGACCCTGATGAGCCTTGCGATGCAAAAAGGTGC  
 TGACGACACAT CGAAACGGTGAAAGCCGTAGGTCT  
 Primer 1: MK.08.66.B (SEQ ID:334)  
 5' AGACCTACGGCTTTCACCGTTTCG  
 15 Primer 2: MK.08.130.B (SEQ ID:335)  
 5'ATACGACTCACTATAGGATGTCCAGTCGCTTGCAATGCCCTTTTAGACC  
 CTGATGAG

20 A trace amount (~ 100 ng) of template dsDNA was combined with 10-100 picomoles of each primer, e.g., 1 uL each of 100 uM stocks. Using taq DNA polymerase, run 15 cycles of PCR using a typical (95-55-75 °C) temperature-step profile was run. The PCR products were cleaned up using a desalting/size separation column, such as a Qiagen Qiaquick PCR cleanup kit. The resuspended volume of cleaned up PCR products was ~ 35 uL. The products can be further concentrated, or added directly to a transcription reaction.

25 After synthesis and amplification of the appropriate sensor construct via PCR, the dsDNA was converted to RNA via *in vitro* transcription. This was typically accomplished using T7 RNA polymerase and pre-mixed NTPs and buffer, such as are provided in the Epicentre High Yield Transcription Kit. The transcription reaction, consisting of ~ 16-20 uL of cleaned up PCR products in a 40-50 uL total reaction volume, was typically run at 37 °C  
 30 for ~ 8-12 hrs. Following the transcription reaction, the products were usually purified via PAGE or using a desalting/size separation column, such as a Centriscp column (Princeton Separations).

Following production and purification of the RNA NASMs, the fluorescent acceptor labels were attached to the sensor molecules. The fluorescent donor label (in this case a



FAM label) was attached to the 5'-biotinylated capture oligo during solid-phase synthesis on a commercial DNA synthesizer, via standard phosphoramidite chemistry. The labeling procedure was accomplished by a specific oxidation of the 3' terminus of the RNA using 20 mM NaIO<sub>4</sub>, followed by reaction with a hydrazide-conjugated fluorescent label. The  
5 resulting covalent bond is extremely stable under the assay conditions. For the oxidation reaction, one typically 15 µL RNA (~ 200-400 pmoles in reaction), 15 µL 0.3 M NaOAc pH 5.4, and 30 µL 20 mM stock NaIO<sub>4</sub> in a were combined 1.5 mL reaction tube, which was then placed on ice in the dark for ~ 1 hr. After the oxidation reaction, the reaction products were run through a Centriscp desalting column. Alternatively the sample can be ethanol  
10 precipitated and resuspended in the original volume.

The oxidized RNA uses then reacted with the dye-(in this example, AlexaFluor 568)-hydrazide conjugate. A typical reaction includes 60 µL oxidized RNA (resuspended in H<sub>2</sub>O), 60 µL 0.3 M NaOAc pH 5.4, and 10 µL 20 mM stock AlexaFluor 568 hydrazide (made fresh in DMSO). This 130 uL reaction was incubated at room temperature for 2 hrs.  
15 After the labeling reaction, the sample was ethanol precipitated, resuspended, gel purified (8% PAGE), and quantitated to determine concentration (using the absorbance at 260 nm). Finally, a 1 uM stock solution of the labeled NASM is prepared in water or buffer.

The substrate and plate-reader/detection system were then prepared for the assay reaction. A typical substrate was a 96 well Pierce Black Neutravidin Plate and a typical  
20 plate reader was the Packard Fusion microplate reader (Packard/Perkin-Elmer). Typical settings for this instrument and this assay (as entered in the user interface menus in the instrument control software) are as follows:

- 2-channel fluorescence assay (ex:FAM\*/em:FAM & TAMRA\*\*)
- 25 - emission detection set to top-read
- integration time/well: 1 sec (1 read/well)
- 25 serial reads of each well, 1 min delay between reads
- lamp intensity: 10/20
- PMT voltage: 950/1100 V
- 30 - PMT gain: 1.0/200
- \*FAM excitation/emission filters: 485 nm/535 nm
- \*\*TAMRA/ or AlexaFluor 568 emission filters: 590 nm

The following reaction mixture of NASM + biotinylated capture oligo was prepared in a 1.5 mL tube:

	Reagent	Stock conc.	Final conc.
5	1 ul FAM-biotin Probe	5uM	500 nM
	1.2 ul Alexa 568 cGMP RNA	5uM	600 nM
	7.8 uL dH2O		
	10.0 uL (total volume = 10 uL x # of wells)		

- 10 The mixture was heat annealed to ~ 85 °C in 1X annealing buffer (30mM Tris, 50mM NaCl), and then cooled to room temperature. The assay reactions were run in the presence/absence of target.

To run a solid-phase FRET assay in a standard 96-well microplate, coated with neutravidin, the following procedure was used:

- 15 First, neutravidin plate surfaces were pre-blocked to prevent nonspecific adsorption of sensor molecules:

-pipet 30 ul of PBS preblocking buffer (0.05% tRNA + 0.025% Tween-20/1X PBS) into wells

-cover and shake for 15 min at room temperature on thermomixer

- 20 -rinse wells with PBS from squirt bottle

-invert plate and tap wells dry on absorbent surface

Next assay reagents were added to each well with multichannel pipet to pre-immobilized sensor molecules in plate wells:

- 25 -pipet 10 ul of annealed RNA+biotinylated probe into wells

-pipet 30 ul of PBS preblocking buffer (0.05% tRNA + 0.025% Tween-20/1X PBS) into wells & -mix by serial pipetting ~ 4x

-cover and shake for 15 min at room temperature on thermomixer

-rinse wells with PBS from squirt bottle

- 30 -invert plate and tap wells dry on absorbent surface

Next assay reaction buffers were added:

-1X HH buffer: 50 mM Tris pH 7.5, 20 mM MgCl<sub>2</sub>, 0.1 mM EDTA

Finally, assay reactions were initiated and the plate was loaded into reader/detector:

-start all reactions simultaneously by adding 50 ul of cGMP in 1X HH Buffer

-shake on thermomixer for ~ 3 sec & load directly into microplate reader

- 5            Kinetic assay data is assigned as (target-dependent) cleavage reaction proceeds. In the case of the Packard Fusion reader described above, data points were plotted on-screen as they were acquired, giving a real-time readout as the assay proceeds.

10        **Example 42. Real-time solid-phase nucleic acid sensor array for assays with surface plasmon resonance (SPR) detection—SPR chip:**

- Surface plasmon resonance (SPR) can be used to detect target-activated ligation/cleavage events of nucleic acid sensor molecules. In SPR, a light beam is directed, via a prism or grating, onto the back surface of a thin metallic layer (typically ~ 50-100 nm of gold) that has been deposited onto a glass or plastic (dielectric) substrate. The incident beam is s-polarized and directed at an angle of incidence which satisfies the condition for total internal reflection (TIR) at the metal/dielectric interface. At certain angles near the TIR critical angle, electromagnetic energy is coupled from the incident beam into surface plasmon excitation modes in the metallic layer. For a fixed excitation wavelength, the precise angular position of the primary surface plasmon resonance depends very strongly on the index of refraction of the medium (typically a liquid) in contact with the front surface of the metallic layer. By monitoring the intensity of the TIR-reflected incident beam as a function of angle, the angular position of the primary surface plasmon resonance can be found. The adsorption or desorption of molecular species from the front surface of the metallic layer (the SPR sensor surface) will modify the refractive index of the medium in its immediate vicinity (*i.e.*, within ~100 nm); this will result in a measurable angular shift in the position of the surface plasmon resonance. SPR is thus a sensitive technique for detecting the binding (including hybridization, etc.) and/or release of molecular species from a surface. Measurements can be performed in real time, allowing the kinetic characterization of binding events, as shown in Figure 67 panels C + D. A system utilizing a large collimated incident beam and an area detector (such as a CCD detector) could be used to generate SPR images of two-dimensional arrays of surface-immobilized species, such as nucleic acid sensor molecules.
- 15  
20  
25  
30

The demonstration of an SPR assay format utilizing nucleic acid sensor molecules was accomplished by coupling hammerhead (HH) nucleic acid sensor molecules (Figure 67, Panel B) to the native gold surface of a commercially available integrated single-channel (*i.e.*, non-imaging) SPR module, called SPReeta (see Figure 67, Panel A). The SPReeta modules are manufactured by Texas Instruments, Inc., and marketed by Nomadics, Inc. (Stillwater, OK). The SPReeta is an integrated package comprised of an optical plastic housing, within which is embedded a light emitting diode (LED) with a polarizer, two reflecting surfaces, *i.e.*, the SPR surface and a mirror, and a linear diode array detector, where the mirror directs the TIR-reflected beam onto diode array. The entire package is built onto a printed circuit board (PCB), with standard dual inline pins which allow the sensor module to be plugged directly into another PCB containing preamplifier and signal processing circuits. The real-time processed data is sent to a computer (PC) via a standard serial port connector. A software application logs, analyzes, and displays the data to the user.

Numerous immobilization schemes for nucleic acid sensor molecules are possible on the gold SPR surface substrate: (1) direct attachment to the gold SPR surface of a construct with a closed stem I, and extended 3' and 5' termini on an open stem III, via a terminal 5' gamma-thiol GTP incorporated during transcription (see Figure 68); (2) direct attachment to a passively adsorbed neutravidin-on-gold layer on the SPR surface using constructs with a truncated, open stem I, and a closed stem III, post-transcriptionally modified by specific periodate oxidation of the 3' terminus, followed by subsequent reaction with biotin-XX-hydrazide to produce a 3' biotinylated hammerhead nucleic acid sensor molecule (see Figure 69); (3) indirect attachment to the gold SPR surface via hybridization of a nucleic acid sensor molecule to a thiol-modified capture oligo via a 3' terminal extension on an open stem I, with stem III closed (see Figure 70). The latter two strategies are flexible in terms of application, allowing easy thermal or chemical stripping and recharging of the sensor surface with nucleic acid sensor molecules *in situ*, whereas direct attachment of the nucleic acid sensor molecules via incorporated thiol-groups would limit the sensor surface to a single use, or require the use of extremely harsh stripping agents.

All of the above nucleic acid sensor molecule constructs were synthesized and/or amplified from existing PCR products. Synthetic DNA templates were prepared by standard solid-phase methods and were purified by denaturing (8 M urea) polyacrylamide gel electrophoresis (PAGE). RNAs were generated by *in vitro* transcription of the appropriate DNA templates that were generated by PCR.

Standard *in vitro* transcription reactions were conducted in a total volume of 50  $\mu$ l containing ~20 pmol template DNA, 50 mM Tris-HCl (pH 7.5 at 23 °C), 10 mM  $MgCl_2$ , 50 mM dithiothreitol (DTT), 20 mM spermidine, 2 mM each of the four ribonucleoside 5'-triphosphates (NTPs), and 35 units/ $\mu$ l T7 RNA polymerase (T7 RNAP) by incubation at 37°C for 2 h. 5'-thiol-modified RNA was prepared by *in vitro* transcription wherein GTP was replaced by guanosine 5'-O-(3-thiotriphosphate) (Sigma Chemical Co., St. Louis, MO). 3'-biotinylated RNA was prepared by post-transcriptionally modification by specific periodate oxidation of the 3' terminus, followed by subsequent reaction with biotin-XX-hydrazide. The resulting RNA products were purified by denaturing 10% PAGE and isolated from the gel by elution with 10 mM Tris-HCl (pH 7.5 at 23 °C), 200 mM NaCl, and 1 mM ethylenediamine tetraacetic acid (EDTA). The recovered RNA was precipitated with ethanol, resuspended in deionized water ( $dH_2O$ ), and stored at -20 °C until use. In addition, the thiol-modified DNA capture oligo was synthesized and purified.

Initial surface immobilization experiments using a kinased (5' 32P-labeled) sample of the thiol-modified DNA capture oligo were conducted on Au/Cr coated microscope slides in order to optimize the thiol-gold coupling procedure. Initial results indicated that blocking of the surface with mercaptohexanol (MCH) and control of pH during the surface coupling reaction were critical. Good surface coupling and binding capacities were observed.

The passive adsorption of neutravidin to the gold sensor surface (via cysteine residues) was investigated, prior to subsequent attachment of a biotinylated hammerhead nucleic acid sensor molecule. Purified neutravidin (Pierce, Rockford, IL) was acquired and flowed over the SPReeta sensor surface at a concentration of 50-100  $\mu$ g/mL. The adsorption of the neutravidin was monitored in real time. cGMP and cAMP hammerhead nucleic acid sensor molecules were generated, amplified, and transcribed with a truncated, open stem I, and a closed stem III. The constructs were then post-transcriptionally modified by specific periodate oxidation of the 3' terminus, followed by subsequent reaction with biotin-XX-hydrazide. These constructs allowed for a maximal cleavage fragment (~ 65 nt=21.5 kD), and thus maximal dynamic range in the observed SPR signals, with only 5 nt retained on the surface. As with the capture oligo and direct-attachment thiol constructs, the melting temperature,  $T_m$ , of the base-paired region retaining the ribozyme body after cleavage is only ~ 25 °C, facilitating rapid dissociation and removal from the SPR sensor surface upon effector-induced cleavage.

After characterizing the SPR signature and kinetic parameters for neutravidin adsorption, the surface immobilization of 3'-biotinylated cGMP nucleic acid sensor molecules (see Figure 71, Panel A). Initial experiments showed a significant reduction in RIUs upon introduction of excess (1 mM) effector solution was introduced into the flow cell (see Figure 71, Panel B). For a 65 nt mass change, the maximum expected signal change (*i.e.*, for complete cleavage of all surface-immobilized nucleic acid sensor molecules) was ~ 2145 RIU; the observed signal change was ~ 1487 RIU, with a signal to noise ratio (SNR) > 100. Subsequent negative control experiments (using cAMP) confirmed the effector-dependent origin of the observed signal. The cleavage course was fitted to a pseudo-first order rate function (correlation coefficient = 0.9977), with an observed rate constant of ~ 1 per minute.

The usual scheme for attachment of the nucleic acid sensor molecule to the capture oligo and subsequent target - induced activity is shown in Panel C of Figure 71. In order to verify the generality of the surface attachment strategy, as well as the catalytic activity of surface-immobilized nucleic acid sensor molecules, the 3'-biotinylated cGMP-dependent hammerhead was immobilized via neutravidin to a SPREETA sensor chip surface. The cGMP HH Riboreporter surface was then exposed to a range of concentrations of cGMP (0-1 mM) and varying  $Mg^{2+}$  concentration in the ambient buffer (2-20 mM). Target-modulated cleavage was observed for  $Mg^{2+}$  concentrations > 10 mM, and cGMP concentrations greater than ~ 10  $\mu$ M. The time-course for a portion of this type of experiment is shown in Figure 72. This data was generated using a single surface-adsorption step with neutravidin, followed by sequential addition of increasing concentrations of cGMP, with buffer flushes and/or active buffer pumping (peristaltic) in between each cGMP addition. All sample fluids were injected at volumes of ~ 25-40  $\mu$ L, while buffer washes were ~ 100  $\mu$ L of 1X buffer. For active pumping, volume exchange was ~ 500  $\mu$ L of 1X buffer.

**Example 43. Solid-phase nucleic acid sensor array for assays with fluorescent detection—ligase chip**

Nucleic acid sensors can be tethered to solid supports while still maintaining their activity, as shown in Figure 66. The plot in Panel B of Figure 66 shows the fraction of this endonuclease-based nucleic acid sensor molecule that is released by self-cleavage over time in presence of its analyte. When exposed to its target analyte, the solid-phase nucleic acid sensor molecule increased in activity. In the case of ligase based NASMs, the sensor molecule ligated itself to a labeled oligonucleotide in the presence of target, and the

detected signal increase. In the case of the endonucleolytic nucleic acid sensor, the molecule cleave a labeled oligonucleotide from itself, resulting in a decrease in the detected signal.

These ligase based nucleic acid sensor molecules can be used in solid-state

5 scintillation proximity assays, membrane assays, chip-based assays with fluorescent detection chip based assays with electrochemical detection, and in assays using intrinsic (non labeled detection such as, surface plasmon resonance. Figure 74 presents the two principal solid-phase array (chip) formats used for ligase-based nucleic acid sensors. (a) In the capture format, the ligation reaction in the presence of target is performed *in vitro*,

10 followed by an optional amplification step, hybridization of the reaction products (amplified or not) to a capture oligo array, and finally washing and detection/readout of the array. Here, each capture tag oligonucleotide is a unique sequence address, allowing spatial multiplexing of the assay over numerous different NASM/target combinations. (b) In the *in situ* format, the ligase sensors are pre-immobilized on the chip via unique capture tags; the

15 ligation reaction in the presence of target is performed *in situ* on the chip surface, followed by washing and detection/readout of the array. Under ordinary solution-phase reaction conditions for ligase-based nucleic acid sensors, a external effector oligo (the non-immobilized analog of the capture oligo used to immobilize the sensor to a surface) is typically present. In order for the capture chip assay format (*i.e.*, *in vitro* reaction followed

20 by on-chip capture and readout, as shown in Figure 74A) to be a viable the case where no isothermal amplification step is performed prior to hybridization, it is necessary either to (a) exclude such a effector oligo from the *in vitro* reaction, so that the appropriate portion of the sensor sequence would be free to hybridize to the surface-immobilized capture oligo in the subsequent capture step; or (b) to include some form of effector oligonucleotide that could

25 be easily separated or competitively displaced from the sensor molecule after the *in vitro* reaction, allowing hybridization to the surface-immobilized capture oligo in the subsequent capture step. Both strategies were shown to be viable. In particular, Figure 73 shows an example, of an ERK-dependent ligase-based nucleic acid sensor system ligase-based NASMs can function effectively in solution-based assays without the presence of a effector

30 oligonucleotide. The ligation signals from 2 separate reactions that were run in the presence of target (ERK) and the presence and absence of an external effector oligonucleotide, respectively; showed *i.e.*, that the reaction products amplified equally in the same number of PCR cycles, that the sensor functions equally well in the presence or absence of a capture effector oligonucleotides. A parallel result is shown for the case of the reactions run in the

absence of target (ERK) and the presence and absence of an external passivating capture oligo, respectively. The results, which have been shown to be valid for all ligase-based sensor systems studied (> 10 different ligase sensor systems), is that the ligase-based sensors function in solution-based reactions equally well in the presence or absence of a  
5 effector oligo. The *in vitro* reaction steps for all capture chip data shown and referred to in the following examples were all conducted in the absence of an effector oligo.

For chip-based assays, the activity of the nucleic acid sensor molecules can be detected using many methods shown schematically in Figure 14. In one embodiment of the invention, the nucleic acid sensor molecule has a self cleaving activity. These nucleic acid  
10 sensors can be attached to a solid substrate in such a way that cleavage results in the release into solution of a nucleic acid fragment of known size. This fragment can then be detected through the use of agarose gels using stains that bind to nucleic acids, including ethidium bromide. The activity of the nucleic acid sensor molecule with self-cleaving activity could also be detected through attaching a fluorophore and a quencher group to either side of the  
15 cleavage site, as detailed above, and in Figure 62. In another embodiment of the invention, the nucleic acid sensor molecule has ligation activity. The components of a ligase-based nucleic acid sensor molecule immobilized to a chip surface are shown in Figure 73. The activity of this sensor in the presence of target molecules can be detected through fluorescence, chemiluminescence, radioactivity, or electrochemical means. The labeling of  
20 the sensor can be accomplished either during the assay reaction, during a subsequent amplification reaction (e.g., via RT), or after the sensor (as a component of the assay reaction products) has been captured to a microarray chip for read out, as shown in Figure 74.

Figure 75 shows a multiplex *in situ* ligase sensor chip, with pre-immobilized  
25 radiolabeled sensors activatable by lysozyme (LYS) and FMN. The left panel shows the pre-hybridized sensor-substrate complexes, in this case immobilized via a biotin linker on the substrate oligo on a streptavidin/glass chip. The right panel shows the array after it has been exposed to target mixtures consisting of lysozyme only, FMN only, and lysozyme plus FMN, as indicated in the figure. Following the *in situ* ligation reaction, the chip was  
30 subjected to a denaturing wash, in order to remove any sensor molecules not ligated to their substrates. Note the retention of labeled sensors only in spots that had been exposed to the corresponding target, and that both protein (lysozyme) and small molecule (FMN) sensors function in the presence of nontarget analytes.



Figure 13 shows a solid-phase nucleic acid sensor molecule array on a streptavidin membrane. In this case, 6 different ligase sensors were immobilized on a streptavidin membrane, and assayed in an *in situ* format, with each NAMS being reacted with an analyte mixture containing all 6 targets. The plot shows the detected retention of ligation signal following target exposure and washing. Here detection was accomplished using a  
5 radiolabeled sensor, with the ligase immobilized via a biotin tag on the 5' terminus of the substrate oligo

Figure 76 shows dose-response data for ERK-dependent ligase-based nucleic acid sensor molecules. The data plotted in the left panel was generated from a PAGE-based  
10 assay, while the data in the right panel was generated from a capture chip assay (right panel). The reaction incubation time for both assays was approximately 2 hrs. Detection in both cases was via radiolabeled sensors and substrate oligonucleotides. Note that the capture chip assay recapitulates the apparent affinity constant ( $K_d$ ) of  $\sim 1 \mu\text{M}$  measured from the gel-assay data, indicating that the chip mode of capture and detection accurately reflects all  
15 reaction parameters observed in the *in vitro* case.

Figure 77 shows dose-response data for an *in situ* ligase-based nucleic acid sensor molecule array populated with ERK-dependent unlabeled ligase sensors. The left panel shows an image of the *in situ* chip after exposure to target (ERK) plus radiolabeled substrate and subsequent washing to remove unligated substrate. The reaction incubation time for  
20 both assays was approximately 2 hrs. Each array spot corresponds to a different concentration of target, giving rise to the observed pattern of positive/negative controls (10/0  $\mu\text{M}$ , respectively), and the concentration profile (0-10  $\mu\text{M}$ ) shown. The retained ligation signal from each spot in the concentration profile is plotted vs. its corresponding target concentration in the right panel. The linear dose-response observed here is a result of  
25 two effects: (1) the surface-immobilization of the NASM reduces its effective catalytic rate over the rate observed *in vitro*; and (2) the much higher surface to volume (S/V) ratio in the *in situ* chip format, relative to the *in vitro* reaction/capture chip format, has been shown to promote nonspecific interactions of protein targets (*e.g.*, ERK) with the chip surface coating. This leads to a reduced effective target concentration in the reaction volume at each  
30 sensor spot.

Figure 78 shows the dose-response data for a ERK-dependent ligase-based nucleic acid sensor molecule capture array. The NASMs in this case were fluorescently 3' labeled with Cy3-labeled ligase sensors, and the substrate oligos were 5' labeled with biotin. After 2 hr. *in vitro* reactions in the presence of varying concentrations of target (ERK), the reaction

products were hybridized to the capture array. After hybridization and washing, tyramide signal amplification (TSA) was performed using a streptavidin-conjugated horseradish peroxidase (HRP) pulse Cy5-labeled tryramide HRP substrates. The captured and amplified fluorescent ligation signal for each spot is plotted in the right panel vs. its corresponding target concentration. This data demonstrates the high sensitivity of fluorescence detection for ligase-based nucleic acid sensor molecule arrays in both *in situ* and capture formats.

Figure 79 shows the components of a generalized construct for an amplifiable ligase-based nucleic acid sensor molecule. Each sensor in an array would have a unique 3' capture tag sequence domain, and a common RT primer binding site. The external substrate oligonucleotide contains a signal generation label, which may comprise a direct label (*e.g.*, fluorescent, radionuclide, enzyme-linked, etc.) or an affinity tag (*e.g.*, biotin) for subsequent indirect labeling or amplification (*e.g.*, via TSA), or a specific sequence (*e.g.*, the T7 RNA polymerase promoter sequence) which will facilitate selective amplification of ligated sensors via RT or RT-PCR. The amplification process (RT or RT-PCR) can also be used to incorporate multiple direct or indirect labels or affinity tags, thus increasing the sensitivity of the sensor system. In one embodiment, detection can be accomplished by using a chimeric (RNA/DNA) external substrate oligonucleotide which contains a DNA sequence that is necessary for subsequent amplification of the nucleic acid, such as the T7 RNA polymerase promoter sequence, as shown in Figure 79. An arbitrary number (*e.g.*, 10, as shown) of unique analyte-specific sensors are each outfitted (*e.g.*, via PCR) with an arbitrary number (*e.g.*, 10, as shown) of unique sequence capture tags. In the present example, the result would be a panel of 100 uniquely addressed sensors. Thus, ten different targets could be simultaneously assayed under ten different conditions, with the reaction products RT-amplified (and simultaneously labeled with multiple fluorophores per sensor molecule) and spatially sorted by hybridization to a capture array chip, and finally detected using standard techniques, such as fluorescence. Upon ligation of such an external substrate, RT can be performed to synthesize a complementary strand to the sensor, as shown in Figure 81. A given NASM is reacted *in vitro* with its corresponding target analyte and a chimeric (*i.e.*, containing both DNA and RNA bases) substrate oligo containing the T7 promoter sequence comprised of DNA bases. Following the ligation reaction, an RT step is performed to synthesize a complementary cDNA strand to the RNA sensor molecule. The NASM is then hydrolyzed or degraded with RNaseH. In the case of NASM that ligated its substrate in the presence of target, this leaves the cDNA strand with the DNA T7 promoter sequence annealed. This complex forms a suitable platform for the T7 RNA polymerase to

transcribe multiple copies of those cDNA sequences corresponding to sensors which were triggered by their respective targets. During transcription, NTPs with direct fluorescent labels or affinity tags (e.g., biotin) are multiply incorporated into the transcribed cRNA molecules. After hybridization to the capture array, the amplified cRNA sensors can be

5 detected via their direct labels, or by subsequent staining with a conjugate to the incorporated affinity tags, e.g., streptavidin-phycoerythrin. After hydrolysis of the original RNA sensor strand, the double-stranded DNA portion containing the T7 promoter sequence will allow *in vitro* transcription of the cDNA to cRNA by standard methods. This transcription step will amplify the sensor sequence, and can be used to simultaneously

10 incorporate nucleotides that contain direct or indirect (e.g., biotin, or other affinity tag) labels for subsequent fluorescence- or luminescence-based detection, as shown in Figure 80. This detection strategy can be implemented in a multiplexed format by using PCR to incorporate arbitrary sequence capture oligo domains, as well as common RT primer binding domains, to a panel of ligase-based sensors. In particular, Figure 80 shows a

15 generalized strategy for performing a multiplexed capture chip formatted assay with ligase-based nucleic acid sensors. An arbitrary number (e.g., 10, as shown) of unique analyte-specific sensors are each outfitted (e.g., via PCR) with an arbitrary number (e.g., 10, as shown) of unique sequence capture tags. In the present example, the result would be a panel of 100 uniquely addressed sensors. Thus, 10 different targets could be simultaneously

20 assayed under 10 different conditions, with the reaction products RT-amplified (and simultaneously labeled with multiple fluorophores per sensor molecule) and spatially sorted by hybridization to a capture array chip, and finally detected using standard techniques, such as fluorescence. In one embodiment, shown in Figure 80, a generalized strategy for performing a highly sensitive capture chip formatted assay with ligase-based nucleic acid

25 sensors. A given sensor molecule is reacted *in vitro* with its corresponding target analyte and a chimeric (RNA) substrate oligo containing the T7 promoter sequence comprised of DNA bases. Following the ligation reaction, an RT step is performed to synthesize a complementary cDNA strand to the RNA sensor molecule. The RNA sensor molecule is then hydrolyzed or degraded with RNaseH. In the case of a sensor which ligated its

30 substrate in the presence of target, this leaves the cDNA strand with the DNA T7 promoter sequence annealed. This complex forms a suitable platform for the T7 RNA polymerase to transcribe multiple copies of those cDNA sequences corresponding to sensors which were triggered by their respective targets. During transcription, NTPs with direct fluorescent labels or affinity tags (e.g., biotin) are multiply incorporated into the transcribed cRNA

molecules. After hybridization to the capture array, the amplified cRNA sensors can be detected via their direct labels, or by subsequent staining with a conjugate to the incorporated affinity tags, *e.g.*, streptavidin-phycoerythrin.

Highly multiplexed *in vitro* or *in situ* assays can then be performed, using the specific hybridization of each sensor's unique capture oligo to address its signal to a feature spot whose position on the array is known. The presence or relative concentration of a given analyte can then be determined by the presence/absence of a signal in the expected array position, and the relative concentration determined by comparison of the relative signal intensity to that of other sensor spots in the array.

Listed below are selected DNA template sequences that can be used to generate ligase-based nucleic acid sensor molecules via PCR, as described above:

**Synthetic DNA universal templates for 2- & 3-piece ligase constructs**  
(underlined regions = primer annealing regions)

15

**ERK.A.LIG.TMPLT (DG.20.58A):** SEQ ID NO: 109

5'CTATAGGACTTCGGCGAAAGCCGTTTCGACCACGCTAAGGAGGATTTC  
GAAAGCGGCTACGGTCCGCCAGTGTCTTAGACAGGAGGTTAGGTGC

**ERK.E.LIG.TMPLT (DG.20.58E):** SEQ ID NO: 111

20

5'-  
CTATAGGACTTCGGCGAAAGCCGTTTCGACCACGCTAAGGAGGATTTCGAAAGC  
GGCTACGGTCCGCCAGCTCTTAGACAGGAGGTTAGGTGC

**LYS.LIG.TMPLT.L1-11.2 :** SEQ ID NO: 336

25 5'CTATAGGACTTCGGCGAAAGCTAACGTCTCATGGCTAAATTGCCATGT  
TGCTACAAATGATATGACTAGAGAGGTTAGGTGC

**FMN.LIG.TMPLT.L1-R7C13:** SEQ ID: 337

5'CTATAGGACTTCGGTCCAGTGCTCGTGCACTAGGCCGTTTCGACCTTCAG  
GATATGCTTCGGCAGAAGGGAACCTTAGACAGGAGGTTAGGTGC3'

**THEO.LIG.TMPLT.L1-D1:** SEQ ID: 338

30 5'CTATAGGACTTCGGTCCAGTGCTCGTGCACTAGGCCGTTTCGACCATGAT  
ACCAGCATCGTCTTGATGCCCTTGGCAGCATCTTAGACAGGAGGTTAGGT  
GC

For 3-piece system:

5'-primer: TK.16.32.A SEQ ID: 339  
 5'-TTCTAATACGACTCACTATAGGACTTC  
 3'-primer (5 nt sub): TK.16.32.B SEQ ID: 340  
 5'-ATTCGAGATGTCCTTGGACCAAAGCCGCACCTAACCTC  
 3'-primer (15 nt sub): TK.16.32.15NT SEQ ID NO: 341  
 5'-ATTCGAGATGTCCTTGGACCAAAGCCTCCATCGTGCGCACCTAACCTC  
 Substrate (5 nt): TK.04.82.A SEQ ID NO: 342  
 5'-Biotin/Fluor-CATGCGACCTTACGATCAGATGACCTugcacu  
 Substrate (15 nt): MK.08.125A SEQ ID NO: 343  
 5'-Biotin-TCCATCGTGCGCACu  
For 2-piece system:  
 5' primer: TK16.32A SEQ ID NO: 344  
 5'-TTCTAATACGACTCACTATAGGACTTC  
 3' primer: MK.08.125B SEQ ID NO: 345  
 5'-TCCATCGTGCGCACCTAACCTC  
 Substrate: MK.08.125A SEQ ID NO: 346  
 5'-Biotin-TCCATCGTGCGCACu

**Example 43. Conversion of nucleic acid sensor molecules into unimolecular ligase NASMs**

The Erk and pERK dependent nucleic acid sensor molecules can be converted to unimolecular ligase NASMs. The engineered NASM ERK B, ERK CW45-33-C08 and ERK CW45-33-D09 were converted by PCR as described for Figure 28 with primers GGACTTCGGCGAAAGC and AGTGCTCTCGCACCTAACCTCCTGTCT (SEQ ID NO:363). NASMs pERK CW45-33-C04, CW45-33-D05 and CW45-33-H03 were all converted with primers GGACTTCGGCGAAAGC (SEQ ID NO:364) and AGTGCTCTCGCACCTAACCTCCTGTCT (SEQ ID NO:365). Circularization can be detected by RTPCR for ERK B (GCTACGGTCCGCCAGTTCTT) (SEQ ID NO:366) and CGCTTTCGGAAATCCTCCTT) (SEQ ID NO:367), ERK CW45-33-C08 (GCTACGGTCCGCCAGGGGCT (SEQ ID NO:368) and CGCTTTCGGAAATCCTCCTT (SEQ ID NO:369)), ERK CW45-33-D09 (GCTACGGTCCGCCAAAAGCT (SEQ ID NO:370) and CGCTTTCGGAAATCCTCCTT (SEQ ID NO:371)) and pERK CW45-33-C04, CW45-33-D05 and CW45-33-H03 (AAGGGGAAAGCGTTATTAAG (SEQ ID NO:372) and TCGAGGAACCAATTCGCTAG (SEQ ID NO:373)).

### References

- 5 Andersson, K. E., H. Gemalmaz, et al. (1999). "The effect of sildenafil on apomorphine-evoked increases in intracavernous pressure in the awake rat." J Urol 161(5): 1707-12.
- Arora, A. S., P. C. de Groen, et al. (1996). "Hepatocellular carcinoma cells resist necrosis during anoxia by preventing phospholipase-mediated calpain activation." J Cell Physiol 167(3): 434-42.
- 10 Asher, et al., PCT 98/08974 (1996)
- Ballard, S. A., C. J. Gingell, et al. (1998). "Effects of sildenafil on the relaxation of human corpus cavernosum tissue *in vitro* and on the activities of cyclic nucleotide phosphodiesterase isozymes." J Urol 159(6): 2164-71.
- 15 Barlocco, D., G. Cignarella, et al. (2001). "Phenylpiperazinylalkylamino substituted pyridazinones as potent alpha(1) adrenoceptor antagonists." J Med Chem 44(15): 2403-10.
- 20 Basararsky, T. et al., "Overview of a Microarray Scanner: Design Essentials for an Integrated Acquisition and Analysis Platform", in *Microarray Biochip Technology*, ed. by M. Schena, Eaton Publishing Co., 2000, Natick, MA, pp. 265-284.
- Beaudet L. et al., "Homogeneous Assays for Single-Nucleotide Polymorphism Typing Using AlphaScreen", Genome Res. 11:600 (2001).
- 25 Bell, et al., J. Biol. Chem. 273: 14309 (1998).
- Berzal-Herranz A; Joseph S; Burke JM. *In vitro* selection of active hairpin ribozymes by sequential RNA-catalyzed cleavage and ligation reactions. Genes and Development, 1992 Jan, 6(1):129-34.
- 30 Berzas Nevado, J. J., J. Rodriguez Flores, et al. (2001). "Micellar electrokinetic capillary chromatography for the determination of Viagra and its metabolite (UK-103,320) in human serum." Electrophoresis 22(10): 2004-9.
- 35 Bianchini, M., M. Radrizzani, et al. (2001). "Specific oligobodies against ERK-2 that recognize both the native and the denatured state of the protein." J Immunol Methods 252(1-2): 191-7.
- 40 Bloom, T. J. and J. A. Beavo (1996). "Identification and tissue-specific expression of PDE7 phosphodiesterase splice variants." Proc Natl Acad Sci U S A 93(24): 14188-92.
- Bock, et al., Nature, 355(6360): 564-6 (1992).
- 45 Bolger, G. B., S. Erdogan, et al. (1997). "Characterization of five different proteins produced by alternatively spliced mRNAs from the human cAMP-specific phosphodiesterase PDE4D gene." Biochem J 328(Pt 2): 539-48.
- 50 Breaker, PCT WO 98/43993 (1998)

- Breaker, PCT WO 98/27104 (1998)
- 5 Bruchez, M. et al., "Semiconductor Nanocrystals as Fluorescent labels", Science 281, 2013 (1998).
- 10 Burt, A. R., M. Sautel, et al. (1998). "Agonist occupation of an alpha2A-adrenoreceptor-Gi1alpha fusion protein results in activation of both receptor-linked and endogenous Gi proteins. Comparisons of their contributions to GTPase activity and signal transduction and analysis of receptor-G protein activation stoichiometry." J Biol Chem 273(17): 10367-75.
- Caceci, et al., Byte 9: 340-362 (1984).
- 15 Chan, V. et al., Effect of Hydrophobicity and Electrostatics on Adsorption and Surface Diffusion of DNA Oligonucleotides at Liquid/Solid Interfaces", J. Colloid and Interface Sci. 203:197 (1998).
- 20 Chan, W.C.W and Nie, S., "Quantum Dot Bioconjugates for Ultrasensitive Nonisotopic Detection", Science 281: 2016 (1998).
- Coleman, D. E. and S. R. Sprang (1998). "Crystal structures of the G protein Gi alpha 1 complexed with GDP and Mg<sup>2+</sup>: a crystallographic titration experiment." Biochemistry 37(41): 14376-85.
- 25 Ciesiolka et al., Methods in Enzymology 267: 3 15-335 (1996).
- Conti, M. (2000). "Phosphodiesterases and cyclic nucleotide signaling in endocrine cells." Mol Endocrinol 14(9): 1317-27.
- 30 Cook, N.D. et al., "SPA: a revolutionary new technique for drug screening", *Pharmaceutical Manufacturing International*, 1992, pp. 49-53.
- Cooper, et al., Biochemistry 29: 926 1-9268 (1990).
- 35 Cotten, et al., Nuc. Acid Res. 19: 2629-2635 (1991).
- Cox JC; Rudolph P; Ellington AD. Automated RNA selection. Biotechnology Progress, 1998 Nov-Dec, 14(6):845-50.
- 40 DeRisi et al., Science 278: 680-686 (1997).
- Ellington and Szostak, Nature 346(6287): 818-822 (1990).
- Fehrentz, J.A. et al., Biochem. Biophys. Res. Commun. 188:865 (1992).
- 45 Froehler, et al., Nucleic Acids Research 14: 53 99-5467 (1986a).
- Froehler, Tet. Lett. 27: 5575-5578 (1986b).

- Franke, B., J. W. Akkerman, et al. (1997). "Rapid  $\text{Ca}^{2+}$ -mediated activation of Rap1 in human platelets." Embo J 16(2): 252-9.
- 5 Gold, et al., Annual Review of Biochemistry 64: 763-797 (1995).
- Gold, et al., Proc. Natl. Acad. Sci. 94: 59-64 (1997).
- Griffiths, et al. EMBO. J 13: 3245-3260 (1994).
- 10 Goueli, S. A., K. Hsiao, et al. (2001). "Assaying activity of individual protein kinases in crude tissue or cellular extracts." Methods Enzymol 333: 16-27.
- Green R; Szostak JW. Selection of a ribozyme that functions as a superior template in a self-copying reaction. Science, 1992 Dec 18, 258(5090):1910-15.
- 15 Guatelli, et al., Proc. Nati. Acad. Sci. 87: 1874 (1990).
- Hamaguchi, N. et al., "Aptamer Beacons for the Direct Detection of Proteins", Anal. Biochem. 294: 126 (2001).
- 20 Han, M. et al., "Quantum-dot-tagged microbeads for multiplexed optical coding of biomolecules", Nature Biotechnol. 19, 631 (2001).
- Heath, R. et al., "cAMP [ $^{125}\text{I}$ ] scintillation proximity assay (SPA) – a homogeneous radioimmunoassay for cAMP", in *Cell Signalling: Experimental Strategies*, ed. Reid, E., Cook, G.M.W., and Luzio, J.R., 1992, pp. 193-194.
- 25 Heller and Morrison, in *Rapid Detection and Identification of Infections Agents*, Academic Press, Inc., San Diego, Calif., pages 245-256 (1985).
- 30 Heller, R.A. et al., "Gene Chips and microarrays: applications in disease profiles, drug target discovery, drug action and toxicity", in *DNA Microarrays: A Practical Approach*, ed. by M. Schena, Oxford University Press, 1999, Oxford, pp. 187-202.
- 35 Hirose, et al., Tet. Lett. 28: 2449 (1978).
- Hobbs, et al., Biochemistry 12: 5138-5145 (1973).
- 40 Hu, B., J. Ellingboe, et al. (2001). "(4-Piperidin-1-yl)phenyl amides: potent and selective human beta(3) agonists." J Med Chem 44(9): 1456-66.
- Jameson, D.M. and Sawyer, W.H., "Fluorescence Anisotropy Applied to Biomolecular Interactions", Methods in Enzymology 246:283 (1995).
- 45 Jameson, D.M. and Seifried, S.E., "Quantificatin of Protein-Protein Interactions Using Fluorescence Polarization", METHODS 19:222 (1999).
- Jellinek, et al., Biochemistry 34: 11363-11372 (1995).



- Jenne A; Hartig JS; Piganeau N; Tauer A; Samarsky DA; Green MR; Davies J; Famulok M. Rapid identification and characterization of hammerhead-ribozyme inhibitors using fluorescence-based technology. Nature Biotechnology, 2001 Jan, 19(1):56-61.
- 5 Jeremy, J. Y., S. A. Ballard, et al. (1997). "Effects of sildenafil, a type-5 cGMP phosphodiesterase inhibitor, and papaverine on cyclic GMP and cyclic AMP levels in the rabbit corpus cavernosum *in vitro*." Br J Urol 79(6): 958-63.
- Jhaveri, et al. J. Am. Chem. Soc. 122: 2469-2473 (2000).
- 10 Jenison, et al. Science, 263(5 152): 1425-9 (1994).
- Jolley, M., "Fluorescence Polarization: an analytical tool for immunoassay and drug discovery", Comb. Chem. High Throughput Screen. 2(4):177 (1999).
- 15 Joyce, in Molecular Biology of RNA: UCLA Symposia on Molecular and Cellular Biology, T. R.Cech (ed.), Liss, N.Y., 1989, pp. 361-371.
- Koizumi, M., G. A. Soukup, et al. (1999). "Allosteric selection of ribozymes that respond to the second messengers cGMP and cAMP." Nat Struct Biol 6(11): 1062-71.
- 20 Koizumi, M., J. N. Kerr, et al. (1999). "Allosteric ribozymes sensitive to the second messengers cAMP and cGMP." Nucleic Acids Symp Ser 42: 275-6.
- 25 Kozasa, T. and A. G. Gilman (1995). "Purification of recombinant G proteins from Sf9 cells by hexahistidine tagging of associated subunits. Characterization of alpha 12 and inhibition of adenylyl cyclase by alpha z." J Biol Chem 270(4): 1734-41.
- Kraus, et al., Journal of Immunology 160: 5209-52 12 (1998).
- 30 Krishnan, et al., Science 282(53 88): 484-7 (1998).
- Kwoh, et al., Proc. Natl. Acad. Sci. 86: 1173 (1989).
- 35 Kubik, et al., Nucleic Acids Res. 22(13): 2619-26 (1994).
- Lakowicz, J.R. and Berndt, K., "Lifetime-selective fluorescence imaging using an rf phase-sensitive camera", Rev. Sci. Instr. 62(7):1727 (1991).
- 40 Lakowicz, J.R., Principles of Fluorescence Spectroscopy, Second Edition, 1999. Kluwer Academic/Plenum Publishers, New York.
- Lee, E., M. E. Linder, et al. (1994). "Expression of G-protein alpha subunits in Escherichia coli." Methods Enzymol 237: 146-64.
- 45 Li, Y., P. M. Sternweis, et al. (1998). "Sites for Galpha binding on the G protein beta subunit overlap with sites for regulation of phospholipase Cbeta and adenylyl cyclase." J Biol Chem 273(26): 16265-72.
- 50 Lin, et al. Nucleic Acids Research 22: 5229-5234 (1994).

- Liu, X. and Tan, W., "A fiberoptic evanescent wave DNA biosensor based on novel molecular beacons", Anal. Chem. 71:5054 (1999).
- 5 Lockhart, et al., Science, 274(5287): 6 10-4 (1996).
- Mace, M.L. et al., "Novel Microarray Printing and Detection Technologies", in *Microarray Biochip Technology*, ed. by M. Schena, Eaton Publishing Co., 2000, Natick, MA, pp. 39-64.
- 10 Morrison, in *Nonisotopic DNA Probe Techniques*, Kricka, ed., Academic Press, Inc., San Diego, Calif., chapter 13 (1992).
- Mansour, S. J., J. M. Candia, et al. (1996). "Interdependent domains controlling the enzymatic activity of mitogen-activated protein kinase kinase 1." Biochemistry 35(48):  
15 15529-36.
- Min, K., N. N. Kim, et al. (2000). "Sildenafil augments pelvic nerve-mediated female genital sexual arousal in the anesthetized rabbit." Int J Impot Res 12 Suppl 3: S32-9.
- 20 Moreland, R. B., I. Goldstein, et al. (1998). "Sildenafil, a novel inhibitor of phosphodiesterase type 5 in human corpus cavernosum smooth muscle cells." Life Sci 62(20): 309-18.
- Morrison, et al., Anal. Biochem 183: 23 1-244 (1989).
- 25 Muller, T., P. Engels, et al. (1996). "Subtypes of the type 4 cAMP phosphodiesterases: structure, regulation and selective inhibition." Trends Pharmacol Sci 17(8): 294-8.
- 30 Nagaoka, T., T. Shirakawa, et al. (1998). "Cyclic nucleotide phosphodiesterase 3 expression *in vivo*: evidence for tissue-specific expression of phosphodiesterase 3A or 3B mRNA and activity in the aorta and adipose tissue of atherosclerosis-prone insulin-resistant rats." Diabetes 47(7): 1135-44.
- 35 Obata, T., M. B. Yaffe, et al. (2000). "Peptide and protein library screening defines optimal substrate motifs for AKT/PKB." J Biol Chem 275(46): 36108-15.
- Oberholte, R., J. Ratzliff, et al. (1997). "Multiple splice variants of phosphodiesterase PDE4C cloned from human lung and testis." Biochim Biophys Acta 1353(3): 287-97.
- 40 O'Shaughnessy, D.J., "Hydrazido-derivatized supports in affinity chromatography", J. Chromatography, 510:13 (1990).
- Owicki, J.C. et al., "Application of Fluorescence Polarization Assays in High-Throughput Screening", Genetic Engineering News 17(19), 1997.
- 45 Pagratis, et al., Nat Biotechnol 15: 68-73 (1997).
- Pieken, et al., Science 253: 314-3 17 (1991).
- 50

- Pillai, R., K. Kytle, et al. (1993). "Use of a yeast expression system for the isolation and analysis of drug-resistant mutants of a mammalian phosphodiesterase." Proc Natl Acad Sci U S A 90(24): 11970-4.
- 5 Potyrailo, et al. Anal Chem, 70(16): 34 19-25 (1998).
- Proudnikov, D. and Mirzabekov, A., "Chemical Methods of DNA and RNA fluorescent labeling", Nucleic Acids Res. 24(22):4535 (1996).
- 10 Richter, W., T. Hermsdorf, et al. (2000). "Refolding, purification, and characterization of human recombinant PDE4A constructs expressed in *Escherichia coli*." Protein Expr Purif 19(3): 375-83.
- Robertson, D. and Davidson, N., "Covalent coupling of ribonucleic acid to agarose",  
15 Biochemistry 11(4):533 (1972).
- Robertson DL; Joyce GF. Selection *in vitro* of an RNA enzyme that specifically cleaves single-stranded DNA. Nature, 1990 Mar 29, 344(6265):467-8.
- 20 Robertson, M.P. and Ellington, A.D., "Design and optimization of effector-activated ribozyme ligases", Nucleic Acids Res. 28(8):1751 (2000).
- Ruhn, P.F. et al., "Development of dihydrazide-activated silica supports for high-performance affinity chromatography", J. Chromatography A 669:9 (1994).
- 25 Saiki, et al., Science 230: 1350-54 (1985).
- Saiki, et al., Science 239: 487-49 1 (1988).
- 30 Sako, Y. et al., "Single-molecule imaging of EGFR signaling on the surface of living cells", Nature Cell Bio. 2:168 (2000).
- Schena, M. and Davis, R.W., "Technology Standards for Microarray Research", in *Microarray Biochip Technology*, ed. by M. Schena, Eaton Publishing Co., 2000, Natick,  
35 MA, pp1-18.
- Schermer, M. J., "Confocal scanning microscopy in microarray detection", in *DNA Microarrays: A Practical Approach*, ed. by M. Schena, Oxford University Press, 1999, Oxford, pp. 17-42.
- 40 Schluep, T. and Cooney, C.L., "Immobilization of oligonucleotides on a large pore support for plasmid purification by triplex affinity interaction", Bioseparation 7:317 (1999).
- Seelig B; Jaschke A. A small catalytic RNA motif with Diels-Alderase activity. Chemistry and Biology, 1999 Mar, 6(3):167-76.
- 45 Seetherman, S., et al., "Immobilized RNA switches for the analysis of complex and biological mixtures", Nature Biotech. 19:336 (2001).
- 50 Serivce, "Coming soon: the pocket DNA sequencer." Science 282, 399-401

- Seiwert, S. D., T. Stines Nahreini, et al. (2000). "RNA aptamers as pathway-specific MAP kinase inhibitors." Chem Biol 7(11): 833-43.
- 5 Simone, et al. Trends Genet 14(7): 272-6 (1998).
- Singh, K.K. et al., "Rapid kinetic characterization of hammerhead ribozymes by real-time monitoring of fluorescence resonance energy transfer (FRET)", RNA 5:1348 (1999).
- 10 Singh, K.K., et al., "Fluorescence Polarization for Monitoring Ribozyme Reactions in Real Time", BioTechniques 29:344 (2000).
- Sirinarumit, T., P. S. Paul, et al. (1997). "Rapid in situ hybridization technique for the detection of ribonucleic acids in tissues using radiolabelled and fluorescein-labelled riboprobes." Mol Cell Probes 11(4): 273-80.
- 15 Sokol, et al., Proc. Natl. Acad. Sci. 95(20): 11538-43 (1998).
- Sood, et al., Nucl. Acid Res. 4:2557 (1977).
- 20 Soukup, et al. Journal of Molecular Biology 298: 623-632 (2000).
- Sproat, et al., Nucl. Acid Res. 19:733-738 (1991).
- 25 Soukup, G., et al., "Generating new ligand-binding RNAs by affinity maturation and disintegration of allosteric ribozymes", RNA 7:524 (2001).
- Stanton, M., Hamaguchi, N., and Ellington, A. Poster Presentation at the National Cancer Institute Innovative Molecular Analysis Technologies P1 Meeting, July 6-8, Chantilly, Virginia
- 30 (2000)
- Stanton, J. A. and M. S. Beer (1997). "Characterisation of a cloned human 5-HT1A receptor cell line using [35S]GTP gamma S binding." Eur J Pharmacol 320(2-3): 267-75.
- 35 Tasset, et al. J Mol Biol, 272(5): 688-98 (1997).
- Truskey, G.A. et al., "Total internal reflection fluorescence microscopy (TIRFM): Topographical mapping of relative cell/substratum separation distances", J. Cell Science
- 40 103:491 (1992).
- Tuerk and Gold, Science 249: 505(1990)
- Taylor, S. J., R. J. Resnick, et al. (2001). "Nonradioactive determination of Ras-GTP levels using activated ras interaction assay." Methods Enzymol 333: 333-42.
- 45 Tyagi, et al., Nature Biotechnology 14: 303-308 (1996).
- Usman, N. and L. M. Blatt (2000). "Nuclease-resistant synthetic ribozymes: developing a new class of therapeutics." J Clin Invest 106(10): 1197-202.
- 50

- Walter, N.G. and Burke, J.M., "Real-time monitoring of hairpin ribozyme kinetics through base-specific quenching of fluorescein-labeled substrates", RNA 3:392 (1997).
- 5 Walter, N.G. et al., "Tertiary structure formation in the hairpin ribozyme monitored by fluorescence resonance energy transfer", The EMBO Journal 17(8):2378 (1998).
- Wang, P., J. G. Myers, et al. (1997). "Expression, purification, and characterization of human cAMP-specific phosphodiesterase (PDE4) subtypes A, B, C, and D." Biochem
- 10 Biophys Res Commun 234(2): 320-4.
- Wenzel-Seifert, K. and R. Seifert (2000). "Molecular analysis of beta(2)-adrenoceptor coupling to G(s)-, G(i)-, and G(q)-proteins." Mol Pharmacol 58(5): 954-66.
- 15 Williams KP; Imahori H; Fujimoto DN; Inoue T. Selection of novel forms of a functional domain within the Tetrahymena ribozyme. Nucleic Acids Research, 1994 Jun 11, 22(11):2003-9.
- Wilson C; Szostak JW. *In vitro* evolution of a self-alkylating ribozyme. Nature, 1995Apr 27, 374(6525):777-82.
- 20 Wu, P. and Brand, L., "Resonance Energy Transfer: Methods and Applications", Anal. Biochem. 218:1 (1994)
- 25 Wu, T.P. et al., "A fluorescent-labeling method for sequencing small RNA on polyacrylamide gel", Nucleic Acids Res. 24(17):3472 (1996).
- Zammatteo, N. et al., "Comparison between Different Strategies of Covalent Attachment of DNA to Glass Surfaces to Build DNA Microarrays", Anal. Biochem. 280:143 (2000).
- 30 Zhang B; Cech TR. Peptide bond formation by *in vitro* selected ribozymes. Nature, 1997 Nov 6, 390(6655):96-100.
- Bassi, G.S., de Oliveira, D.M., White, M.F. and Weeks, K.M. (2002) Proc Natl Acad Sci USA 99, 128-33.
- 35 Chang, L. and Karin, M. (2001) Nature 410, 37-40.
- Doudna, J.A. and Cech, T.R. (2002) Nature 418, 222-8.
- 40 English, J.M. and Cobb, M.H. (2002) Trends Pharm. Sci. 23, 40-45.
- Koizumi, M., Soukup, G.A., Kerr, J.Q. and Breaker, R.R. (1999) Nat. Struct. Biol. 6, 1062-71.
- 45 Myers, C.A., Kuhla, B., Cusack, S. and Lambowitz, A.M. (2002) Proc Natl Acad Sci USA 99, 2630-5.
- Pan, T. (1995) Biochemistry 34, 902-9.
- 50

- Robertson, M.P. and Ellington, A.D. (1999) Nat. Biotech. 17, 62-66.
- Robertson, M.P. and Ellington, A.D. (2000) Nucleic Acids Res. 28, 1751-9.
- 5 Russell R. and Herschlag, D. (1999) J. Mol. Biol. 291, 1155-67.
- Sassanfar and Szostak (1993) Nature 363, 550-553.
- Soukup, G.A. and Breaker, R.R. (1999) Proc Natl Acad Sci USA 96, 3584-9.
- 10 Soukup, G.A., DeRose, E.C., Koizumi, M. and Breaker, R.R. (2001) RNA 7, 524-36.
- Tang, J. and Breaker, R.R. (1997) Chem. Biol. 4, 453-9.
- 15 Walter, N.G., Harris, D.A., Pereira, M.J. and Rueda D. (2001-2) Biopolymers 61, 224-42.
- Knusel, B., Rabin, S.J., Hefti, F., Kaplan, D.R. 1994. Regulated neurotrophin receptor responsiveness during neuronal migration and early differentiation. J. Neurosci 14:1542
- 20 Fox, T., Coll, J.T., Xie, Xiaoling, Ford, P.J., Germann, U.A., Porter, M.D., Pazhanisamy, S., Fleming, M.A., Galullo, V., Su, M.S. and Wilson, K.P. (1998). "A single amino acid substitution makes ERK2 susceptible to pyridinyl imidazole inhibitors of p38 MAP kinase". Protein Science 7: 2249-55.
- 25 Meggio, F., Donella, D.A., Ruzzene, M., Brunati, A.M., Cesaro, L., Guerra, B., Meyer, T., Mett, H., Fabbro, D., Furet, P., *et al.* (1995). "Different susceptibility of protein kinases to staurosporine inhibition". Eur J Biochem 234: 317-22.

Variations, modifications, and other implementations of what is described herein

30 will occur to those of ordinary skill in the art without departing from the spirit and scope as claimed. Accordingly, the invention is to be defined not by the preceding illustrative description but instead by the spirit and scope of the following claims.

We claim:

1. A nucleic acid sensor molecule comprising
  - (a) a target modulation domain, wherein said target modulation domain recognizes a target molecule;
  - 5 (b) a linker domain; and
  - (c) a catalytic domain,wherein said nucleic acid sensor molecule comprises an optical signal generating unit.
- 10 2. The nucleic acid sensor molecule of claim 1, wherein said optical signal generating unit includes at least one signaling moiety.
3. The nucleic acid sensor molecule of claim 1, wherein said optical signal generating unit comprises at least a first signaling moiety and a second signaling moiety.
- 15 4. The nucleic acid sensor molecule of claim 3, wherein said first and second signaling moieties change proximity to each other upon recognition of a target by the target modulation domain.
- 20 5. The nucleic acid sensor molecule of claim 4, wherein said first and second signaling moieties comprise a fluorescent donor and a fluorescent quencher, and recognition of a target by the target modulation domain results in an increase in detectable fluorescence of said fluorescent donor.
- 25 6. The nucleic acid sensor molecule of claim 4, wherein said first signaling moiety and said second signaling moiety comprise fluorescent energy transfer (FRET) donor and acceptor groups, and recognition of a target by the target modulation domain results in a change in distance between said donor and acceptor groups, thereby changing optical properties of said molecule.
- 30 7. The nucleic acid sensor molecule of claim 1, wherein said optical signal generating unit consists essentially of a first signaling moiety, wherein said first

signaling moiety changes conformation upon recognition of a target by the target modulation domain, thereby resulting in a detectable optical signal.

5 8. The nucleic acid sensor molecule of claim 1, wherein said nucleic acid sensor molecule includes at least one modified nucleic acid.

9. The nucleic acid sensor molecule of claim 1, wherein said catalytic domain comprises an endonucleolytic ribozyme.

10 10. The nucleic acid sensor molecule of claim 9, wherein said endonucleolytic ribozyme is a *cis*-endonucleolytic ribozyme or a *trans*-endonucleolytic ribozyme.

11. The nucleic acid sensor molecule of claim 9, wherein said  
15 endonucleolytic ribozyme is a hammerhead ribozyme.

12. The nucleic acid sensor molecule of claim 1, wherein said catalytic domain comprises a self-ligating ribozyme.

20 13. The nucleic acid sensor molecule of claim 12, wherein said self-ligating ribozyme is a *cis*-ligase ribozyme or a *trans*-ligase ribozyme.

14. The nucleic acid sensor molecule of claim 12, wherein said self-ligating  
25 ribozyme is a 1-piece ligase, 2-piece ligase or 3-piece ligase.

15. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a target selected from the group consisting of  
30 proteins, post-translationally modified forms of proteins, peptides, nucleic acids, oligosaccharides, nucleotides, metabolites, drugs, toxins, biohazards, ions, carbohydrates, polysaccharides, hormones, receptors, antigens, antibodies, viruses, metabolites, co-factors, drugs, dyes, nutrients, and growth factors.



16. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a target selected from the group consisting of a protein and a post-translationally modified protein.

5           17. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a target that is a post-translationally modified protein, and wherein the post-translation modifications are selected from the group consisting of: phosphorylation, prenylation, glycosylation, methionine removal, N-acetylation, acylation, acylation of cysteines, myristoylation, alkylation,  
10           ubiquitinylation, prolyl-4-hydroxylation, carboxylation of glutaminy residues, advanced glycosylation, deamination of glutamine and asparagine, addition of glycoposphatidylinositol, disulfide bond formation, hydroxylation, and lipidation.

15           18. The nucleic acid sensor molecule of claim 1, wherein said target is a protein kinase.

19. The nucleic acid sensor molecule of claim 1, wherein said target is a phosphorylated protein kinase.

20           20. The nucleic acid sensor molecule of claim 19, wherein said phosphorylated protein kinase is a monophosphorylated protein kinase or a diphosphorylated protein kinase.

25           21. The nucleic acid sensor molecule of claim 18, wherein said protein is ERK.

22. The nucleic acid sensor molecule of claim 19, wherein said post-translationally modified protein is pp ERK.

30           23. The nucleic acid sensor molecule of claim 18 wherein said protein kinase is ERK1 or ERK2.

24. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a component of a MAP kinase pathway, a product of

a MAP kinase pathway, a MAP kinase pathway associated protein, or an extracellular component of a MAP kinase pathway.

5           25. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a component of the ERK1/2 MAP kinase pathway.

          26. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a component of the JNK MAP kinase pathway.

10           27. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a component of the P38 MAP kinase pathway.

          28. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes an endogenous form of a MAP Kinase (MEK), an endogenous form of a MAP Kinase Kinase (MEKK), or an endogenous form of MAP Kinase Kinase Kinase, (MEKKK).

          29. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes an endogenous form of RAF kinase.

20           30. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a Ras protein.

          31. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a phosphatase.

          32. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a GTP binding protein.

30           33. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a GPCR.

          34. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a cytokine.

35. The nucleic acid sensor molecule of claim 1, wherein said target modulation domain recognizes a growth factor.

5           36. The nucleic acid sensor molecule of claim 1, wherein said target recognition domain recognizes a cellular metabolite.

          37. The nucleic acid sensor molecule of claim 1, wherein said target recognition domain recognizes a small molecule.

10           38. The nucleic acid sensor molecule of claim 1, wherein said nucleic acid sensor molecule comprises RNA, DNA, or both RNA and DNA.

          39. A composition comprising the nucleic acid sensor molecule of any one of claims 1-38 or claims 74-99 and a buffer.

          40. A composition comprising the nucleic acid sensor molecule of any one of claims 1-38 or claims 74-99 and a tissue extract, a cell extract or an *in vitro* cell culture.

20           41. The composition of claim 39 or 40, further comprising an RNase inhibitor.

          42. The composition of claim 41, wherein said RNase inhibitor is selected from the group consisting of Va-riboside, vanadyl, tRNA, polyU, RNaseIn and RNaseOut.

          43. The composition of claim 39 or 40, wherein said composition is substantially RNase-free.

30           44. A composition comprising at least one nucleic acid sensor molecule according to any one of claims 1-38 or claims 74-99, affixed to a substrate.

45. The composition of claim 44, wherein said substrate is glass, gold or other metal, silicon or other semiconductor material, nitrocellulose, nylon, or plastic.

5 46. The composition of claim 44, wherein the nucleic acid sensor molecule is covalently attached to said substrate.

47. The composition of claim 44, wherein the nucleic acid sensor molecule is non-covalently attached to said substrate.

10 48. The composition of claim 44, wherein the nucleic acid sensor molecule is immobilized to the substrate via hybridization of a terminal portion of the nucleic acid sensor molecule to an oligonucleotide that is bound to the surface of the substrate.

15 49. The composition of claim 44, wherein said composition comprises a plurality of nucleic acid sensor molecules immobilized to the substrate via hybridization of a terminal portion of the nucleic acid sensor molecule to an array of oligonucleotides bound to the substrate at spatially discrete regions.

20 50. The composition of claim 44, wherein at least two members of said plurality each recognize different target molecules.

51. The substrate of claim 44, wherein said substrate comprises at least 50 nucleic acid sensor molecules.

25 52. The substrate of claim 44, wherein said substrate comprises at least 250 nucleic acid sensor molecules.

30 53. The substrate of claim 44, wherein said substrate comprises at least 500 nucleic acid sensor molecules.

54. The substrate of claim 44, wherein said substrate comprises at least 5000 nucleic acid sensor molecules.

55. A system for detecting a target molecule comprising a composition according to any one of claims 41-54, and a detector in optical communication with said composition, wherein said detector detects changes in the optical properties of said composition.

5

56. The system of claim 55, further comprising a light source in optical communication with said composition.

10

57. The system of claim 56, further comprising a processor for processing optical signals detected by the detector.

15

58. The system of claim 55, wherein said system comprises a plurality of nucleic acid sensor molecules, wherein at least two of said biosensor molecules each recognize different target molecules.

20

59. A method of identifying or detecting a target molecule in a sample, the method comprising:  
contacting a sample suspected of containing a target molecule with a nucleic acid sensor molecule according to any one of claims 1-38 or claims 74-99, wherein a change in the signal generated by the optical signal generating unit indicates the presence of said target in said sample.

25

60. The method of claim 59 further comprising quantifying the change signal generated by the optical signal generating unit to quantify the amount of target molecule in the sample.

30

61. The method of claim 59 or 60 wherein the sample is selected from the group consisting of: environmental samples, biohazard materials, organic samples, drugs and toxins, flavors and fragrances, and biological samples.

62. The method of claim 59 or 60 wherein the sample is a biological sample, including cells, cell extracts or lysates, tissues or tissue extracts, bodily fluids, serum, blood and blood products.

5 63. The method of claim 59 or 60 wherein the target is selected from the group consisting of proteins, post-translationally modified forms of proteins, peptides, nucleic acids, oligosaccharides, nucleotides, metabolites, drugs, toxins, biohazards, ions, carbohydrates, polysaccharides, hormones, receptors, antigens, antibodies, viruses, metabolites, co-factors, drugs, dyes, nutrients, and growth factors.

10 64. The method of claim 59 or 60 wherein the target is selected from the group consisting of proteins and post-translationally modified forms of proteins.

15 65. The method of claim 64, wherein said target is a post-translationally modified protein, and wherein the post-translation modifications are selected from the group consisting of: phosphorylation, prenylation, glycosylation, methionine removal, N-acetylation, acylation, acylation of cysteines, myristoylation, alkylation, ubiquitinylation, prolyl-4-hydroxylation, carboxylation of glutaminy residues, advanced glycosylation, deamination of glutamine and asparagine, addition of glycophosphatidylinositol, disulfide bond formation, hydroxylation, and lipidation.

20 66. The method of claim 59 or 60 wherein the target is a protein kinase.

67. The method of claim 59 or 60, wherein said target is a phosphorylated protein kinase.

25 68. A diagnostic system for identifying or detecting a target molecule, the diagnostic system comprising  
a nucleic acid sensor molecule according to any one of claims 1-38 or 74-99;  
and  
a detector in communication with said nucleic acid sensor molecule, wherein  
said detector detects changes in the signal generated by the optical signal generating  
30 unit of said nucleic acid sensor.

69. The diagnostic system of claim 68, further comprising a processor for processing signals detected by the detector.

70. A method of identifying or detecting a protein kinase in a sample, the method comprising:

contacting a sample suspected of containing a protein kinase with a nucleic acid sensor molecule according to claim 1, wherein said nucleic acid sensor molecule has a target recognition domain that recognizes a protein kinase, wherein a change in the signal generated by the optical signal generating unit indicates the presence of protein kinase in said sample.

71. The method of claim 70, further comprising quantifying the amount of signal generated by the optical signal generating unit to quantify the amount of protein kinase in the sample.

72. A method of identifying a modulator of protein kinase activity, the method comprising:

contacting a test agent with a protein kinase and nucleic acid sensor molecule according to claim 1, wherein said nucleic acid sensor molecule has a target recognition domain that recognizes a protein kinase, wherein recognition of the protein kinase by the target recognition domain of said nucleic acid sensor molecule results in a change in the signal generated by the optical signal generating unit and further wherein changes in the signal generated by the optical signal generating unit in the presence and absence of said test agent indicates the test agent is a modulator of said protein kinase activity.

73. The method of claim 70 or 72, wherein the catalytic domain of said nucleic acid sensor molecule comprises a *cis*-ligase ribozyme or a *trans*-ligase ribozyme.

74. A nucleic acid sensor molecule comprising:  
a target modulation domain that recognizes ERK;  
a catalytic domain that comprises a ligase or *cis*-hammerhead; and  
a linker domain that links said target modulation domain and said catalytic domain.

75. A nucleic acid sensor molecule comprising:

a target modulation domain that recognizes phosphoERK;  
a catalytic domain that comprises a ligase or a cis-hammerhead; and  
a linker domain that links said target modulation domain and said catalytic domain.

5

76. A nucleic acid sensor molecule comprising:  
a target modulation domain that recognizes lysozyme;  
a catalytic domain that comprises a 1-piece cis-ligase; and  
a linker domain that links said target modulation domain and said catalytic

10

domain.

77. A nucleic acid sensor molecule comprising:  
a target modulation domain that recognizes any one of cCMP, cAMP, or cGMP;  
a catalytic domain; and  
a linker domain that links said target modulation domain and said catalytic domain, wherein said nucleic acid sensor molecule comprises an optical signal generating unit or a non-radioactive detectable label.

15

20

78. The nucleic acid sensor molecule of any one of claims 74-76 wherein the nucleic acid sensor molecule comprises an optical signal generating unit.

79. The nucleic acid sensor molecule of any one of claims 74-76, wherein said nucleic acid sensor molecule comprises a detectable label.

25

80. The nucleic acid sensor molecule of claim 79, wherein the label is a radioactive label.

30

81. The nucleic acid sensor molecule of claim 79, wherein the radioactive label is  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{14}\text{C}$ ,  $^{35}\text{S}$ ,  $^3\text{H}$ , or  $^{125}\text{I}$ .

82. The nucleic acid sensor molecule of any one of claims 74-77, wherein said nucleic acid sensor molecule comprises a fluorescent label.



83. The nucleic acid sensor molecule of claim 82, wherein the fluorescent label is fluorescein, DABCYL, or a green fluorescent protein (GFP) moiety.

5 84. The nucleic acid sensor molecule of claim 77 or claim 78, wherein the optical signal generating unit comprises a fluorescent moiety and a quenching moiety, wherein recognition by the target modulation domain causes a change in detectable fluorescence by the optical signal generating unit.

10 85. The nucleic acid sensor molecule of any one of claims 74-77, wherein said nucleic acid sensor molecule comprises an enzymatic label.

86. The nucleic acid sensor molecule of any one of claims 74-77, wherein said nucleic acid sensor molecule comprises an affinity capture tag label.

15 87. The nucleic acid sensor molecule of claim 74, wherein the target modulation domain recognizes ERK1, ERK2 or both.

20 88. The nucleic acid sensor molecule of claim 74, wherein the target modulation domain and the catalytic domain are as shown in any one of SEQ ID NO. 80 and the linker is randomized.

25 89. The nucleic acid sensor molecule of claim 74, wherein the target modulation domain and the catalytic domain are as shown in any one of SEQ ID NOS. 47, 118 and 119 and the linker is randomized.

90. A nucleic acid sensor molecule that recognizes ERK comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 90-95, 108-116, 131-133, 140-295, 349, 351, and 356.

30 91. A nucleic acid sensor molecule that recognizes phosphoERK comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 5-8, 37-39, 44-45, 81-89, 96-100, 121-130, 352, and 353.

92. A nucleic acid sensor molecule that recognizes any one of cCMP, cAMP or cGMP comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 40-43, 103, and 135-139.

5           93. A nucleic acid sensor molecule that recognizes lysozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 46, 47, 76, 105-107.

10           94. A 1-piece ligase ribozyme comprising  
a target modulation domain that recognizes a target;  
a linker domain;  
a catalytic domain,  
wherein the 5' and 3' ends of the ligase ligate to each other upon recognition  
of the target by the modulation domain.

15           95. A 2-piece ligase ribozyme comprising  
a target modulation domain that recognizes a target;  
a linker domain;  
a catalytic domain comprising an oligonucleotide substrate ligation site and  
20 and oligonucleotide supersubstrate binding domain, wherein upon recognition of the  
target by the modulation domain, the 3' end of the an oligonucleotide supersubstrate  
is ligated to the 5' end of the oligonucleotide substrate ligation site.

25           96. A 3-piece ligase ribozyme comprising  
a target modulation domain that recognizes a target;  
a linker domain;  
a catalytic domain comprising comprising an oligonucleotide substrate  
binding domain capable of binding an oligonucleotide substrate and an effector-  
oligonucleotide binding site capable of binding an effector oligonucleotide, wherein  
30 upon recognition of the target by the modulation domain, and in the presence of  
binding of the effector oligonucleotide to the effector-oligonucleotide binding site,  
then the 3' end of the oligonucleotide substrate is ligated to the 5' end of the ligase.

97. A 1-piece ligase ribozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 47, 105-107, 119.

5 98. A 2-piece ligase ribozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 347, 349, and 351.

99. A 3-piece ligase ribozyme comprising the nucleic acid sensor molecule shown in any one of SEQ ID NOS. 46, 75, 76, 108-116, 118, 121-130, and 352.





FIGURE 1A

Figure 1B



*R signifies the hammerhead ribozyme*







FIGURE 2



### UNQUENCHED LIGAND-BOUND FORM



### QUENCHED NO-LIGAND FORM

**FIGURE 3**




FIGURE 4






FIGURE 5

A                    A    CC    CA                    SEQ ID NO:49  
A GGC UUGG GUGGUAU -D  
A CCG AACCC A-P  
G    --    AU

**UNQUENCHED LIGAND-BOUND FORM**

B                    CC    GA                    SEQ ID NO:49  
G UUGGC UGGUAU-D  
G AGCCG ACCAUA-F  
AA            -A

**QUENCHED NO-LIGAND FORM****FIGURE 6**




FIGURE 7




FIGURE 8




Figure 9.




Figure 10.




FIGURE 11




FIGURE 12



Figure 13






Figure 14.





RNA sensor SEQ ID NO:78  
capture oligo SEQ ID NO:79




FIGURE 16




FIGURE 17




FIGURE 18

SEQ ID NO:80

Target modulation domain






Figure 19


	top -5'NNNN 3' bottom-3'NNNN 5'	Activity
1-14	ACGU GCUGG	+++
1-13	UACGU UUCU	+++
1-2	ACGU UUC	++
1-6	ACGU UAUC	+
2-7	CAGU UUC	+++
2-2	CUAG AUCC	+
2-3	UUAA UCAA	+
2-13	CUGG AAAU	+
2-14	GAUC UUUU	+
2-20	UUAA AGCG	+

Figure 20A





**Figure 20B**



y = m1*(1 - exp(-m2*t))		
	Value	Error
m1	0.92027	0.0004329
m2	0.043998	0.00022795
ChiSq	1.9713e-05	NA
R	0.99998	NA

y = m1 + m2 * t		
	Value	Error
m1	0.01036	0.008938
m2	0.0005519	0.0001592
ChiSq	0.00026998	NA
R	0.94807	NA

y = m1*(1 - exp(-m2*t))		
	Value	Error
m1	0.85753	0.011573
m2	0.000051	0.00163
ChiSq	0.0001716	NA
R	0.99995	NA

Figure 20C




Figure 21




Figure 22A




Figure 22B

SEQ ID NO: 353

# Target Modulation Domain




Figure 23A


Construct 6, SEQ ID NO: 81  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUGUJACGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 7, SEQ ID NO: 82  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUUUACGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 8, SEQ ID NO: 83  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUGUUCGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 9, SEQ ID NO: 84  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUUUACGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 10, SEQ ID NO: 85  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUGUUCGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 11, SEQ ID NO: 86  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUGCGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 12, SEQ ID NO: 87  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUGGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 13, SEQ ID NO: 88  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUCAACGAAACGGUGAAAGGCCGUAAGGUUGCC  
Construct 14, SEQ ID NO: 89  
GGCGACCCUGAUGAGUCACGCGACGCUAGCGAAUUGGUUCCUCCGAAAGGGGAAAGCGUUUAAGAAACCAAAUUGUCUACGAAACGGUGAAAGGCCGUAAGGUUGCC

**Figure 23B**


**Figure 24**

	sequences	activity	stability
Construct 6	GUGUUA CGCACU	++	-32.41Kcal
Construct 7	UGUUA GCACU	Constitutively active	-29.11Kcal
Construct 8	GUGUU CGCAC	Constitutively active	No predicted duplex formation
Construct 9	GUUUA CGACU	Constitutively active	-29.41Kcal
Construct 10	GUGUUG CGCACU	++	-32.7Kcal
Construct 11	GUG CGU	Constitutively active	-30.71Kcal
Construct 12	GUACGU CGUCC	+++	No predicted duplex formation
Construct 13	GUCAUA CGGUCU	Constitutively active	-32.41Kcal
Construct 14	GUCUUA CGGUCU	Constitutively active	No predicted duplex formation






**B**



**Figure 25**

Figure 26






Figure 27








Figure 28



**Figure 29**

Figure 30





## Figure 31


**SEQ ID NO: 18**

## Sequences of ERK-modulated ligase nucleic acid sensor molecules


DG.20.58A SEQ ID NO: 109 construct 17  
GGACUUCGGCGAAAGCCGUUUCGACCCACGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUGUCUUAGACAG  
GAGGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58B SEQ ID NO: 110 construct 18  
GGACUUCGGCGAAAGCCGUUUCGACCCACGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUGUCUUAGACAGGA  
GGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58C SEQ ID NO: 112 construct 20  
GGACUUCGGCGAAAGCCGUUUCGACCCACGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCCAUUGUCUUAGACAG  
GAGGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58D SEQ ID NO: 113 construct 21  
GGACUUCGGCGAAAGCCGUUUCGACCCACGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUCUUAGACAG  
GAGGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58E SEQ ID NO: 111 construct 19  
GGACUUCGGCGAAAGCCGUUUCGACCCACGCUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGCUCUUAGACAGGA  
GGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58F SEQ ID NO: 114 construct 22  
GGACUUCGGCGAAAGCCGUUUCGACCCGCGUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUGCUUAGACAGGA  
GGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58G SEQ ID NO: 115 construct 23  
GGACUUCGGCGAAAGCCGUUUCGACCCGCGUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUCUUAGACAGGAGG  
UUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58H SEQ ID NO: 117 construct 25  
GGACUUCGGCGAAAGCCGUUUCGACCCGCGUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAUUGCUUAGACAGGA  
GGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58I SEQ ID NO: 118 construct 26  
GGACUUCGGCGAAAGCCGUUUCGACCCGCGUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGUCUUAGACAGGA  
GGUUAGGUGCCGUCGACUGAUCUGCGAGUUAACG  
DG.20.58J SEQ ID NO: 116 construct 24  
GGACUUCGGCGAAAGCCGUUUCGACCCGCGUAAGGAGGAUUUCCGAAAGCGGCUACGGUCCGCCAGCCUUAGACAGGAGG  
UUAGGUGCCGUCGACUGAUCUGCGAGUUAACG


Figure 32





**Figure 33**


**Figure 34**

**Figure 35**

20 nM ERK:  $k_{obs} = 8 \times 10^{-5} \text{ min}^{-1}$

200 nM ERK:  $k_{obs} = 5 \times 10^{-4} \text{ min}^{-1}$

**Figure 36**






Figure 37






Figure 38




**Figure 39A**



**Figure 39B**



**Figure 40**

**SEQ ID NO: 352**

GCAAUUGAGGCUUAGUCAGCCUGC U — Sequence tag

U-A oligonucleotide substrate  
G-C  
G II

## Target-modulation domain

## Figure 41

# Construct sequence Data

Constructs (these are RNA transcript sequences given as DNA sequences)

Construct 31 (TK.16.118.K) + SEQ ID NO: 121  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
AGTGCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 32 (TK.16.118.L) + SEQ ID NO: 122  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
GTCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 33 (TK.16.118.M) ++ SEQ ID NO: 123  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
ATTGCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 34 (TK.16.118.N) I SEQ ID NO: 124  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
ACGTTCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 35 (TK.16.118.O) C SEQ ID NO: 125  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
GCTCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 36 (TK.16.118.P) I SEQ ID NO: 126  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
GTGCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 37 (TK.16.118.Q) ++ SEQ ID NO: 127  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATGAG  
TCCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 38 (TK.16.118.R) ++ SEQ ID NO: 128  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
TTGCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 39 (TK.16.118.S) I SEQ ID NO: 129  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATG  
CGTCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

Construct 40 (TK.16.118.T) + SEQ ID NO: 130  
GGACTTCGGCGAAAGCCGTTCCGACCGCTAGCGAATTGGTTCCTCGAAAGGGGAAAGCGTTATTAAAGAAACCAAAATGAG  
CCTTAGACACAGGAGGTTAGGTGCGGCTTTGGTCCAAGGACATCTCGAAT

I Inactive  
C Constitutively Active  
+ Effector-dependent  
++ Greater Effector-Dependence than +

Figure 42

## Figure 43

## •ppERK library

5'GGACTTCGGCGAAGCCGTTTCGACCNNNN(N)(N)CAGACGCTA  
GCGAATTGGTTCCTCGAAGGGAAGCGTTATTAAAGAAACCAAAA  
TGNNNN(N)(N)CTTAGACAGGAGGTAGGTGCGTCAATGCTGCAA  
GTIACTG 3' SEQ ID NO: 354

## •ERK library

GGACTTCGGCGAAGCCGTTTCGACCNNNN(N)(N)AAGGAGGATTC  
CGAAGCGGCTACGGTCCGCCNN(N)(N)CTTAGACAGGAGGTTA  
GGTGCGTAGGTAAACCGATAGTTCCG SEQ ID NO:355

## Figure 44A

SEQ ID NO:356	AAAAG UUUU	CW45-33-D09 (17.3, 5/18)	SEQ ID NO:133
AGAGGCU	ACAAG UGUCC	CW45-33-H08 (36, 1/18)	SEQ ID NO:91
UCUC CC	AGCGA UCGCU	CW45-33-H09 (33, 1/18)	SEQ ID NO:92
CW45-33-A08 (3.18, 2/18)	GGUUG UCAA	CW45-33-A10 (11.4, 1/18)	SEQ ID NO:93
AGGGG3'	AACAG UAGU	CW45-33-F09 (14.2, 2/18)	SEQ ID NO:94
UCUCC 5'	AGUAG UCAU	CW45-33-G08 (11.4, 1/18)	SEQ ID NO:95
CW45-33-C08 (11.3, 1/18)			
SEQ ID NO:131			
CW45-33-C09 (18.6, 1/18)			
SEQ ID NO:132			
CW45-33-F08 (9.5, 1/18)			
SEQ ID NO:90			

## Designed Library:

3-5 nt stem  
25% of each A, C, U, G  
37.5% complementary

## Result of selection:

4-5 nt stem  
27.8% A  
27.8% U  
25.6% G  
18.9% C  
1st pair 90% A-U  
2nd pair 60% GC  
95.5% complementary

GAACCGCU GUUGG CC CW45-33-A02 (4.46, 1/31) One C->U mutation in 5' ligase SEQ ID NO:44	UAGA <sup>A</sup> ---C <sup>C</sup> AGCUU-G <sup>C-G</sup> T <sup>T</sup> G <sup>G</sup>	UGAUCG3' GUUGG5' CW45-33-B04(5.05, 5/31) SEQ ID NO:45 CGUGU GUACG CW45-33-C04(6.2, 1/31) SEQ ID NO:5 UUAUCG GAUGG CW45-33-D04(9.0, 1/31) One deletion, one A to G mutation in aptamer structure SEQ ID NO:6 CGCAGG GGUGUC CW45-33-F03(4, 1/31) SEQ ID NO:7	CW45-33-D01(2, 1/31) AGGG UCAUC SEQ ID NO:37 CW45-33-D02(3, 1/31) AGUAAG UCUU CW45-33-D05(2, 1/31) CGGUGG GACGC CW45-33-E01(2, 1/31) GCAGAU UGUUUA CW45-33-G02(3, 1/31) AUUAGA UGAUUU CW45-33-G03(3, 1/31) AUCGG UCGC CW45-33-H03(2, 1/31)	Designed Library: 4-6 nt stem 25% of each A, C, U, G 37.5% complementary Result of selection: 4-6 nt stem 4-6 pairing 18.0% A 28.9% U 36.7% G 16.4% C 89.8% complementary GUGUGG AACGC CW45-33-D01(2, 1/31) CAAU GUUG CW45-33-H1(1, 3/31) CUAA GAUU CW45-33-B05(1, 1/31)	SEQ ID NO:8 SEQ ID NO:99 SEQ ID NO:100 SEQ ID NO:101 SEQ ID NO:102 SEQ ID NO:103 SEQ ID NO:104 SEQ ID NO:105 SEQ ID NO:106 SEQ ID NO:107 SEQ ID NO:108 SEQ ID NO:109 SEQ ID NO:110 SEQ ID NO:111 SEQ ID NO:112 SEQ ID NO:113 SEQ ID NO:114 SEQ ID NO:115 SEQ ID NO:116 SEQ ID NO:117 SEQ ID NO:118 SEQ ID NO:119 SEQ ID NO:120 SEQ ID NO:121 SEQ ID NO:122 SEQ ID NO:123 SEQ ID NO:124 SEQ ID NO:125 SEQ ID NO:126 SEQ ID NO:127 SEQ ID NO:128 SEQ ID NO:129 SEQ ID NO:130 SEQ ID NO:131 SEQ ID NO:132 SEQ ID NO:133 SEQ ID NO:134 SEQ ID NO:135 SEQ ID NO:136 SEQ ID NO:137 SEQ ID NO:138 SEQ ID NO:139 SEQ ID NO:140 SEQ ID NO:141 SEQ ID NO:142 SEQ ID NO:143 SEQ ID NO:144 SEQ ID NO:145 SEQ ID NO:146 SEQ ID NO:147 SEQ ID NO:148 SEQ ID NO:149 SEQ ID NO:150 SEQ ID NO:151 SEQ ID NO:152 SEQ ID NO:153 SEQ ID NO:154 SEQ ID NO:155 SEQ ID NO:156 SEQ ID NO:157 SEQ ID NO:158 SEQ ID NO:159 SEQ ID NO:160 SEQ ID NO:161 SEQ ID NO:162 SEQ ID NO:163 SEQ ID NO:164 SEQ ID NO:165 SEQ ID NO:166 SEQ ID NO:167 SEQ ID NO:168 SEQ ID NO:169 SEQ ID NO:170 SEQ ID NO:171 SEQ ID NO:172 SEQ ID NO:173 SEQ ID NO:174 SEQ ID NO:175 SEQ ID NO:176 SEQ ID NO:177 SEQ ID NO:178 SEQ ID NO:179 SEQ ID NO:180 SEQ ID NO:181 SEQ ID NO:182 SEQ ID NO:183 SEQ ID NO:184 SEQ ID NO:185 SEQ ID NO:186 SEQ ID NO:187 SEQ ID NO:188 SEQ ID NO:189 SEQ ID NO:190 SEQ ID NO:191 SEQ ID NO:192 SEQ ID NO:193 SEQ ID NO:194 SEQ ID NO:195 SEQ ID NO:196 SEQ ID NO:197 SEQ ID NO:198 SEQ ID NO:199 SEQ ID NO:200 SEQ ID NO:201 SEQ ID NO:202 SEQ ID NO:203 SEQ ID NO:204 SEQ ID NO:205 SEQ ID NO:206 SEQ ID NO:207 SEQ ID NO:208 SEQ ID NO:209 SEQ ID NO:210 SEQ ID NO:211 SEQ ID NO:212 SEQ ID NO:213 SEQ ID NO:214 SEQ ID NO:215 SEQ ID NO:216 SEQ ID NO:217 SEQ ID NO:218 SEQ ID NO:219 SEQ ID NO:220 SEQ ID NO:221 SEQ ID NO:222 SEQ ID NO:223 SEQ ID NO:224 SEQ ID NO:225 SEQ ID NO:226 SEQ ID NO:227 SEQ ID NO:228 SEQ ID NO:229 SEQ ID NO:230 SEQ ID NO:231 SEQ ID NO:232 SEQ ID NO:233 SEQ ID NO:234 SEQ ID NO:235 SEQ ID NO:236 SEQ ID NO:237 SEQ ID NO:238 SEQ ID NO:239 SEQ ID NO:240 SEQ ID NO:241 SEQ ID NO:242 SEQ ID NO:243 SEQ ID NO:244 SEQ ID NO:245 SEQ ID NO:246 SEQ ID NO:247 SEQ ID NO:248 SEQ ID NO:249 SEQ ID NO:250 SEQ ID NO:251 SEQ ID NO:252 SEQ ID NO:253 SEQ ID NO:254 SEQ ID NO:255 SEQ ID NO:256 SEQ ID NO:257 SEQ ID NO:258 SEQ ID NO:259 SEQ ID NO:260 SEQ ID NO:261 SEQ ID NO:262 SEQ ID NO:263 SEQ ID NO:264 SEQ ID NO:265 SEQ ID NO:266 SEQ ID NO:267 SEQ ID NO:268 SEQ ID NO:269 SEQ ID NO:270 SEQ ID NO:271 SEQ ID NO:272 SEQ ID NO:273 SEQ ID NO:274 SEQ ID NO:275 SEQ ID NO:276 SEQ ID NO:277 SEQ ID NO:278 SEQ ID NO:279 SEQ ID NO:280 SEQ ID NO:281 SEQ ID NO:282 SEQ ID NO:283 SEQ ID NO:284 SEQ ID NO:285 SEQ ID NO:286 SEQ ID NO:287 SEQ ID NO:288 SEQ ID NO:289 SEQ ID NO:290 SEQ ID NO:291 SEQ ID NO:292 SEQ ID NO:293 SEQ ID NO:294 SEQ ID NO:295 SEQ ID NO:296 SEQ ID NO:297 SEQ ID NO:298 SEQ ID NO:299 SEQ ID NO:300 SEQ ID NO:301 SEQ ID NO:302 SEQ ID NO:303 SEQ ID NO:304 SEQ ID NO:305 SEQ ID NO:306 SEQ ID NO:307 SEQ ID NO:308 SEQ ID NO:309 SEQ ID NO:310 SEQ ID NO:311 SEQ ID NO:312 SEQ ID NO:313 SEQ ID NO:314 SEQ ID NO:315 SEQ ID NO:316 SEQ ID NO:317 SEQ ID NO:318 SEQ ID NO:319 SEQ ID NO:320 SEQ ID NO:321 SEQ ID NO:322 SEQ ID NO:323 SEQ ID NO:324 SEQ ID NO:325 SEQ ID NO:326 SEQ ID NO:327 SEQ ID NO:328 SEQ ID NO:329 SEQ ID NO:330 SEQ ID NO:331 SEQ ID NO:332 SEQ ID NO:333 SEQ ID NO:334 SEQ ID NO:335 SEQ ID NO:336 SEQ ID NO:337 SEQ ID NO:338 SEQ ID NO:339 SEQ ID NO:340 SEQ ID NO:341 SEQ ID NO:342 SEQ ID NO:343 SEQ ID NO:344 SEQ ID NO:345 SEQ ID NO:346 SEQ ID NO:347 SEQ ID NO:348 SEQ ID NO:349 SEQ ID NO:350 SEQ ID NO:351 SEQ ID NO:352 SEQ ID NO:353 SEQ ID NO:354 SEQ ID NO:355 SEQ ID NO:356 SEQ ID NO:357 SEQ ID NO:358 SEQ ID NO:359 SEQ ID NO:360 SEQ ID NO:361 SEQ ID NO:362 SEQ ID NO:363 SEQ ID NO:364 SEQ ID NO:365 SEQ ID NO:366 SEQ ID NO:367 SEQ ID NO:368 SEQ ID NO:369 SEQ ID NO:370 SEQ ID NO:371 SEQ ID NO:372 SEQ ID NO:373 SEQ ID NO:374 SEQ ID NO:375 SEQ ID NO:376 SEQ ID NO:377 SEQ ID NO:378 SEQ ID NO:379 SEQ ID NO:380 SEQ ID NO:381 SEQ ID NO:382 SEQ ID NO:383 SEQ ID NO:384 SEQ ID NO:385 SEQ ID NO:386 SEQ ID NO:387 SEQ ID NO:388 SEQ ID NO:389 SEQ ID NO:390 SEQ ID NO:391 SEQ ID NO:392 SEQ ID NO:393 SEQ ID NO:394 SEQ ID NO:395 SEQ ID NO:396 SEQ ID NO:397 SEQ ID NO:398 SEQ ID NO:399 SEQ ID NO:400 SEQ ID NO:401 SEQ ID NO:402 SEQ ID NO:403 SEQ ID NO:404 SEQ ID NO:40
--	--	---	---	--	--


## Figure 45

•CW45.33.A02 sequence SEQ ID NO:44

5'GGACUUCGGCGAAAGUCGUUCGACCGGUUGCAGACGCUAGCGA  
AUUGGUUCCUCGAAAGGGGAAAGCGUUAUUAAGAAACCAAAUUGGAA  
CCGCUUAGACAGGAGGUUAGGUGC3'


•CW45.33.D04 SEQ ID NO:6

5'GGACUUCGGCGAAAGCCGUUCGACCGGUAGCAGACGCUAGCGA  
AUUGGU\_CCUCGAAAGGGGAAAGCGUUAUUAAGAAACCAAAUUGUUA  
UCGCUUAGACAGGAGGUUAGGUGC3'



**Figure 46**





Switch factor =  $k_{act} / k_{unact}$

Figure 47




Figure 48





Figure 49



zyme]/uM  
amplicon  
Unligated

Reticulocyte Lysate    HeLa Lysate    Water    0.1x "Contaminant"  
RNasin, tRNA, 1h 25C, Phenol extraction, SDS-PAGE

Figure 50





Figure 51



**Figure 52**




Figure 53







Figure 54






**Figure 55**

Figure 56



Panel A: Specificity for phosphorylation



Panel B: Specificity for MAP kinase component




Figure 57

Figure 58





Figure 59

<u>ERK</u> <u>sensor</u>	<u>Switch</u> <u>factor</u>	<u>K<sub>D,act</sub></u> <u>μM</u>	<u>K<sub>D,ERK</sub></u> <u>μM</u>	<u>K<sub>unact</sub></u> <u>nM/min</u>	<u>Det.</u> <u>limit</u>
ERK aptamer	N/A	N/A	0.0043 ± 0.0005	N/A	N/A
Ligase 1	≥800	≥5	≥5	~0.004	40 nM
Ligase 2	76 ±20	0.75 ±0.08	—	0.04 ±0.01	10 nM
Ligase 3	65 ±16	0.28 ±0.08	1.2 ±0.1	~0.02	10 nM
Ligase 4	5 ±1	0.04 ±0.02	0.17 ±0.07	~0.3	40 nM

Figure 60






Figure 61






FIGURE 62




A.



B.



Expected time-course :



C.





FIGURE 63

A.



B.




FIGURE 64.









FIGURE 65A and 65B

**C.****D.****E.****FIGURE 65C, 65D and 65E**


A.



B.



C.




D.

E.


FIGURE 66

A.

## SPReeta SPR sensor




## NASM loading



C.


B.

## HH NASM



SEQ ID NO: 103 (same as cAMP)

## Analyte-induced cleavage



D.

FIGURE 67

cGMP (90 nt) SEQ ID NO: 135

5' thiol-GGAUAAUAGCCGUAGGUUGCGAAAGCGACCCUGAUGA  
GCCCUGCGAUGCAGAAAGGUGCUGACGACACAUCGAAACGGUAGCGAGAG  
CUC 3'

cCMP (88 nt): SEQ ID NO: 136

5' thiol-GGAUAAUAGCCGUAGGUUGCGAAAGCGACCCUGAUGA  
CCUGUGGAAACAGACGUGGCACAUGACUACGU  
GAAACGGUAGCGAGAGCUC 3'

cAMP (88 nt): SEQ ID NO: 137

5' thiol- GGAUAAUAGCCGUAGGUUGCGAAAGCGACCCUGAUGA  
CCUUGCGAUGCAAAAAGGUGCUGACGACACAU  
GAAACGGUAGCGAGAGCUC 3'

SEQ ID NO: 137




FIGURE 68

cCMP ((71 nt)): SEQ ID NO: 138

5' GGGACCCUGAUGAG CCUUUAGGGCCAAGUGUGGUGAAAGACACACGU  
CGAAACGGUGAAAGCCGUAGGUCC-Biotin 3'

cAMP ((70 nt)): SEQ ID NO: 103

5' GGGACCCUGAUGAG CCUGUGGAAACAGACGUGGCACAUGACUACGU  
CGAAACGGUGAAAGCCGUAGGUCC-Biotin 3'

cGMP (70 nt): SEQ ID NO: 139

5' GGGACCCUGAUGAG CCUUGCGAUGCAAAAAGGUGCUGACGACACAU  
CGAAACGGUGAAAGCCGUAGGUCC-Biotin 3'

SEQ ID NO: 103




FIGURE 69



cCMP: SEQ ID NO: 40

5' GGACCCUGAUGAG CCUUUAGGGCCAAGUGUGGUGAAAGACACACGU  
CGAAACGGUGAAAGCCGUAGGUCCUUGCGUGGUUCUGUUCCCUUCUUCG 3'

cAMP: SEQ ID NO: 41

5' GGACCCUGAUGAG CCUGUGGAAACAGACGUGGCACAUGACUACGU  
CGAAACGGUGAAAGCCGUAGGUCCUUGCGUGGUUCUGUUCCCUUCUUCG 3'

cGMP: SEQ ID NO: 42

5' GGACCCUGAUGAG CCUUGCGAUGCAAAAAGGUGCUGACGACACAU  
CGAAACGGUGAAAGCCGUAGGUCCUUGCGUGGUUCUGUUCCCUUCUUCG 3'

Capture Oligo: SEQ ID NO: 43

3'-ACGCACCAAGACAAGGGAAGAAGC-Biotin-5'








FIGURE 70

**A.** cGMP HH Binding to  
neutravidin-gold surface



$$K_{ads} \sim 0.41 \text{ min}^{-1}$$

**B.** cGMP HH cleavage  
with 1 mM cGMP



$$K_{dv} \sim 1.31 \text{ min}^{-1}$$

**FIGURE 71**




FIGURE 72




FIGURE 73





FIGURE 74


**Figure 75**

**B**

Hybridized & Washed



Ligated & Denatured






FIGURE 76





FIGURE 77



**A.****Captured Cy3-ligase:  
100% signal****TSA-Cy5 substrate:  
ligation signal****B.****FIGURE 78**




FIGURE 79




FIGURE 80





FIGURE 81

Figure 82



## SEQUENCE LISTING

<110> Archemix Corporation  
Stanton, Martin  
Epstein, David  
Hamaguchi, Nobuko  
Kurz, Markus  
Keefe, Tony  
Wilson, Charles  
Grate, Dilara  
Marshall, Kristin  
McCauley, Thomas  
Kurz, Jeffrey

<120> NUCLEIC ACID SENSOR MOLECULES AND METHODS OF USING SAME

<130> 23239-501-061 CIP

<140> Not yet assigned

<141> 2002-08-09

<150> 60/232,454

<151> 2000-09-13

<150> 09/952,680

<151> 2001-09-13

<150> 60/311,378

<151> 2001-08-09

<150> 60/313,932

<151> 2001-08-21

<150> 60/338,186

<151> 2001-11-13

<150> 60/349,959

<151> 2002-01-18

<150> 60/364,486

<151> 2002-03-13

<150> 60/376,744

<151> 2002-05-01

<150> 60/367,991

<151> 2002-03-25

<150> 60/369,887

<151> 2002-04-04

<150> 60/385,097

<151> 2002-05-31

<160> 372

<170> PatentIn Ver. 2.1

<210> 1

<211> 128

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 5'  
fixed:random:3'-fixed sequence

<220>

<221> misc\_feature

<222> (40)..(89)

<223> Wherein n is an a or u or c or g, and some may be  
missing.

<400> 1

ggacuucggu ccagugcucg ugcacuaggc cguucgaccn nnnnnnnnnn nnnnnnnnnn 60  
nnnnnnnnnn nnnnnnnnnn nnnnnnnnnc uuagacagga gguuaggugc cucgugaugu 120  
ccagucgc 128

<210> 2

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 3' primer  
sequence

<400> 2

gcgactggac atcacgag

18

<210> 3

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: substrate  
primer

<400> 3

aaaaaatgca ctggact

17

<210> 4

<211> 103

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: 5'  
fixed:random:3' fixed A-3'-fixed B molecule

<220>

<221> misc\_feature

<222> (22)..(71)

<223> Wherein n is an a or u or c or g and some may be  
missing.

<400> 4

gggcgaccu gaugagccug gnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 60

nnnnnnnnnn nuuagacgaa acggugaaag ccguagguug ccc 103

<210> 5  
 <211> 132  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: CW45-33-B04  
 ERK specific nucleic acid sensor molecule

<400> 5  
 ggacuucggc gaaagccguu cgaccgcaug cagacgcuag cgaauugguu ccucgaaagg 60  
 gaaagcguaa uuaagaaacc aaaaugcgug ucuuagacag gagguuaggu gcgucaaugc 120  
 ugcaaguac ug 132

<210> 6  
 <211> 133  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: CW45-33-D04  
 ERK specific nucleic acid sensor molecule

<400> 6  
 ggacuucggc gaaagccguu cgaccgguag cagacgcuag cgaauugguu ccucgaaagg 60  
 gaaagcguaa uuaagaaacc aaaauguuau cgcuuagaca ggagguuagg ugcgucaaug 120  
 cugcaaguua cug 133

<210> 7  
 <211> 134  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: CW45-33-F03  
 ERK specific nucleic acid sensor molecule

<400> 7  
 ggacuucggc gaaagccguu cgaccgugug gcagacgcuu gcgaauuggu uccucgaaag 60  
 ggaaagcgua auuaagaaac caaaaugcgc aggcuuagac aggagguuag gugcucaau 120  
 gcugcaagu acug 134

<210> 8  
 <211> 133  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: CW45-33-D01  
 ERK specific nucleic acid sensor molecule

<400> 8  
 ggacuucggc gaaagccguu cgaccgcaa cagacgcuag cgaauugguu ccucgaaagg 60  
 gaaagcguaa uuaagaaacc aaaauggugu ggcuuagaca ggagguuagg ugcgucaaug 120  
 cugcaaguua cug 133



<210> 9  
 <211> 2092  
 <212> DNA  
 <213> Homo sapiens

<400> 9  
 gaattccaaa atttgtatgt ttcttgatt tttgatgaag gagaaatact gtaatgatca 60  
 ctgtttacac tatgtacact ttaggccagc cctttgtagc gttatacaaa ctgaaagcac 120  
 accggaccgc caggctccgc gggcagggcc ggggccagag ctgcgctgtc ggcgggacat 180  
 gcgctgcgtc gcctctaacc tcgggctgtg ctctttttcc aggtggcccg ccggtttctg 240  
 agccttctgc cctgcgggga caggtctgc accctgcccc cgccacgga ccatgaccat 300  
 gaccctccac accaaaagcat ctgggatggc cctactgcat cagatccaag ggaacgagct 360  
 ggagccccctg aaccgtccgc agtcaagat cccctggag cggccccctg gcgaggtgta 420  
 cctggacagc agcaagcccg ccgtgtacaa ctaccccgag ggcgcgcctt acgagttcaa 480  
 cgccgcggcc gcgcgcaacg cgcaggtcta cggtcagacc ggcctccctt acggcccccg 540  
 gtctgaggct gcggcgcttc gctccaacgg cctggggggg tccccccac tcaacagcgt 600  
 gtctccgagc cgcctgatgc tactgcacc gccgcgcag ctgtcgctt tctgcagcc 660  
 ccacggccag caggtgccct actacctgga gaggaagagc tgccaggcct gccggctccg 720  
 ggccggcccg ccggcattct acaggccaaa ttcagataat cgacgccagg gtggcagaga 780  
 aagattggcc agtaccaatg acaaggaag tatggctatg gaatctgcca aggagactcg 840  
 ctactgtgca gtgtgcaatg actatgcttc aggtaccat tatggagtct ggtcctgtga 900  
 gggctgcaag gccttcttca agagaagtat tcaaggacat aacgactata tgtgtccagc 960  
 caccaaccag tgcaccattg ataaaaacag gaggaagagc tgccaggcct gccggctccg 1020  
 caaatgctac gaagtgggaa tgatgaaagg tgggatacga aaagaccgaa gaggagggag 1080  
 aatgttgaia cacaagcgcc agagagatga tggggagggc aggggtgaag tggggtctgc 1140  
 tggagacatg agagctgcca acctttggcc aagcccgctc atgatcaaac gctctaagaa 1200  
 gaacagcctg gccttctccc tgacggccga ccagatggtc agtgccttgt tggatgctga 1260  
 gcccccata ctctattccg agtatgatcc taccagaccc ttcagtgaag cttecatgat 1320  
 gggcttactg accaactctg cagacaggga gctgggtcac atgatcaact gggcgaagag 1380  
 ggtgccaggc tttgtggatt tgaccteca tgatcaggtc caccttctag aatgtgcctg 1440  
 gctagagatc ctgatgattg gtctcgtctg gcgctccatg gagcaccag tgaagctact 1500  
 gtttgcctct aacttgcctt tggacaggaa ccagggaiaa tgtgtagagg gcatgggtga 1560  
 gatcttcgac atgctgctgg ctacatcatc tcggttccgc atgatgaatc tgcagggaga 1620  
 ggagtttgtg tgcctcaaat ctattatctt gcttaattct ggagtgtaca catttctgtc 1680  
 cagcaccctg aagctctctg aagagaagga ccatatccac cgagtccctg acaagatcac 1740  
 agacactttg atccactga tggccaaggc aggcctgacc ctgcagcagc agcaccagcg 1800  
 gctggcccag ctctctctca tctctctcca catcaggcac atgagtaaca aaggcatgga 1860  
 gcatctgtac agcatgaagt gcaagaacgt ggtgccctc tatgacctgc tgcaggagat 1920  
 gctggacgcc caccgcctac atgcgccac tagccgtgga ggggcatccg tggaggagac 1980  
 ggaccaaagc cacttgccca ctgcgggctc tacttcatcg cattccttgc aaaagtatta 2040  
 catcacgggg gaggcagagg gtttccctgc cacagtctga gagtccctg gc 2092

<210> 10  
 <211> 595  
 <212> PRT  
 <213> Homo sapiens

<400> 10  
 Met Thr Met Thr Leu His Thr Lys Ala Ser Gly Met Ala Leu Leu His  
 1 5 10 15  
 Gln Ile Gln Gly Asn Glu Leu Glu Pro Leu Asn Arg Pro Gln Leu Lys  
 20 25 30  
 Ile Pro Leu Glu Arg Pro Leu Gly Glu Val Tyr Leu Asp Ser Ser Lys  
 35 40 45  
 Pro Ala Val Tyr Asn Tyr Pro Glu Gly Ala Ala Tyr Glu Phe Asn Ala

50                      55                      60  
 Ala Ala Ala Ala Asn Ala Gln Val Tyr Gly Gln Thr Gly Leu Pro Tyr  
 65                      70                      75                      80  
 Gly Pro Gly Ser Glu Ala Ala Ala Phe Gly Ser Asn Gly Leu Gly Gly  
 85                      90                      95  
 Phe Pro Pro Leu Asn Ser Val Ser Pro Ser Pro Leu Met Leu Leu His  
 100                      105                      110  
 Pro Pro Pro Gln Leu Ser Pro Phe Leu Gln Pro His Gly Gln Gln Val  
 115                      120                      125  
 Pro Tyr Tyr Leu Glu Asn Glu Pro Ser Gly Tyr Thr Val Arg Glu Ala  
 130                      135                      140  
 Gly Pro Pro Ala Phe Tyr Arg Pro Asn Ser Asp Asn Arg Arg Gln Gly  
 145                      150                      155                      160  
 Gly Arg Glu Arg Leu Ala Ser Thr Asn Asp Lys Gly Ser Met Ala Met  
 165                      170                      175  
 Glu Ser Ala Lys Glu Thr Arg Tyr Cys Ala Val Cys Asn Asp Tyr Ala  
 180                      185                      190  
 Ser Gly Tyr His Tyr Gly Val Trp Ser Cys Glu Gly Cys Lys Ala Phe  
 195                      200                      205  
 Phe Lys Arg Ser Ile Gln Gly His Asn Asp Tyr Met Cys Pro Ala Thr  
 210                      215                      220  
 Asn Gln Cys Thr Ile Asp Lys Asn Arg Arg Lys Ser Cys Gln Ala Cys  
 225                      230                      235                      240  
 Arg Leu Arg Lys Cys Tyr Glu Val Gly Met Met Lys Gly Gly Ile Arg  
 245                      250                      255  
 Lys Asp Arg Arg Gly Gly Arg Met Leu Lys His Lys Arg Gln Arg Asp  
 260                      265                      270  
 Asp Gly Glu Gly Arg Gly Glu Val Gly Ser Ala Gly Asp Met Arg Ala  
 275                      280                      285  
 Ala Asn Leu Trp Pro Ser Pro Leu Met Ile Lys Arg Ser Lys Lys Asn  
 290                      295                      300  
 Ser Leu Ala Leu Ser Leu Thr Ala Asp Gln Met Val Ser Ala Leu Leu  
 305                      310                      315                      320  
 Asp Ala Glu Pro Pro Ile Leu Tyr Ser Glu Tyr Asp Pro Thr Arg Pro  
 325                      330                      335  
 Phe Ser Glu Ala Ser Met Met Gly Leu Leu Thr Asn Leu Ala Asp Arg  
 340                      345                      350  
 Glu Leu Val His Met Ile Asn Trp Ala Lys Arg Val Pro Gly Phe Val  
 355                      360                      365  
 Asp Leu Thr Leu His Asp Gln Val His Leu Leu Glu Cys Ala Trp Leu  
 370                      375                      380

Glu Ile Leu Met Ile Gly Leu Val Trp Arg Ser Met Glu His Pro Val  
 385 390 395 400  
 Lys Leu Leu Phe Ala Pro Asn Leu Leu Leu Asp Arg Asn Gln Gly Lys  
 405 410 415  
 Cys Val Glu Gly Met Val Glu Ile Phe Asp Met Leu Leu Ala Thr Ser  
 420 425 430  
 Ser Arg Phe Arg Met Met Asn Leu Gln Gly Glu Glu Phe Val Cys Leu  
 435 440 445  
 Lys Ser Ile Ile Leu Leu Asn Ser Gly Val Tyr Thr Phe Leu Ser Ser  
 450 455 460  
 Thr Leu Lys Ser Leu Glu Glu Lys Asp His Ile His Arg Val Leu Asp  
 465 470 475 480  
 Lys Ile Thr Asp Thr Leu Ile His Leu Met Ala Lys Ala Gly Leu Thr  
 485 490 495  
 Leu Gln Gln Gln His Gln Arg Leu Ala Gln Leu Leu Leu Ile Leu Ser  
 500 505 510  
 His Ile Arg His Met Ser Asn Lys Gly Met Glu His Leu Tyr Ser Met  
 515 520 525  
 Lys Cys Lys Asn Val Val Pro Leu Tyr Asp Leu Leu Leu Glu Met Leu  
 530 535 540  
 Asp Ala His Arg Leu His Ala Pro Thr Ser Arg Gly Gly Ala Ser Val  
 545 550 555 560  
 Glu Glu Thr Asp Gln Ser His Leu Ala Thr Ala Gly Ser Thr Ser Ser  
 565 570 575  
 His Ser Leu Gln Lys Tyr Tyr Ile Thr Gly Glu Ala Glu Gly Phe Pro  
 580 585 590  
 Ala Thr Val  
 595

&lt;210&gt; 11

&lt;211&gt; 1970

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

gcaccgcgag cccctagcac cgcacaagct gagtgtgcag gacgagtcac caccacaccc 60  
 acaccacagc cgctgaatga ggcttccagg cgtccgctcg cggcccgag agccccgcg 120  
 tgggtccgcc cgctgaggcg cccccagcca gtgcgcttac ctgccagact gcgcgccatg 180  
 gggcaacccg ggaacggcag cgccttcttg ctggcaccca atggaagcca tgcgcgggac 240  
 cagcagctca cgcaggaaag ggacgaggtg tgggtggtgg gcatgggcat cgtcatgtct 300  
 ctcatcgctc tggccatcgt gtttggcaat gtgctggtca tcacagccat tgccaagtcc 360  
 gagcgtctgc agacggtcac caactacttc atcacttcac tggcctgtgc tgatctggtc 420  
 atgggcctgg cagtgggtgcc ctttggggcc gcccatattc ttatgaaaat gtggactttt 480  
 ggcaacttct ggtgcgagtt ttggacttcc attgatgtgc tgtgcgtcac ggccagcatt 540  
 gagaccctgt gcgtgatcgc agtggatcgc tactttgccca ttacttcacc tttcaagtac 600  
 cagagcctgc tgaccaagaa taaggcccg gtgatcattc tgatgggtgtg gattgtgtca 660

```

ggccttacct ccttcttgcc cattcagatg cactgggtacc gggccaccca ccaggaagcc 720
atcaactgct atgccaatga gacctgctgt gacttcttca cgaaccaagc ctatgccatt 780
gcctcttcca tcgtgtcctt ctacgttccc ctgggtgatca tgggtcttctg ctactccagg 840
gtcttttcagg aggcacaaag gcagctccag aagattgaca aatctgaggg ccgcttccat 900
gtccagaacc ttagccagggt ggagcaggat gggcgagcgg ggcagtgact ccgcagatct 960
tccaagttct gcttgaagga gcacaaagcc ctcaagacgt taggcatcat catgggcact 1020
ttcacccctc gctggctgcc cttcttcacg gttaacattg tgcatgtgat ccaggataac 1080
ctcatccgta aggaagttaa catcctccta aattggatag gctatgtcaa ttctggtttc 1140
aatcccccta tctactgccg gagcccgatg ttcaggattg ccttccagga gcttctgtgc 1200
ctgctcagggt cttctttgaa ggcctatggg aatggctact ccagcaacgg caacacaggg 1260
gagcagagtg gatatacagt ggaacaggag aaagaaaata aactgctgtg tgaagacctc 1320
ccaggcacgg aagactttgt gggccatcaa ggtactgtgc ctagegataa cattgattca 1380
caaggaggga attgtagtac aaatgactca ctgctgtaaa gcagtttttc tacttttaaa 1440
gacccccccc cccccacag aacactaaac agactattta acttgagggt aataaactta 1500
gaataaaatt gtaaaaattg tatagagata tgcagaagga agggcatcct tctgcctttt 1560
ttattttttt aagctgtaaa aagagagaaa acttatttga gtgattattt gttatttgta 1620
cagttcagtt cctctttgca tggaaattgt aagtttatgt ctaaagagct ttagtcctag 1680
aggacctgag tctgctatat ttccatgact ttccatgta tctacctcac tattcaagta 1740
ttaggggtaa tatattgctg gtaatttgta tctgaaggag attttccttc ctacacctt 1800
ggacttgagg attttgagta tctcggacct ttcagctgtg aacatggact cttccccac 1860
tcctcttatt tgctcacagc ggggtatttta ggcagggtt tgaggagcag cttcagttgt 1920
tttcccgagc aaaggctcaa agtttacagt aaataaaatg tttgaccatg 1970

```

&lt;210&gt; 12

&lt;211&gt; 413

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

```

Met Gly Gln Pro Gly Asn Gly Ser Ala Phe Leu Leu Ala Pro Asn Gly
  1                      5                      10                      15
Ser His Ala Pro Asp His Asp Val Thr Gln Gln Arg Asp Glu Val Trp
                      20                      25                      30
Val Val Gly Met Gly Ile Val Met Ser Leu Ile Val Leu Ala Ile Val
                      35                      40                      45
Phe Gly Asn Val Leu Val Ile Thr Ala Ile Ala Lys Phe Glu Arg Leu
                      50                      55                      60
Gln Thr Val Thr Asn Tyr Phe Ile Thr Ser Leu Ala Cys Ala Asp Leu
                      65                      70                      75                      80
Val Met Gly Leu Ala Val Val Pro Phe Gly Ala Ala His Ile Leu Met
                      85                      90                      95
Lys Met Trp Thr Phe Gly Asn Phe Trp Cys Glu Phe Trp Thr Ser Ile
                      100                      105                      110
Asp Val Leu Cys Val Thr Ala Ser Ile Glu Thr Leu Cys Val Ile Ala
                      115                      120                      125
Val Asp Arg Tyr Phe Ala Ile Thr Ser Pro Phe Lys Tyr Gln Ser Leu
                      130                      135                      140
Leu Thr Lys Asn Lys Ala Arg Val Ile Ile Leu Met Val Trp Ile Val
                      145                      150                      155                      160
Ser Gly Leu Thr Ser Phe Leu Pro Ile Gln Met His Trp Tyr Arg Ala

```

165 170 175  
 Thr His Gln Glu Ala Ile Asn Cys Tyr Ala Asn Glu Thr Cys Cys Asp  
 180 185 190  
 Phe Phe Thr Asn Gln Ala Tyr Ala Ile Ala Ser Ser Ile Val Ser Phe  
 195 200 205  
 Tyr Val Pro Leu Val Ile Met Val Phe Val Tyr Ser Arg Val Phe Gln  
 210 215 220  
 Glu Ala Lys Arg Gln Leu Gln Lys Ile Asp Lys Ser Glu Gly Arg Phe  
 225 230 235 240  
 His Val Gln Asn Leu Ser Gln Val Glu Gln Asp Gly Arg Thr Gly His  
 245 250 255  
 Gly Leu Arg Arg Ser Ser Lys Phe Cys Leu Lys Glu His Lys Ala Leu  
 260 265 270  
 Lys Thr Leu Gly Ile Ile Met Gly Thr Phe Thr Leu Cys Trp Leu Pro  
 275 280 285  
 Phe Phe Ile Val Asn Ile Val His Val Ile Gln Asp Asn Leu Ile Arg  
 290 295 300  
 Lys Glu Val Tyr Ile Leu Leu Asn Trp Ile Gly Tyr Val Asn Ser Gly  
 305 310 315 320  
 Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe Arg Ile Ala Phe  
 325 330 335  
 Gln Glu Leu Leu Cys Leu Arg Arg Ser Ser Leu Lys Ala Tyr Gly Asn  
 340 345 350  
 Gly Tyr Ser Ser Asn Gly Asn Thr Gly Glu Gln Ser Gly Tyr His Val  
 355 360 365  
 Glu Gln Glu Lys Glu Asn Lys Leu Leu Cys Glu Asp Leu Pro Gly Thr  
 370 375 380  
 Glu Asp Phe Val Gly His Gln Gly Thr Val Pro Ser Asp Asn Ile Asp  
 385 390 395 400  
 Ser Gln Gly Arg Asn Cys Ser Thr Asn Asp Ser Leu Leu  
 405 410

<210> 13  
 <211> 394  
 <212> PRT  
 <213> Homo sapiens

<400> 13  
 Met Gly Cys Leu Gly Asn Ser Lys Thr Glu Asp Gln Arg Asn Glu Glu  
 1 5 10 15  
 Lys Ala Gln Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys  
 20 25 30  
 Asp Lys Gln Val Tyr Arg Ala Thr His Arg Leu Leu Leu Leu Gly Ala

35 40 45  
 Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His  
 50 55 60  
 Val Asn Gly Phe Asn Gly Gly Gly Glu Glu Asp Pro Gln Ala Ala  
 65 70 75 80  
 Arg Ser Asn Ser Asp Gly Glu Lys Ala Thr Lys Val Gln Asp Ile Lys  
 85 90 95  
 Asn Asn Leu Lys Glu Ala Ile Glu Thr Ile Val Ala Ala Met Ser Asn  
 100 105 110  
 Leu Val Pro Pro Val Glu Leu Ala Asn Pro Glu Asn Gln Phe Arg Val  
 115 120 125  
 Asp Tyr Ile Leu Ser Val Met Asn Val Pro Asp Phe Asp Phe Pro Pro  
 130 135 140  
 Glu Phe Tyr Glu His Ala Lys Ala Leu Trp Glu Asp Glu Gly Val Arg  
 145 150 155 160  
 Ala Cys Tyr Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala Gln  
 165 170 175  
 Tyr Phe Leu Asp Lys Ile Asp Val Ile Lys Gln Ala Asp Tyr Val Pro  
 180 185 190  
 Ser Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile Phe  
 195 200 205  
 Glu Thr Lys Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp Val  
 210 215 220  
 Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn Asp  
 225 230 235 240  
 Val Thr Ala Ile Ile Phe Val Val Ala Ser Ser Ser Tyr Asn Met Val  
 245 250 255  
 Ile Arg Glu Asp Asn Gln Thr Asn Arg Leu Gln Glu Ala Leu Asn Leu  
 260 265 270  
 Phe Lys Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Val Ile  
 275 280 285  
 Leu Phe Leu Asn Lys Gln Asp Leu Leu Ala Glu Lys Val Leu Ala Gly  
 290 295 300  
 Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Phe Ala Arg Tyr Thr Thr  
 305 310 315 320  
 Pro Glu Asp Ala Thr Pro Glu Pro Gly Glu Asp Pro Arg Val Thr Arg  
 325 330 335  
 Ala Lys Tyr Phe Ile Arg Asp Glu Phe Leu Arg Ile Ser Thr Ala Ser  
 340 345 350  
 Gly Asp Gly Arg His Tyr Cys Tyr Pro His Phe Thr Cys Ala Val Asp  
 355 360 365

Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile Ile Gln  
 370 375 380

Arg Met His Leu Arg Gln Tyr Glu Leu Leu  
 385 390

<210> 14  
 <211> 395  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Met Gly Cys Leu Gly Asn Ser Lys Thr Glu Asp Gln Arg Asn Glu Glu  
 1 5 10 15

Lys Ala Gln Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys  
 20 25 30

Asp Lys Gln Val Tyr Arg Ala Thr His Arg Leu Leu Leu Leu Gly Ala  
 35 40 45

Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His  
 50 55 60

Val Asn Gly Phe Asn Gly Glu Gly Glu Glu Asp Pro Gln Ala Ala  
 65 70 75 80

Arg Ser Asn Ser Asp Gly Ser Glu Lys Ala Thr Lys Val Gln Asp Ile  
 85 90 95

Lys Asn Asn Leu Lys Glu Ala Ile Glu Thr Ile Val Ala Ala Met Ser  
 100 105 110

Asn Leu Val Pro Pro Val Glu Leu Ala Asn Pro Glu Asn Gln Phe Arg  
 115 120 125

Val Asp Tyr Ile Leu Ser Val Met Asn Val Pro Asp Phe Asp Phe Pro  
 130 135 140

Pro Glu Phe Tyr Glu His Ala Lys Ala Leu Trp Glu Asp Glu Gly Val  
 145 150 155 160

Arg Ala Cys Tyr Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala  
 165 170 175

Gln Tyr Phe Leu Asp Lys Ile Asp Val Ile Lys Gln Ala Asp Tyr Val  
 180 185 190

Pro Ser Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile  
 195 200 205

Phe Glu Thr Lys Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp  
 210 215 220

Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn  
 225 230 235 240

Asp Val Thr Ala Ile Ile Phe Val Val Ala Ser Ser Ser Tyr Asn Met  
 245 250 255

Val Ile Arg Glu Asp Asn Gln Thr Asn Arg Leu Gln Glu Ala Leu Asn  
 260 265 270  
 Leu Phe Lys Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Val  
 275 280 285  
 Ile Leu Phe Leu Asn Lys Gln Asp Leu Leu Ala Glu Lys Val Leu Ala  
 290 295 300  
 Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Phe Ala Arg Tyr Thr  
 305 310 315 320  
 Thr Pro Glu Asp Ala Thr Pro Glu Pro Gly Glu Asp Pro Arg Val Thr  
 325 330 335  
 Arg Ala Lys Tyr Phe Ile Arg Asp Glu Phe Leu Arg Ile Ser Thr Ala  
 340 345 350  
 Ser Gly Asp Gly Arg His Tyr Cys Tyr Pro His Phe Thr Cys Ala Val  
 355 360 365  
 Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile Ile  
 370 375 380  
 Gln Arg Met His Leu Arg Gln Tyr Glu Leu Leu  
 385 390 395

<210> 15  
 <211> 379  
 <212> PRT  
 <213> Homo sapiens

<400> 15  
 Met Gly Cys Leu Gly Asn Ser Lys Thr Glu Asp Gln Arg Asn Glu Glu  
 1 5 10 15  
 Lys Ala Gln Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys  
 20 25 30  
 Asp Lys Gln Val Tyr Arg Ala Thr His Arg Leu Leu Leu Leu Gly Ala  
 35 40 45  
 Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His  
 50 55 60  
 Val Asn Gly Phe Asn Gly Asp Glu Lys Ala Thr Lys Val Gln Asp Ile  
 65 70 75 80  
 Lys Asn Asn Leu Lys Glu Ala Ile Glu Thr Ile Val Ala Ala Met Ser  
 85 90 95  
 Asn Leu Val Pro Pro Val Glu Leu Ala Asn Pro Glu Asn Gln Phe Arg  
 100 105 110  
 Val Asp Tyr Ile Leu Ser Val Met Asn Val Pro Asp Phe Asp Phe Pro  
 115 120 125  
 Pro Glu Phe Tyr Glu His Ala Lys Ala Leu Trp Glu Asp Glu Gly Val  
 130 135 140



Arg Ala Cys Tyr Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala  
 145 150 155 160  
 Gln Tyr Phe Leu Asp Lys Ile Asp Val Ile Lys Gln Ala Asp Tyr Val  
 165 170 175  
 Pro Ser Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile  
 180 185 190  
 Phe Glu Thr Lys Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp  
 195 200 205  
 Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn  
 210 215 220  
 Asp Val Thr Ala Ile Ile Phe Val Val Ala Ser Ser Ser Tyr Asn Met  
 225 230 235 240  
 Val Ile Arg Glu Asp Asn Gln Thr Asn Arg Leu Gln Glu Ala Leu Asn  
 245 250 255  
 Leu Phe Lys Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Val  
 260 265 270  
 Ile Leu Phe Leu Asn Lys Gln Asp Leu Leu Ala Glu Lys Val Leu Ala  
 275 280 285  
 Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Phe Ala Arg Tyr Thr  
 290 295 300  
 Thr Pro Glu Asp Ala Thr Pro Glu Pro Gly Glu Asp Pro Arg Val Thr  
 305 310 315 320  
 Arg Ala Lys Tyr Phe Ile Arg Asp Glu Phe Leu Arg Ile Ser Thr Ala  
 325 330 335  
 Ser Gly Asp Gly Arg His Tyr Cys Tyr Pro His Phe Thr Cys Ala Val  
 340 345 350  
 Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile Ile  
 355 360 365  
 Gln Arg Met His Leu Arg Gln Tyr Glu Leu Leu  
 370 375

&lt;210&gt; 16

&lt;211&gt; 380

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

Met Gly Cys Leu Gly Asn Ser Lys Thr Glu Asp Gln Arg Asn Glu Glu  
 1 5 10 15  
 Lys Ala Gln Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu Gln Lys  
 20 25 30  
 Asp Lys Gln Val Tyr Arg Ala Thr His Arg Leu Leu Leu Leu Gly Ala  
 35 40 45

Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His  
 50 55 60  
 Val Asn Gly Phe Asn Gly Asp Ser Glu Lys Ala Thr Lys Val Gln Asp  
 65 70 75 80  
 Ile Lys Asn Asn Leu Lys Glu Ala Ile Glu Thr Ile Val Ala Ala Met  
 85 90 95  
 Ser Asn Leu Val Pro Pro Val Glu Leu Ala Asn Pro Glu Asn Gln Phe  
 100 105 110  
 Arg Val Asp Tyr Ile Leu Ser Val Met Asn Val Pro Asp Phe Asp Phe  
 115 120 125  
 Pro Pro Glu Phe Tyr Glu His Ala Lys Ala Leu Trp Glu Asp Glu Gly  
 130 135 140  
 Val Arg Ala Cys Tyr Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys  
 145 150 155 160  
 Ala Gln Tyr Phe Leu Asp Lys Ile Asp Val Ile Lys Gln Ala Asp Tyr  
 165 170 175  
 Val Pro Ser Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly  
 180 185 190  
 Ile Phe Glu Thr Lys Phe Gln Val Asp Lys Val Asn Phe His Met Phe  
 195 200 205  
 Asp Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe  
 210 215 220  
 Asn Asp Val Thr Ala Ile Ile Phe Val Val Ala Ser Ser Ser Tyr Asn  
 225 230 235 240  
 Met Val Ile Arg Glu Asp Asn Gln Thr Asn Arg Leu Gln Glu Ala Leu  
 245 250 255  
 Asn Leu Phe Lys Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser  
 260 265 270  
 Val Ile Leu Phe Leu Asn Lys Gln Asp Leu Leu Ala Glu Lys Val Leu  
 275 280 285  
 Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Phe Ala Arg Tyr  
 290 295 300  
 Thr Thr Pro Glu Asp Ala Thr Pro Glu Pro Gly Glu Asp Pro Arg Val  
 305 310 315 320  
 Thr Arg Ala Lys Tyr Phe Ile Arg Asp Glu Phe Leu Arg Ile Ser Thr  
 325 330 335  
 Ala Ser Gly Asp Gly Arg His Tyr Cys Tyr Pro His Phe Thr Cys Ala  
 340 345 350  
 Val Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile  
 355 360 365

Ile Gln Arg Met His Leu Arg Gln Tyr Glu Leu Leu  
 370 375 380

<210> 17

<211> 388

<212> PRT

<213> Homo sapiens

<400> 17

Met Glu Ile Ser Gly Pro Pro Phe Glu Ile Gly Ser Ala Pro Ala Gly  
 1 5 10 15

Val Asp Asp Thr Pro Val Asn Met Asp Ser Pro Pro Ile Ala Leu Asp  
 20 25 30

Gly Pro Pro Ile Lys Val Ser Gly Ala Pro Asp Lys Arg Glu Arg Ala  
 35 40 45

Glu Arg Pro Pro Val Glu Glu Glu Ala Ala Glu Met Glu Gly Ala Ala  
 50 55 60

Asp Ala Ala Glu Gly Gly Lys Val Pro Ser Pro Gly Tyr Gly Ser Pro  
 65 70 75 80

Ala Ala Gly Ala Ala Ser Ala Asp Thr Ala Ala Arg Ala Ala Pro Ala  
 85 90 95

Ala Pro Ala Asp Pro Asp Ser Gly Ala Thr Pro Glu Asp Pro Asp Ser  
 100 105 110

Gly Thr Ala Pro Ala Asp Pro Asp Ser Gly Ala Phe Ala Ala Asp Pro  
 115 120 125

Asp Ser Gly Ala Ala Pro Ala Ala Pro Ala Asp Pro Asp Ser Gly Ala  
 130 135 140

Ala Pro Asp Ala Pro Ala Asp Pro Asp Ser Gly Ala Ala Pro Asp Ala  
 145 150 155 160

Pro Ala Asp Pro Asp Ala Gly Ala Ala Pro Glu Ala Pro Ala Ala Pro  
 165 170 175

Ala Ala Ala Glu Thr Arg Ala Ala His Val Ala Pro Ala Ala Pro Asp  
 180 185 190

Ala Gly Ala Pro Thr Ala Pro Ala Ala Ser Ala Thr Arg Ala Ala Gln  
 195 200 205

Val Arg Arg Ala Ala Ser Ala Ala Pro Ala Ser Gly Ala Arg Arg Lys  
 210 215 220

Ile His Leu Arg Pro Pro Ser Pro Glu Ile Gln Ala Ala Asp Pro Pro  
 225 230 235 240

Thr Pro Arg Pro Thr Arg Ala Ser Ala Trp Arg Gly Lys Ser Glu Ser  
 245 250 255

Ser Arg Gly Arg Arg Val Tyr Tyr Asp Glu Gly Val Ala Ser Ser Asp  
 260 265 270

Asp Asp Ser Ser Gly Asp Glu Ser Asp Asp Gly Thr Ser Gly Cys Leu  
 275 280 285  
 Arg Trp Phe Gln His Arg Arg Asn Arg Arg Arg Arg Lys Pro Gln Arg  
 290 295 300  
 Asn Leu Leu Arg Asn Phe Leu Val Gln Ala Phe Gly Gly Cys Phe Gly  
 305 310 315 320  
 Arg Ser Glu Ser Pro Gln Pro Lys Ala Ser Arg Ser Leu Lys Val Lys  
 325 330 335  
 Lys Val Pro Leu Ala Glu Lys Arg Arg Gln Met Arg Lys Glu Ala Leu  
 340 345 350  
 Glu Lys Arg Ala Gln Lys Arg Ala Glu Lys Lys Arg Ser Lys Leu Ile  
 355 360 365  
 Asp Lys Gln Leu Gln Asp Glu Lys Met Gly Tyr Met Cys Thr His Arg  
 370 375 380  
 Leu Leu Leu Leu  
 385

<210> 18  
 <211> 381  
 <212> PRT  
 <213> Homo sapiens

<400> 18  
 Met Gly Cys Leu Gly Gly Asn Ser Lys Thr Thr Glu Asp Gln Gly Val  
 1 5 10 15  
 Asp Glu Lys Glu Arg Arg Glu Ala Asn Lys Lys Ile Glu Lys Gln Leu  
 20 25 30  
 Gln Lys Glu Arg Leu Ala Tyr Lys Ala Thr His Arg Leu Leu Leu Leu  
 35 40 45  
 Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile  
 50 55 60  
 Leu His Val Asn Gly Phe Asn Pro Glu Glu Lys Lys Gln Lys Ile Leu  
 65 70 75 80  
 Asp Ile Arg Lys Asn Val Lys Asp Ala Ile Val Thr Ile Val Ser Ala  
 85 90 95  
 Met Ser Thr Ile Ile Pro Pro Val Pro Leu Ala Asn Pro Glu Asn Gln  
 100 105 110  
 Phe Arg Ser Asp Tyr Ile Lys Ser Ile Ala Pro Ile Thr Asp Phe Glu  
 115 120 125  
 Tyr Ser Gln Glu Phe Phe Asp His Val Lys Lys Leu Trp Asp Asp Glu  
 130 135 140  
 Gly Val Lys Ala Cys Phe Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp  
 145 150 155 160

Cys Ala Gln Tyr Phe Leu Glu Arg Ile Asp Ser Val Ser Leu Val Asp  
 165 170 175  
 Tyr Thr Pro Thr Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser  
 180 185 190  
 Gly Ile Phe Glu Thr Arg Phe Gln Val Asp Lys Val Asn Phe His Met  
 195 200 205  
 Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys  
 210 215 220  
 Phe Asn Asp Val Thr Ala Ile Ile Tyr Val Ala Ala Cys Ser Ser Tyr  
 225 230 235 240  
 Asn Met Val Ile Arg Glu Asp Asn Asn Thr Asn Arg Leu Arg Glu Ser  
 245 250 255  
 Leu Asp Leu Phe Glu Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile  
 260 265 270  
 Ser Ile Ile Leu Phe Leu Asn Lys Gln Asp Met Leu Ala Glu Lys Val  
 275 280 285  
 Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Tyr Ala Asn  
 290 295 300  
 Tyr Thr Val Pro Glu Asp Ala Thr Pro Asp Ala Gly Glu Asp Pro Lys  
 305 310 315 320  
 Val Thr Arg Ala Lys Phe Phe Ile Arg Asp Leu Phe Leu Arg Ile Ser  
 325 330 335  
 Thr Ala Thr Gly Asp Gly Lys His Tyr Cys Tyr Pro His Phe Thr Cys  
 340 345 350  
 Ala Val Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp  
 355 360 365  
 Ile Ile Gln Arg Met His Leu Lys Gln Tyr Glu Leu Leu  
 370 375 380

&lt;210&gt; 19

&lt;211&gt; 354

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

Met Gly Cys Thr Leu Ser Ala Glu Asp Lys Ala Ala Val Glu Arg Ser  
 1 5 10 15  
 Lys Met Ile Asp Arg Asn Leu Arg Glu Asp Gly Glu Lys Ala Ala Arg  
 20 25 30  
 Glu Val Lys Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr  
 35 40 45  
 Ile Val Lys Gln Met Lys Ile Ile His Glu Ala Gly Tyr Ser Glu Glu  
 50 55 60

Glu Cys Lys Gln Tyr Lys Ala Val Val Tyr Ser Asn Thr Ile Gln Ser  
 65 70 75 80  
 Ile Ile Ala Ile Ile Arg Ala Met Gly Arg Leu Lys Ile Asp Phe Gly  
 85 90 95  
 Asp Ser Ala Arg Ala Asp Asp Ala Arg Gln Leu Phe Val Leu Ala Gly  
 100 105 110  
 Ala Ala Glu Glu Gly Phe Met Thr Ala Glu Leu Ala Gly Val Ile Lys  
 115 120 125  
 Arg Leu Trp Lys Asp Ser Gly Val Gln Ala Cys Phe Asn Arg Ser Arg  
 130 135 140  
 Glu Tyr Gln Leu Asn Asp Ser Ala Ala Tyr Tyr Leu Asn Asp Leu Asp  
 145 150 155 160  
 Arg Ile Ala Gln Pro Asn Tyr Ile Pro Thr Gln Gln Asp Val Leu Arg  
 165 170 175  
 Thr Arg Val Lys Thr Thr Gly Ile Val Glu Thr His Phe Thr Phe Lys  
 180 185 190  
 Asp Leu His Phe Lys Met Phe Asp Val Gly Gly Gln Arg Ser Glu Arg  
 195 200 205  
 Lys Lys Trp Ile His Cys Phe Glu Gly Val Thr Ala Ile Ile Phe Cys  
 210 215 220  
 Val Ala Leu Ser Asp Tyr Asp Leu Val Leu Ala Glu Asp Glu Glu Met  
 225 230 235 240  
 Asn Arg Met His Glu Ser Met Lys Leu Phe Asp Ser Ile Cys Asn Asn  
 245 250 255  
 Lys Trp Phe Thr Asp Thr Ser Ile Ile Leu Phe Leu Asn Lys Lys Asp  
 260 265 270  
 Leu Phe Glu Glu Lys Ile Lys Lys Ser Pro Leu Thr Ile Cys Tyr Pro  
 275 280 285  
 Glu Tyr Ala Gly Ser Asn Thr Tyr Glu Glu Ala Ala Ala Tyr Ile Gln  
 290 295 300  
 Cys Gln Phe Glu Asp Leu Asn Lys Arg Lys Asp Thr Lys Glu Ile Tyr  
 305 310 315 320  
 Thr His Phe Thr Cys Ala Thr Asp Thr Lys Asn Val Gln Phe Val Phe  
 325 330 335  
 Asp Ala Val Thr Asp Val Ile Ile Lys Asn Asn Leu Lys Asp Cys Gly  
 340 345 350  
 Leu Phe

<210> 20  
 <211> 355  
 <212> PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

Met Gly Cys Thr Val Ser Ala Glu Asp Lys Ala Ala Ala Glu Arg Ser  
 1 5 10 15  
 Lys Met Ile Asp Lys Asn Leu Arg Glu Asp Gly Glu Lys Ala Ala Arg  
 20 25 30  
 Glu Val Lys Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr  
 35 40 45  
 Ile Val Lys Gln Met Lys Ile Ile His Glu Asp Gly Tyr Ser Glu Glu  
 50 55 60  
 Glu Cys Arg Gln Tyr Arg Ala Val Val Tyr Ser Asn Thr Ile Gln Ser  
 65 70 75 80  
 Ile Met Ala Ile Val Lys Ala Met Gly Asn Leu Gln Ile Asp Phe Ala  
 85 90 95  
 Asp Pro Ser Arg Ala Asp Asp Ala Arg Gln Leu Phe Ala Leu Ser Cys  
 100 105 110  
 Thr Ala Glu Glu Gln Gly Val Leu Pro Asp Asp Leu Ser Gly Val Ile  
 115 120 125  
 Arg Arg Leu Trp Ala Asp His Gly Val Gln Ala Cys Phe Gly Arg Ser  
 130 135 140  
 Arg Glu Tyr Gln Leu Asn Asp Ser Ala Ala Tyr Tyr Leu Asn Asp Leu  
 145 150 155 160  
 Glu Arg Ile Ala Gln Ser Asp Tyr Ile Pro Thr Gln Gln Asp Val Leu  
 165 170 175  
 Arg Thr Arg Val Lys Thr Thr Gly Ile Val Glu Thr His Phe Thr Phe  
 180 185 190  
 Lys Asp Leu His Phe Lys Met Phe Asp Val Gly Gly Gln Arg Ser Glu  
 195 200 205  
 Arg Lys Lys Trp Ile His Cys Phe Glu Gly Val Thr Ala Ile Ile Phe  
 210 215 220  
 Cys Val Ala Leu Ser Ala Tyr Asp Leu Val Leu Ala Glu Asp Glu Glu  
 225 230 235 240  
 Met Asn Arg Met His Glu Ser Met Lys Leu Phe Asp Ser Ile Cys Asn  
 245 250 255  
 Asn Lys Trp Phe Thr Asp Thr Ser Ile Ile Leu Phe Leu Asn Lys Lys  
 260 265 270  
 Asp Leu Phe Glu Glu Lys Ile Thr His Ser Pro Leu Thr Ile Cys Phe  
 275 280 285  
 Pro Glu Tyr Thr Gly Ala Asn Lys Tyr Asp Glu Ala Ala Ser Tyr Ile  
 290 295 300  
 Gln Ser Lys Phe Glu Asp Leu Asn Lys Arg Lys Asp Thr Lys Glu Ile

[illegible]

```
<210> 21
<211> 354
<212> PRT
<213> Homo sapiens
```

<400> 21																	
Met	Gly	Cys	Thr	Leu	Ser	Ala	Glu	Asp	Lys	Ala	Ala	Val	Glu	Arg	Ser		
1				5					10					15			
Lys	Met	Ile	Asp	Arg	Asn	Leu	Arg	Glu	Asp	Gly	Glu	Lys	Ala	Ala	Lys		
			20					25					30				
Glu	Val	Lys	Leu	Leu	Leu	Leu	Gly	Ala	Gly	Glu	Ser	Gly	Lys	Ser	Thr		
		35					40						45				
Ile	Val	Lys	Gln	Met	Lys	Ile	Ile	His	Glu	Asp	Gly	Tyr	Ser	Glu	Asp		
	50					55						60					
Glu	Cys	Lys	Gln	Tyr	Lys	Val	Val	Val	Tyr	Ser	Asn	Thr	Ile	Gln	Ser		
65					70					75							80
Ile	Ile	Ala	Ile	Ile	Arg	Ala	Met	Gly	Arg	Leu	Lys	Ile	Asp	Phe	Gly		
				85					90					95			
Glu	Ala	Ala	Arg	Ala	Asp	Asp	Ala	Arg	Gln	Leu	Phe	Val	Leu	Ala	Gly		
			100					105					110				
Ser	Ala	Glu	Glu	Gly	Val	Met	Thr	Pro	Glu	Leu	Ala	Gly	Val	Ile	Lys		
		115					120					125					
Arg	Leu	Trp	Arg	Asp	Gly	Gly	Val	Gln	Ala	Cys	Phe	Ser	Arg	Ser	Arg		
	130					135					140						
Glu	Tyr	Gln	Leu	Asn	Asp	Ser	Ala	Ser	Tyr	Tyr	Leu	Asn	Asp	Leu	Asp		
145					150					155					160		
Arg	Ile	Ser	Gln	Ser	Asn	Tyr	Ile	Pro	Thr	Gln	Gln	Asp	Val	Leu	Arg		
				165					170					175			
Thr	Arg	Val	Lys	Thr	Thr	Gly	Ile	Val	Glu	Thr	His	Phe	Thr	Phe	Lys		
			180					185					190				
Asp	Leu	Tyr	Phe	Lys	Met	Phe	Asp	Val	Gly	Gly	Gln	Arg	Ser	Glu	Arg		
		195					200					205					
Lys	Lys	Trp	Ile	His	Cys	Phe	Glu	Gly	Val	Thr	Ala	Ile	Ile	Phe	Cys		
	210					215					220						
Val	Ala	Leu	Ser	Asp	Tyr	Asp	Leu	Val	Leu	Ala	Glu	Asp	Glu	Glu	Met		



225                      230                      235                      240  
 Asn Arg Met His Glu Ser Met Lys Leu Phe Asp Ser Ile Cys Asn Asn  
                                  245                                   250                                   255  
 Lys Trp Phe Thr Glu Thr Ser Ile Ile Leu Phe Leu Asn Lys Lys Asp  
                                  260                                   265                                   270  
 Leu Phe Glu Glu Lys Ile Lys Arg Ser Pro Leu Thr Ile Cys Tyr Pro  
                                  275                                   280                                   285  
 Glu Tyr Thr Gly Ser Asn Thr Tyr Glu Glu Ala Ala Ala Tyr Ile Gln  
                                  290                                   295                                   300  
 Cys Gln Phe Glu Asp Leu Asn Arg Arg Lys Asp Thr Lys Glu Ile Tyr  
 305                                   310                                   315                                   320  
 Thr His Phe Thr Cys Ala Thr Asp Thr Lys Asn Val Gln Phe Val Phe  
                                  325                                   330                                   335  
 Asp Ala Val Thr Asp Val Ile Ile Lys Asn Asn Leu Lys Glu Cys Gly  
                                  340                                   345                                   350  
 Leu Tyr

<210> 22  
 <211> 354  
 <212> PRT  
 <213> Homo sapiens

<400> 22  
 Met Gly Cys Thr Leu Ser Ala Glu Glu Arg Ala Ala Leu Glu Arg Ser  
   1                                   5                                   10                                   15  
 Lys Ala Ile Glu Lys Asn Leu Lys Glu Asp Gly Ile Ser Ala Ala Lys  
                                  20                                   25                                   30  
 Asp Val Lys Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr  
                                  35                                   40                                   45  
 Ile Val Lys Gln Met Lys Ile Ile His Glu Asp Gly Phe Ser Gly Glu  
                                  50                                   55                                   60  
 Asp Val Lys Gln Tyr Lys Pro Val Val Tyr Ser Asn Thr Ile Gln Ser  
   65                                   70                                   75                                   80  
 Leu Ala Ala Ile Val Arg Ala Met Asp Thr Leu Gly Ile Glu Tyr Gly  
                                  85                                   90                                   95  
 Asp Lys Glu Arg Lys Ala Asp Ala Lys Met Val Cys Asp Val Val Ser  
                                  100                                   105                                   110  
 Arg Met Glu Asp Thr Glu Pro Phe Ser Ala Glu Leu Leu Ser Ala Met  
                                  115                                   120                                   125  
 Met Arg Leu Trp Gly Asp Ser Gly Ile Gln Glu Cys Phe Asn Arg Ser  
                                  130                                   135                                   140  
 Arg Glu Tyr Gln Leu Asn Asp Ser Ala Lys Tyr Tyr Leu Asp Ser Leu

145                      150                      155                      160  
 Asp Arg Ile Gly Ala Ala Asp Tyr Gln Pro Thr Glu Gln Asp Ile Leu  
                                  165                      170                      175  
 Arg Thr Arg Val Lys Thr Thr Gly Ile Val Glu Thr His Phe Thr Phe  
                                  180                      185                      190  
 Lys Asn Leu His Phe Arg Leu Phe Asp Val Gly Gly Gln Arg Ser Glu  
                                  195                      200                      205  
 Arg Lys Lys Trp Ile His Cys Phe Glu Asp Val Thr Ala Ile Ile Phe  
                                  210                      215                      220  
 Cys Val Ala Leu Ser Gly Tyr Asp Gln Val Leu His Glu Asp Glu Thr  
                                  225                      230                      235                      240  
 Thr Asn Arg Met His Glu Ser Leu Met Leu Phe Asp Ser Ile Cys Asn  
                                  245                      250                      255  
 Asn Lys Phe Phe Ile Asp Thr Ser Ile Ile Leu Phe Leu Asn Lys Lys  
                                  260                      265                      270  
 Asp Leu Phe Gly Glu Lys Ile Lys Lys Ser Pro Leu Thr Ile Cys Phe  
                                  275                      280                      285  
 Pro Glu Tyr Thr Gly Pro Asn Thr Tyr Glu Asp Ala Ala Ala Tyr Ile  
                                  290                      295                      300  
 Gln Ala Gln Phe Glu Ser Lys Asn Arg Ser Pro Asn Lys Glu Ile Tyr  
                                  305                      310                      315                      320  
 Cys His Met Thr Cys Ala Thr Asp Thr Asn Asn Ile Gln Val Val Phe  
                                  325                      330                      335  
 Asp Ala Val Thr Asp Ile Ile Ile Ala Asn Asn Leu Arg Gly Cys Gly  
                                  340                      345                      350  
 Leu Tyr

&lt;210&gt; 23

&lt;211&gt; 355

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

Met Gly Cys Thr Val Ser Ala Glu Asp Lys Ala Ala Ala Glu Arg Ser  
   1                                  5                                  10                                  15  
 Lys Met Ile Asp Lys Asn Leu Arg Glu Asp Gly Glu Lys Ala Ala Arg  
                                   20                                  25                                  30  
 Glu Val Lys Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr  
                                   35                                  40                                  45  
 Ile Val Lys Gln Met Lys Ile Ile His Glu Asp Gly Tyr Ser Glu Glu  
                                   50                                  55                                  60  
 Glu Cys Arg Gln Tyr Arg Ala Val Val Tyr Ser Asn Thr Ile Gln Ser

[illegible]

&lt;400&gt; 24

Met Gly Ala Gly Ala Ser Ala Glu Glu Lys His Ser Arg Glu Leu Glu  
 1 5 10 15  
 Lys Lys Leu Lys Glu Asp Ala Glu Lys Asp Ala Arg Thr Val Lys Leu  
 20 25 30  
 Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln  
 35 40 45  
 Met Lys Ile Ile His Gln Asp Gly Tyr Ser Leu Glu Glu Cys Leu Glu  
 50 55 60  
 Phe Ile Ala Ile Ile Tyr Gly Asn Thr Leu Gln Ser Ile Leu Ala Ile  
 65 70 75 80  
 Val Arg Ala Met Thr Thr Leu Asn Ile Gln Tyr Gly Asp Ser Ala Arg  
 85 90 95  
 Gln Asp Asp Ala Arg Lys Leu Met His Met Ala Asp Thr Ile Glu Glu  
 100 105 110  
 Gly Thr Met Pro Lys Glu Met Ser Asp Ile Ile Gln Arg Leu Trp Lys  
 115 120 125  
 Asp Ser Gly Ile Gln Ala Cys Phe Glu Arg Ala Ser Glu Tyr Gln Leu  
 130 135 140  
 Asn Asp Ser Ala Gly Tyr Tyr Leu Ser Asp Leu Glu Arg Leu Val Thr  
 145 150 155 160  
 Pro Gly Tyr Val Pro Thr Glu Gln Asp Val Leu Arg Ser Arg Val Lys  
 165 170 175  
 Thr Thr Gly Ile Ile Glu Thr Gln Phe Ser Phe Lys Asp Leu Asn Phe  
 180 185 190  
 Arg Met Phe Asp Val Gly Gly Gln Arg Ser Glu Arg Lys Lys Trp Ile  
 195 200 205  
 His Cys Phe Glu Gly Val Thr Cys Ile Ile Phe Ile Ala Ala Leu Ser  
 210 215 220  
 Ala Tyr Asp Met Val Leu Val Glu Asp Asp Glu Val Asn Arg Met His  
 225 230 235 240  
 Glu Ser Leu His Leu Phe Asn Ser Ile Cys Asn His Arg Tyr Phe Ala  
 245 250 255  
 Thr Thr Ser Ile Val Leu Phe Leu Asn Lys Lys Asp Val Phe Phe Glu  
 260 265 270  
 Lys Ile Lys Lys Ala His Leu Ser Ile Cys Phe Pro Asp Tyr Asp Gly  
 275 280 285  
 Pro Asn Thr Tyr Glu Asp Ala Gly Asn Tyr Ile Lys Val Gln Phe Leu  
 290 295 300  
 Glu Leu Asn Met Arg Arg Asp Val Lys Glu Ile Tyr Ser His Met Thr  
 305 310 315 320

Cys Ala Thr Asp Thr Gln Asn Val Lys Phe Val Phe Asp Ala Val Thr  
 325 330 335

Asp Ile Ile Ile Lys Glu Asn Leu Lys Asp Cys Gly Leu Phe  
 340 345 350

<210> 25

<211> 354

<212> PRT

<213> Homo sapiens

<400> 25

Met Gly Ser Gly Ala Ser Ala Glu Asp Lys Glu Leu Ala Lys Arg Ser  
 1 5 10 15

Lys Glu Leu Glu Lys Lys Leu Gln Glu Asp Ala Asp Lys Glu Ala Lys  
 20 25 30

Thr Val Lys Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr  
 35 40 45

Ile Val Lys Gln Met Lys Ile Ile His Gln Asp Gly Tyr Ser Pro Glu  
 50 55 60

Glu Cys Leu Glu Phe Lys Ala Ile Ile Tyr Gly Asn Val Leu Gln Ser  
 65 70 75 80

Ile Leu Ala Ile Ile Arg Ala Met Thr Thr Leu Gly Ile Asp Tyr Ala  
 85 90 95

Glu Pro Ser Cys Ala Asp Asp Gly Arg Gln Leu Asn Asn Leu Ala Asp  
 100 105 110

Ser Ile Glu Glu Gly Thr Met Pro Pro Glu Leu Val Glu Val Ile Arg  
 115 120 125

Arg Leu Trp Lys Asp Gly Gly Val Gln Ala Cys Phe Glu Arg Ala Ala  
 130 135 140

Glu Tyr Gln Leu Asn Asp Ser Ala Ser Tyr Tyr Leu Asn Gln Leu Glu  
 145 150 155 160

Arg Ile Thr Asp Pro Glu Tyr Leu Pro Ser Glu Gln Asp Val Leu Arg  
 165 170 175

Ser Arg Val Lys Thr Thr Gly Ile Ile Glu Thr Lys Phe Ser Val Lys  
 180 185 190

Asp Leu Asn Phe Arg Met Phe Asp Val Gly Gly Gln Arg Ser Glu Arg  
 195 200 205

Lys Lys Trp Ile His Cys Phe Glu Gly Val Thr Cys Ile Ile Phe Cys  
 210 215 220

Ala Ala Leu Ser Ala Tyr Asp Met Val Leu Val Glu Asp Asp Glu Val  
 225 230 235 240

Asn Arg Met His Glu Ser Leu His Leu Phe Asn Ser Ile Cys Asn His  
 245 250 255

Lys Phe Phe Ala Ala Thr Ser Ile Val Leu Phe Leu Asn Lys Lys Asp  
                   260                                  265                                  270  
 Leu Phe Glu Glu Lys Ile Lys Lys Val His Leu Ser Ile Cys Phe Pro  
                   275                                  280                                  285  
 Glu Tyr Asp Gly Asn Asn Ser Tyr Asp Asp Ala Gly Asn Tyr Ile Lys  
                   290                                  295                                  300  
 Ser Gln Phe Leu Asp Leu Asn Met Arg Lys Asp Val Lys Glu Ile Tyr  
                   305                                  310                                  315                                  320  
 Ser His Met Thr Cys Ala Thr Asp Thr Gln Asn Val Lys Phe Val Phe  
                                   325                                  330                                  335  
 Asp Ala Val Thr Asp Ile Ile Ile Lys Glu Asn Leu Lys Asp Cys Gly  
                   340                                  345                                  350  
 Leu Phe

<210> 26  
 <211> 466  
 <212> PRT  
 <213> Homo sapiens

<400> 26  
 Met Val Phe Leu Ser Gly Asn Ala Ser Asp Ser Ser Asn Cys Thr Gln  
           1                                  5                                  10                                  15  
 Pro Pro Ala Pro Val Asn Ile Ser Lys Ala Ile Leu Leu Gly Val Ile  
                   20                                  25                                  30  
 Leu Gly Gly Leu Ile Leu Phe Gly Val Leu Gly Asn Ile Leu Val Ile  
                   35                                  40                                  45  
 Leu Ser Val Ala Cys His Arg His Leu His Ser Val Thr His Tyr Tyr  
                   50                                  55                                  60  
 Ile Val Asn Leu Ala Val Ala Asp Leu Leu Leu Thr Ser Thr Val Leu  
                   65                                  70                                  75                                  80  
 Pro Phe Ser Ala Ile Phe Glu Val Leu Gly Tyr Trp Ala Phe Gly Arg  
                                   85                                  90                                  95  
 Val Phe Cys Asn Ile Trp Ala Ala Val Asp Val Leu Cys Cys Thr Ala  
                   100                                  105                                  110  
 Ser Ile Met Gly Leu Cys Ile Ile Ser Ile Asp Arg Tyr Ile Gly Val  
                   115                                  120                                  125  
 Ser His Pro Leu Arg Tyr Pro Thr Ile Val Thr Gln Arg Arg Gly Leu  
                   130                                  135                                  140  
 Met Ala Leu Leu Cys Val Trp Ala Leu Ser Leu Val Ile Ser Ile Gly  
                   145                                  150                                  155                                  160  
 Pro Leu Phe Gly Trp Arg Gln Pro Ala Pro Glu Asp Glu Thr Ile Cys  
                   165                                  170                                  175

Gln Ile Asn Glu Glu Pro Gly Tyr Val Leu Phe Ser Ala Leu Gly Ser  
 180 185 190  
 Phe Tyr Leu Pro Leu Ala Ile Ile Leu Val Met Tyr Cys Arg Val Tyr  
 195 200 205  
 Val Val Ala Lys Arg Glu Ser Arg Gly Leu Lys Ser Gly Leu Lys Thr  
 210 215 220  
 Asp Lys Ser Asp Ser Glu Gln Val Thr Leu Arg Ile His Arg Lys Asn  
 225 230 235 240  
 Ala Pro Ala Gly Gly Ser Gly Met Ala Ser Ala Lys Thr Lys Thr His  
 245 250 255  
 Phe Ser Val Arg Leu Leu Lys Phe Ser Arg Glu Lys Lys Ala Ala Lys  
 260 265 270  
 Thr Leu Gly Ile Val Val Gly Cys Phe Val Leu Cys Trp Leu Pro Phe  
 275 280 285  
 Phe Leu Val Met Pro Ile Gly Ser Phe Phe Pro Asp Phe Lys Pro Ser  
 290 295 300  
 Glu Thr Val Phe Lys Ile Val Phe Trp Leu Gly Tyr Leu Asn Ser Cys  
 305 310 315 320  
 Ile Asn Pro Ile Ile Tyr Pro Cys Ser Ser Gln Glu Phe Lys Lys Ala  
 325 330 335  
 Phe Gln Asn Val Leu Arg Ile Gln Cys Leu Cys Arg Lys Gln Ser Ser  
 340 345 350  
 Lys His Ala Leu Gly Tyr Thr Leu His Pro Pro Ser Gln Ala Val Glu  
 355 360 365  
 Gly Gln His Lys Asp Met Val Arg Ile Pro Val Gly Ser Arg Glu Ala  
 370 375 380  
 Phe Tyr Gly Ile Ser Arg Thr Asp Gly Val Cys Glu Trp Lys Phe Phe  
 385 390 395 400  
 Ser Ser Met Pro Arg Gly Ser Ala Arg Ile Thr Val Ser Lys Asp Gln  
 405 410 415  
 Ser Ser Cys Thr Thr Ala Arg Val Arg Ser Lys Ser Phe Leu Gln Val  
 420 425 430  
 Cys Cys Cys Val Glu Pro Ser Thr Pro Ser Leu Asp Lys Asn His Gln  
 435 440 445  
 Val Pro Thr Ile Lys Val His Thr Ile Ser Leu Ser Glu Asn Gly Glu  
 450 455 460  
 Glu Val  
 465

&lt;210&gt; 27

&lt;211&gt; 353

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 27

```

Met Ala Cys Cys Leu Ser Glu Glu Ala Lys Glu Ala Arg Arg Ile Asn
  1           5           10           15

Asp Glu Ile Glu Arg Gln Leu Arg Arg Asp Lys Arg Asp Ala Arg Arg
      20           25           30

Glu Leu Lys Leu Leu Leu Leu Gly Thr Gly Glu Ser Gly Lys Ser Thr
      35           40           45

Phe Ile Lys Gln Met Arg Ile Ile His Gly Ser Gly Tyr Ser Asp Glu
      50           55           60

Asp Lys Arg Gly Phe Thr Lys Leu Val Tyr Gln Asn Ile Phe Thr Ala
      65           70           75           80

Met Gln Ala Met Ile Arg Ala Met Asp Thr Leu Lys Ile Pro Tyr Lys
      85           90           95

Tyr Glu His Asn Lys Ala His Ala Gln Leu Val Arg Glu Val Asp Val
      100          105          110

Glu Lys Val Ser Ala Phe Glu Asn Pro Tyr Val Asp Ala Ile Lys Ser
      115          120          125

Leu Trp Asn Asp Pro Gly Ile Gln Glu Cys Tyr Asp Arg Arg Arg Glu
      130          135          140

Tyr Gln Leu Ser Asp Ser Thr Lys Tyr Tyr Leu Asn Asp Leu Asp Arg
      145          150          155          160

Val Ala Asp Pro Ala Tyr Leu Pro Thr Gln Gln Asp Val Leu Arg Val
      165          170          175

Arg Val Pro Thr Thr Gly Ile Ile Glu Tyr Pro Phe Asp Leu Gln Ser
      180          185          190

Val Ile Phe Arg Met Val Asp Val Gly Gly Gln Arg Ser Glu Arg Arg
      195          200          205

Lys Trp Ile His Cys Phe Glu Asn Val Thr Ser Ile Met Phe Leu Val
      210          215          220

Ala Leu Ser Glu Tyr Asp Gln Val Leu Val Glu Ser Asp Asn Glu Asn
      225          230          235          240

Arg Met Glu Glu Ser Lys Ala Leu Phe Arg Thr Ile Ile Thr Tyr Pro
      245          250          255

Trp Phe Gln Asn Ser Ser Val Ile Leu Phe Leu Asn Lys Lys Asp Leu
      260          265          270

Leu Glu Glu Lys Ile Met Tyr Ser His Leu Val Asp Tyr Phe Pro Glu
      275          280          285

Tyr Asp Gly Pro Gln Arg Asp Ala Gln Ala Ala Arg Glu Phe Ile Leu
      290          295          300

```



Lys Met Phe Val Asp Leu Asn Pro Asp Ser Asp Lys Ile Ile Tyr Ser  
 305 310 315 320  
 His Phe Thr Cys Ala Thr Asp Thr Glu Asn Ile Arg Phe Val Phe Ala  
 325 330 335  
 Ala Val Lys Asp Thr Ile Leu Gln Leu Asn Leu Lys Glu Tyr Asn Ala  
 340 345 350

Val

<210> 28  
 <211> 359  
 <212> PRT  
 <213> Homo sapiens

<400> 28  
 Met Thr Leu Glu Ser Met Met Ala Cys Cys Leu Ser Asp Glu Val Lys  
 1 5 10 15  
 Glu Ser Lys Arg Ile Asn Ala Glu Ile Glu Lys Gln Leu Arg Arg Asp  
 20 25 30  
 Lys Arg Asp Ala Arg Arg Glu Leu Lys Leu Leu Leu Gly Thr Gly  
 35 40 45  
 Glu Ser Gly Lys Ser Thr Phe Ile Lys Gln Met Arg Ile Ile His Gly  
 50 55 60  
 Ala Gly Tyr Ser Glu Glu Asp Lys Arg Gly Phe Thr Lys Leu Val Tyr  
 65 70 75 80  
 Gln Asn Ile Phe Thr Ala Met Gln Ala Met Ile Arg Ala Met Glu Thr  
 85 90 95  
 Leu Lys Ile Leu Tyr Lys Tyr Glu Gln Asn Lys Ala Asn Ala Leu Leu  
 100 105 110  
 Ile Arg Glu Val Asp Val Glu Lys Val Thr Thr Phe Glu His Gln Tyr  
 115 120 125  
 Val Ser Ala Ile Lys Thr Leu Trp Glu Asp Pro Gly Ile Gln Glu Cys  
 130 135 140  
 Tyr Asp Arg Arg Arg Glu Tyr Gln Leu Ser Asp Ser Ala Lys Tyr Tyr  
 145 150 155 160  
 Leu Thr Asp Val Asp Arg Ile Ala Thr Leu Gly Tyr Leu Pro Thr Gln  
 165 170 175  
 Gln Asp Val Leu Arg Val Arg Val Pro Thr Thr Gly Ile Ile Glu Tyr  
 180 185 190  
 Pro Phe Asp Leu Glu Asn Ile Ile Phe Arg Met Val Asp Val Gly Gly  
 195 200 205  
 Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu Asn Val Thr  
 210 215 220

Ser Ile Met Phe Leu Val Ala Leu Ser Glu Tyr Asp Gln Val Leu Val  
 225 230 235 240  
 Glu Ser Asp Asn Glu Asn Arg Met Glu Glu Ser Lys Ala Leu Phe Arg  
 245 250 255  
 Thr Ile Ile Thr Tyr Pro Trp Phe Gln Asn Ser Ser Val Ile Leu Phe  
 260 265 270  
 Leu Asn Lys Lys Asp Leu Leu Glu Asp Lys Ile Leu Tyr Ser His Leu  
 275 280 285  
 Val Asp Tyr Phe Pro Glu Phe Asp Gly Pro Gln Arg Asp Ala Gln Ala  
 290 295 300  
 Ala Arg Glu Phe Ile Leu Lys Met Phe Val Asp Leu Asn Pro Asp Ser  
 305 310 315 320  
 Asp Lys Ile Ile Tyr Ser His Phe Thr Cys Ala Thr Asp Thr Glu Asn  
 325 330 335  
 Ile Arg Phe Val Phe Ala Ala Val Lys Asp Thr Ile Leu Gln Leu Asn  
 340 345 350  
 Leu Lys Glu Tyr Asn Leu Val  
 355

<210> 29  
 <211> 381  
 <212> PRT  
 <213> Homo sapiens

<400> 29  
 Met Ser Gly Val Val Arg Thr Leu Ser Arg Cys Leu Leu Pro Ala Glu  
 1 5 10 15  
 Ala Gly Gly Ala Arg Glu Arg Arg Ala Gly Ser Gly Ala Arg Asp Ala  
 20 25 30  
 Glu Arg Glu Ala Arg Arg Arg Ser Arg Asp Ile Asp Ala Leu Leu Ala  
 35 40 45  
 Arg Glu Arg Arg Ala Val Arg Arg Leu Val Lys Ile Leu Leu Leu Gly  
 50 55 60  
 Ala Gly Glu Ser Gly Lys Ser Thr Phe Leu Lys Gln Met Arg Ile Ile  
 65 70 75 80  
 His Gly Arg Glu Phe Asp Gln Lys Ala Leu Leu Glu Phe Arg Asp Thr  
 85 90 95  
 Ile Phe Asp Asn Ile Leu Lys Gly Ser Arg Val Leu Val Asp Ala Arg  
 100 105 110  
 Asp Lys Leu Gly Ile Pro Trp Gln Tyr Ser Glu Asn Glu Lys His Gly  
 115 120 125  
 Met Phe Leu Met Ala Phe Glu Asn Lys Ala Gly Leu Pro Val Glu Pro  
 130 135 140

Ala Thr Phe Gln Leu Tyr Val Pro Ala Leu Ser Ala Leu Trp Arg Asp  
 145 150 155 160

Ser Gly Ile Arg Glu Ala Phe Ser Arg Arg Ser Glu Phe Gln Leu Gly  
 165 170 175

Glu Ser Val Lys Tyr Phe Leu Asp Asn Leu Asp Arg Ile Gly Gln Leu  
 180 185 190

Asn Tyr Phe Pro Ser Lys Gln Asp Ile Leu Leu Ala Arg Lys Ala Thr  
 195 200 205

Lys Gly Ile Val Glu His Asp Phe Val Ile Lys Lys Ile Pro Phe Lys  
 210 215 220

Met Val Asp Val Gly Gly Gln Arg Ser Gln Arg Gln Lys Trp Phe Gln  
 225 230 235 240

Cys Phe Asp Gly Ile Thr Ser Ile Leu Phe Met Val Ser Ser Ser Glu  
 245 250 255

Tyr Asp Gln Val Leu Met Glu Asp Arg Arg Thr Asn Arg Leu Val Glu  
 260 265 270

Ser Met Asn Ile Phe Glu Thr Ile Val Asn Asn Lys Leu Phe Phe Asn  
 275 280 285

Val Ser Ile Ile Leu Phe Leu Asn Lys Met Asp Leu Leu Val Glu Lys  
 290 295 300

Val Lys Thr Val Ser Ile Lys Lys His Phe Pro Asp Phe Arg Gly Asp  
 305 310 315 320

Pro His Gln Leu Glu Asp Val Gln Arg Tyr Leu Val Gln Cys Phe Asp  
 325 330 335

Arg Lys Arg Arg Asn Arg Ser Lys Pro Leu Phe His His Phe Thr Thr  
 340 345 350

Ala Ile Asp Thr Glu Asn Val Arg Phe Val Phe His Ala Val Lys Asp  
 355 360 365

Thr Ile Leu Gln Glu Asn Leu Lys Asp Ile Met Leu Gln  
 370 375 380

&lt;210&gt; 30

&lt;211&gt; 377

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 30

Met Ala Asp Phe Leu Pro Ser Arg Ser Val Leu Ser Val Cys Phe Pro  
 1 5 10 15

Gly Cys Leu Leu Thr Ser Gly Glu Ala Glu Gln Gln Arg Lys Ser Lys  
 20 25 30

Glu Ile Asp Lys Cys Leu Ser Arg Glu Lys Thr Tyr Val Lys Arg Leu  
 35 40 45

Val Lys Ile Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Phe  
 50 55 60  
 Leu Lys Gln Met Arg Ile Ile His Gly Gln Asp Phe Asp Gln Arg Ala  
 65 70 75 80  
 Arg Glu Glu Phe Arg Pro Thr Ile Tyr Ser Asn Val Ile Lys Gly Met  
 85 90 95  
 Arg Val Leu Val Asp Ala Arg Glu Lys Leu His Ile Pro Trp Gly Asp  
 100 105 110  
 Asn Ser Asn Gln Gln His Gly Asp Lys Met Met Ser Phe Asp Thr Arg  
 115 120 125  
 Ala Pro Met Ala Ala Gln Gly Met Val Glu Thr Arg Val Phe Leu Gln  
 130 135 140  
 Tyr Leu Pro Ala Ile Arg Ala Leu Trp Ala Asp Ser Gly Ile Gln Asn  
 145 150 155 160  
 Ala Tyr Asp Arg Arg Arg Glu Phe Gln Leu Gly Glu Ser Val Lys Tyr  
 165 170 175  
 Phe Leu Asp Asn Leu Asp Lys Leu Gly Glu Pro Asp Tyr Ile Pro Ser  
 180 185 190  
 Gln Gln Asp Ile Leu Leu Ala Arg Arg Pro Thr Lys Gly Ile His Glu  
 195 200 205  
 Tyr Asp Phe Glu Ile Lys Asn Val Pro Phe Lys Met Leu Asp Val Gly  
 210 215 220  
 Gly Gln Arg Ser Glu Arg Lys Arg Trp Phe Glu Cys Phe Asp Ser Val  
 225 230 235 240  
 Thr Ser Ile Leu Phe Leu Val Ser Ser Ser Glu Phe Asp Gln Val Leu  
 245 250 255  
 Met Glu Asp Arg Leu Thr Asn Arg Leu Thr Glu Ser Leu Asn Ile Phe  
 260 265 270  
 Glu Thr Ile Val Asn Asn Arg Val Phe Ser Asn Val Ser Ile Ile Leu  
 275 280 285  
 Phe Leu Asn Lys Thr Asp Leu Leu Glu Glu Lys Val Gln Ile Val Ser  
 290 295 300  
 Ile Lys Asp Tyr Phe Leu Glu Phe Glu Gly Asp Pro His Cys Leu Arg  
 305 310 315 320  
 Asp Val Gln Lys Phe Leu Val Glu Cys Phe Arg Asn Lys Arg Arg Asp  
 325 330 335  
 Gln Gln Gln Lys Pro Leu Tyr His His Phe Thr Thr Ala Ile Asn Thr  
 340 345 350  
 Glu Asn Ile Arg Leu Val Phe Arg Asp Val Lys Asp Thr Ile Leu His  
 355 360 365  
 Asp Asn Leu Lys Gln Leu Met Leu Gln

370

375

&lt;210&gt; 31

&lt;211&gt; 355

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

```

Met Ala Gly Cys Cys Cys Leu Ser Ala Glu Glu Lys Glu Ser Gln Arg
 1              5              10              15

Ile Ser Ala Glu Ile Glu Arg Gln Leu Arg Arg Asp Lys Lys Asp Ala
      20              25              30

Arg Arg Glu Leu Lys Leu Leu Leu Leu Gly Thr Gly Glu Ser Gly Lys
      35              40              45

Ser Thr Phe Ile Lys Gln Met Arg Ile Ile His Gly Ser Gly Tyr Ser
      50              55              60

Asp Glu Asp Arg Lys Gly Phe Thr Lys Leu Val Tyr Gln Asn Ile Phe
      65              70              75              80

Thr Ala Met Gln Ala Met Ile Arg Ala Met Asp Thr Leu Arg Ile Gln
      85              90              95

Tyr Val Cys Glu Gln Asn Lys Glu Asn Ala Gln Ile Ile Arg Glu Val
      100              105              110

Glu Val Asp Lys Val Ser Met Leu Ser Arg Glu Gln Val Glu Ala Ile
      115              120              125

Lys Gln Leu Trp Gln Asp Pro Gly Ile Gln Glu Cys Tyr Asp Arg Arg
      130              135              140

Arg Glu Tyr Gln Leu Ser Asp Ser Ala Lys Tyr Tyr Leu Thr Asp Ile
      145              150              155              160

Asp Arg Ile Ala Thr Pro Ser Phe Val Pro Thr Gln Gln Asp Val Leu
      165              170              175

Arg Val Arg Val Pro Thr Thr Gly Ile Ile Glu Tyr Pro Phe Asp Leu
      180              185              190

Glu Asn Ile Ile Phe Arg Met Val Asp Val Gly Gly Gln Arg Ser Glu
      195              200              205

Arg Arg Lys Trp Ile His Cys Phe Glu Ser Val Thr Ser Ile Ile Phe
      210              215              220

Leu Val Ala Leu Ser Glu Tyr Asp Gln Val Leu Ala Glu Cys Asp Asn
      225              230              235              240

Glu Asn Arg Met Glu Glu Ser Lys Ala Leu Phe Lys Thr Ile Ile Thr
      245              250              255

Tyr Pro Trp Phe Leu Asn Ser Ser Val Ile Leu Phe Leu Asn Lys Lys
      260              265              270

Asp Leu Leu Glu Glu Lys Ile Met Tyr Ser His Leu Ile Ser Tyr Phe

```

275                      280                      285  
 Pro Glu Tyr Thr Gly Pro Lys Gln Asp Val Arg Ala Ala Arg Asp Phe  
 290                      295                      300  
 Ile Leu Lys Leu Tyr Gln Asp Gln Asn Pro Asp Lys Glu Lys Val Ile  
 305                      310                      315                      320  
 Tyr Ser His Phe Thr Cys Ala Thr Asp Thr Asp Asn Ile Arg Phe Val  
 325                      330                      335  
 Phe Ala Ala Val Lys Asp Thr Ile Leu Gln Leu Asn Leu Arg Glu Phe  
 340                      345                      350  
 Asn Leu Val  
 355

<210> 32  
 <211> 374  
 <212> PRT  
 <213> Homo sapiens

<400> 32  
 Met Ala Arg Ser Leu Thr Trp Arg Cys Cys Pro Trp Cys Leu Thr Glu  
 1                      5                      10                      15  
 Asp Glu Lys Ala Ala Ala Arg Val Asp Gln Glu Ile Asn Arg Ile Leu  
 20                      25                      30  
 Leu Glu Gln Lys Lys Gln Asp Arg Gly Glu Leu Lys Leu Leu Leu Leu  
 35                      40                      45  
 Gly Pro Gly Glu Ser Gly Lys Ser Thr Phe Ile Lys Gln Met Arg Ile  
 50                      55                      60  
 Ile His Gly Ala Gly Tyr Ser Glu Glu Glu Arg Lys Gly Phe Arg Pro  
 65                      70                      75                      80  
 Leu Val Tyr Gln Asn Ile Phe Val Ser Met Arg Ala Met Ile Glu Ala  
 85                      90                      95  
 Met Glu Arg Leu Gln Ile Pro Phe Ser Arg Pro Glu Ser Lys His His  
 100                      105                      110  
 Ala Ser Leu Val Met Ser Gln Asp Pro Tyr Lys Val Thr Thr Phe Glu  
 115                      120                      125  
 Lys Arg Tyr Ala Ala Ala Met Gln Trp Leu Trp Arg Asp Ala Gly Ile  
 130                      135                      140  
 Arg Ala Cys Tyr Glu Arg Arg Arg Glu Phe His Leu Leu Asp Ser Ala  
 145                      150                      155                      160  
 Val Tyr Tyr Leu Ser His Leu Glu Arg Ile Thr Glu Glu Gly Tyr Val  
 165                      170                      175  
 Pro Thr Ala Gln Asp Val Leu Arg Ser Arg Met Pro Thr Thr Gly Ile  
 180                      185                      190  
 Asn Glu Tyr Cys Phe Ser Val Gln Lys Thr Asn Leu Arg Ile Val Asp

```

          195              200              205
Val Gly Gly Gln Lys Ser Glu Arg Lys Lys Trp Ile His Cys Phe Glu
  210              215              220
Asn Val Ile Ala Leu Ile Tyr Leu Ala Ser Leu Ser Glu Tyr Asp Gln
  225              230              235              240
Cys Leu Glu Glu Asn Asn Gln Glu Asn Arg Met Lys Glu Ser Leu Ala
          245              250              255
Leu Phe Gly Thr Ile Leu Glu Leu Pro Trp Phe Lys Ser Thr Ser Val
          260              265              270
Ile Leu Phe Leu Asn Lys Thr Asp Ile Leu Glu Glu Lys Ile Pro Thr
          275              280              285
Ser His Leu Ala Thr Tyr Phe Pro Ser Phe Gln Gly Pro Lys Gln Asp
          290              295              300
Ala Glu Ala Ala Lys Arg Phe Ile Leu Asp Met Tyr Thr Arg Met Tyr
  305              310              315              320
Thr Gly Cys Val Asp Gly Pro Glu Gly Ser Lys Lys Gly Ala Arg Ser
          325              330              335
Arg Arg Leu Phe Ser His Tyr Thr Cys Ala Thr Asp Thr Gln Asn Ile
          340              345              350
Arg Lys Val Phe Lys Asp Val Arg Asp Ser Val Leu Ala Arg Tyr Leu
          355              360              365
Asp Glu Ile Asn Leu Leu
          370

```

```

<210> 33
<211> 157
<212> PRT
<213> Homo sapiens

```

```

<400> 33
Met Phe Asp Val Gly Gly Gln Arg Ser Glu Arg Lys Lys Trp Ile His
  1              5              10              15
Cys Phe Glu Gly Val Thr Cys Ile Ile Phe Cys Ala Ala Leu Ser Ala
          20              25              30
Tyr Asp Met Val Leu Val Glu Asp Glu Glu Val Asn Arg Met His Glu
          35              40              45
Ser Leu His Leu Phe Asn Ser Ile Cys Asn His Lys Tyr Phe Ser Thr
          50              55              60
Thr Ser Ile Val Leu Phe Leu Asn Lys Lys Asp Ile Phe Gln Glu Lys
          65              70              75              80
Val Thr Lys Val His Leu Ser Ile Cys Phe Pro Glu Tyr Thr Gly Pro
          85              90              95
Asn Thr Phe Glu Asp Ala Gly Asn Tyr Ile Lys Asn Gln Phe Leu Asp

```

100                      105                      110  
 Leu Asn Leu Lys Lys Glu Asp Lys Glu Ile Tyr Ser His Met Thr Cys  
           115                      120                      125  
 Ala Thr Asp Thr Gln Asn Val Lys Phe Val Phe Asp Ala Val Thr Asp  
           130                      135                      140  
 Ile Ile Ile Lys Glu Asn Leu Lys Asp Cys Gly Leu Phe  
 145                      150                      155

<210> 34  
 <211> 75  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
           cAMP-Hammerhead RNA sequence

<400> 34  
 gggcgacccu gaugagccug uggaaacaga cguggcacau gacuacgucg aaacggugaa 60  
 agccguaggu ugccc 75

<210> 35  
 <211> 75  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
           cGMP-Hammerhead RNA sequence

<400> 35  
 gggcgacccu gaugagcccu gcgaugcaga aaggugcuga cgacacaucg aaacggugaa 60  
 agccguaggu ugccc 75

<210> 36  
 <211> 164  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Template for  
           randomized sequence

<220>  
 <221> misc\_feature  
 <222> (23)  
 <223> Wherein n is t or a or c or g.

<220>  
 <221> misc\_feature  
 <222> (36)  
 <223> Wherein n is a or t or a or c or g.

<220>  
 <221> misc\_feature



<222> (42)

<223> Wherein n is t or a or c or g.

<220>

<221> misc\_feature

<222> (55)..(56)

<223> Wherein n is t or a or c or g.

<220>

<221> misc\_feature

<222> (58)

<223> Wherein n is t or a or c or g.

<220>

<221> misc\_feature

<222> (63)

<223> Wherein n is t or a or c or g.

<220>

<221> misc\_feature

<222> (69)..(118)

<223> Wherein n is an a or u or c or g and some may be missing.

<400> 36

```
ggagttacct aacaagaaac agngaagtca accagnaaa cncacggaga cgtgnnanat 60
tacctggttn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnnnnnngg 120
acctactgag ctgacagtcc tgtttgatgc ataccgagta agtg 164
```

<210> 37

<211> 131

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-D02  
ERK specific nucleic acid sensor molecule

<400> 37

```
ggacuucggc gaaagccguu cgacccuacu cagacgcuag cgaauugguu ccucgaaagg 60
gaaagcguua uuaagaaacc aaaugaggg cuuagacagg agguuaggug cgucaaugcu 120
gcaaguua cu g 131
```

<210> 38

<211> 132

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-D05  
ERK specific nucleic acid sensor molecule

<400> 38

```
ggacuucggc gaaagccguu cgaccuucuc agacgcuagc gaauugguuc cucgaaaggg 60
aaagcguuau uaagaaacca aaaugaguaa gcuuagacag gagguuaggu gcucaaugcu 120
ugcaaguua cu g 132
```

<210> 39

<211> 133  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-E01

<400> 39  
ggacuucggc gaaagccguu cgacccgcag cagacgcua gaaauugguu ccucgaaagg 60  
gaaagcguaa uuaagaaacc aaaaugcggu ggcuaagaca ggagguuagg ugcgucaaug 120  
cugcaaguua cug 133

<210> 40  
<211> 95  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cCMP nucleic  
acid sensor molecule

<400> 40  
ggaccugau gagccuuuag ggccaagugu ggugaaagac acacgucgaa acggugaaag 60  
ccguaggucc ugcgugguu cuguucccuu cuucg 95

<210> 41  
<211> 94  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cAMP nucleic  
acid sensor molecule

<400> 41  
ggaccugau gagccugugg aaacagacgu ggcacaugac uacgucgaaa cggugaaagc 60  
cguagguccu ugcgugguuc uguucccuuc uucg 94

<210> 42  
<211> 94  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cGMP nucleic  
acid sensor molecule

<400> 42  
ggaccugau gagccuugcg augcaaaaag gugcugacga cacaucgaaa cggugaaagc 60  
cguagguccu ugcgugguuc uguucccuuc uucg 94

<210> 43  
<211> 24  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Capture oligo

<400> 43

acgcaccaag acaagggaag aagc

24

<210> 44

<211> 133

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-A02  
ppERK nucleic acid sensor molecule

<400> 44

ggacuucggc gaaagccguu cgaccggguug cagacgcua ggaauugguu ccucgaaagg 60  
gaaagcguua uuaagaaacc aaaauggaac cgcuaagaca ggagguuagg ugcgucaaug 120  
cugcaaguua cug 133

<210> 45

<211> 133

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW-45-33-B04  
ppERK nucleic acid sensor molecule

<400> 45

ggacuucggc gaaagccguu cgaccggguug cagacgcua ggaauugguu ccucgaaagg 60  
gaaagcguua uuaagaaacc aaaaugugau cgcuaagaca ggagguuagg ugcgucaaug 120  
cugcaaguua cug 133

<210> 46

<211> 96

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nucleic acid  
sensor precursor molecule

<400> 46

ggcgugaccu gaugagucac gcagacgcua gcgaauuggu uccucaaagg gggaaagcgu 60  
uaauaagaaa caaaaaugug uuacgaaac guuccc 96

<210> 47

<211> 108

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nucleic acid  
sensor precursor molecule

<400> 47

ggacuucggg ccagugcucg ugcacuaggc cguucgacca ugauaccagc aucgucuuga 60

ugcccuuggc agcaucuag acaggagguu aggugccucg ugaugucc

108

&lt;210&gt; 48

&lt;211&gt; 79

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Nucleic acid  
sensor precursor molecule

&lt;400&gt; 48

gggacgacccu gaugagccug gauaccaagc cgaaaggccc uggcaguua gacgaaacgg 60  
ugaaagccgu agguagccc 79

&lt;210&gt; 49

&lt;211&gt; 34

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Nucleic acid  
sensor molecule

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)

&lt;223&gt; First signaling moiety.

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (34)

&lt;223&gt; Second signaling moiety.

&lt;400&gt; 49

nauaccaagc cgaaaggccc uggcagugg uaun

34

&lt;210&gt; 50

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Thrombin  
nucleic acid sensor molecule

&lt;400&gt; 50

ccaaccggtt ggtgtggttg g

21

&lt;210&gt; 51

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Target  
molecule

<400> 51  
gcgactggac atcacgag

18

<210> 52  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKA peptide

<400> 52  
Leu Arg Ala Ser Leu Gly  
1 5

<210> 53  
<211> 16  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKC peptide

<400> 53  
Ala Ala Lys Ile Gln Ala Ser Phe Arg Gly His Met Ala Arg Lys Lys  
1 5 10 15

<210> 54  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: cdc2 peptide

<400> 54  
Pro Lys Thr Pro Lys Lys Ala Lys Lys Leu  
1 5 10

<210> 55  
<211> 15  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: DNA-PK peptide

<400> 55  
Glu Pro Pro Leu Ser Gln Glu Ala Phe Ala Asp Leu Trp Lys Lys  
1 5 10 15

<210> 56  
<211> 10  
<212> PRT  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: CK-1 peptide

&lt;400&gt; 56

Asp	Asp	Asp	Glu	Glu	Ser	Ile	Thr	Arg	Arg
1				5					10

&lt;210&gt; 57

&lt;211&gt; 10

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: CK-2 peptide

&lt;400&gt; 57

Arg	Arg	Arg	Glu	Glu	Glu	Thr	Glu	Glu	Glu
1				5					10

&lt;210&gt; 58

&lt;211&gt; 13

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Cam KII peptide

&lt;400&gt; 58

Lys	Lys	Ala	Leu	Arg	Arg	Gln	Glu	Thr	Val	Asp	Ala	Leu
1				5								10

&lt;210&gt; 59

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: p38 peptide

&lt;400&gt; 59

Ser	Thr	Lys	Val	Pro	Gln	Thr	Pro	Leu	His	Thr	Ser	Arg	Val
1				5									10

&lt;210&gt; 60

&lt;211&gt; 9

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: PKA peptide

&lt;400&gt; 60

Arg	Arg	Arg	Arg	Ser	Ile	Ile	Phe	Ile
1				5				

<210> 61  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKA alpha  
peptide

<400> 61  
Arg Arg Arg Arg Arg Lys Gly Ser Phe Arg Arg Lys Ala  
1 5 10

<210> 62  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKC beta I, II  
peptide

<400> 62  
Arg Lys Leu Lys Arg Lys Gly Ser Phe Arg Arg Lys Ala  
1 5 10

<210> 63  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKC gammaI  
peptide

<400> 63  
Arg Arg Arg Arg Arg Lys Gly Ser Phe Lys Lys Phe Ala  
1 5 10

<210> 64  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PKC delta  
peptide

<400> 64  
Ala Ala Arg Lys Arg Lys Gly Ser Phe Phe Tyr Gly Gly  
1 5 10

<210> 65  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PKC epsilon peptide

<220>

<221> VARIANT

<222> (3)

<223> Wherein Xaa is any amino acid.

<400> 65

Tyr Tyr Xaa Lys Arg Lys Met Ser Phe Phe Glu Phe Asp  
1 5 10

<210> 66

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PKC eta peptide

<220>

<221> VARIANT

<222> (12)

<223> Wherein Xaa is any amino acid.

<400> 66

Ala Arg Leu Arg Arg Arg Arg Ser Phe Arg Arg Xaa Arg  
1 5 10

<210> 67

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PKC zeta peptide

<400> 67

Arg Arg Phe Lys Arg Gln Gly Ser Phe Phe Tyr Phe Phe  
1 5 10

<210> 68

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PKC mu peptide

<400> 68

Ala Ala Leu Val Arg Gln Met Ser Val Ala Phe Phe Phe  
1 5 10



<210> 69  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Cam KII  
peptide

<400> 69  
Lys Arg Gln Gln Ser Phe Asp Leu Phe  
1 5

<210> 70  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: phosphorylase  
kinaseI peptide

<400> 70  
Phe Arg Met Met Ser Phe Phe Leu Phe  
1 5

<210> 71  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: SLK1 peptide

<400> 71  
Arg Arg Phe Gly Ser Leu Arg Arg Phe  
1 5

<210> 72  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: SPRK2 peptide

<400> 72  
Arg Arg Arg His Ser Arg Arg Arg  
1 5

<210> 73  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AKT/PKB

## peptide

&lt;220&gt;

&lt;221&gt; VARIANT

&lt;222&gt; (4)

&lt;223&gt; Wherein Xaa is any amino acid.

&lt;400&gt; 73

Arg	Lys	Arg	Xaa	Arg	Thr	Tyr	Ser	Phe	Gly
1				5					10

&lt;210&gt; 74

&lt;211&gt; 28

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Effector dependent release factor

&lt;400&gt; 74

guccuguuug augcauaccg aguaagug .

28

&lt;210&gt; 75

&lt;211&gt; 146

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Combined DNA/RNA Molecule: Substrate oligonucleotide

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: ERK specific ligase nucleic acid sensor molecule

&lt;400&gt; 75

uaagcucua	caggaccugg	uuuucgcgug	gauuggagga	cagauucugu	gaccgccgug	60
caucggcgaa	agccuuuagg	aggaauccga	ccagcuugcc	gaaagcggcu	ucaggucacg	120
tccagtagac	tagcattcca	gcgtac				146

&lt;210&gt; 76

&lt;211&gt; 145

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Ligase nucleic acid sensor molecule

&lt;400&gt; 76

catgcgacct	tacgatcaga	tgacctgcac	uggacuucgg	cgaagccgu	ucgaccacgc	60
uaaggaggau	uuccgaaagc	ggcuacgugc	cgccaguguc	uuagacagga	gguuaggugc	120
gcuuuggucc	aaggacaucu	cgaau				145

&lt;210&gt; 77

&lt;211&gt; 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Capture oligo

<400> 77

attcgagatg uccttgacc aaagc

25

<210> 78

<211> 121

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Hammerhead  
(endonuclease) nucleic acid sensor molecule

<400> 78

cucgagagcg auggcaaagc ugcaucagua cacggugcag acaaaggugu ccgaguaguc 60  
ccagcgaaag cguuggaugc cgauaaauagg uuuuuucccg uaacguucgc ugaccuguag 120  
g 121

<210> 79

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Capture oligo

<400> 79

cctacaggtc agcgaacgtt acgggt

26

<210> 80

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK2 activated  
nucleic acid sensor

<220>

<221> misc\_feature

<222> (17)..(20)

<223> Wherein n is a or u or c or g or nothing.

<220>

<221> misc\_feature

<222> (58)..(61)

<223> Wherein n is a or u or c or g or nothing.

<400> 80

gggcgacccu gaugagnnnn cuaaggagga uuuccgaaag cggcuacggg ccgccagnnn 60  
ncgaaacggg gaaagccgua gguugccc 88

<210> 81  
<211> 110  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 6

<400> 81  
gggcgacccu gaugagucac gcagacgcua gcgaauggu uccucgaaag gggaaagcgu 60  
uauuaagaaa caaaaaugug uuacgaaacg gugaaaggcc guagguagcc 110

<210> 82  
<211> 108  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 7

<400> 82  
gggcgacccu gaugagucac gagacgcua gcaauuggu ccucgaaagg ggaaagcgu 60  
auuaagaaac caaaauuguu acgaaacggu gaaaggccgu agguagcc 108

<210> 83  
<211> 108  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 8

<400> 83  
gggcgacccu gaugagcacg cagacgcua gcaauuggu ccucgaaagg ggaaagcgu 60  
auuaagaaac caaaugugu ucgaaacggu gaaaggccgu agguagcc 108

<210> 84  
<211> 108  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 9

<400> 84  
gggcgacccu gaugagucag cagacgcua gcaauuggu ccucgaaagg ggaaagcgu 60  
auuaagaaac caaauguuu acgaaacggu gaaaggccgu agguagcc 108

<210> 85

<211> 110  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 10

<400> 85  
gggcgacccu gaugagucac gcagacgcua gcgaauggu uccucgaaag gggaaagcgu 60  
uauuaagaaa ccaaaaugug uugcgaaacg gugaaaggcc guagguagcc 110

<210> 86  
<211> 104  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 11

<400> 86  
gggcgacccu gaugagugca gacgcuagcg aauggguucc ucgaaagggg aaagcguaau 60  
uaagaaacca aaugugcga aacggugaaa ggccguaggu agcc 104

<210> 87  
<211> 110  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 12

<400> 87  
gggcgacccu gaugagccuu gcagacgcua gcgaauggu uccucgaaag gggaaagcgu 60  
uauuaagaaa ccaaaaugua cgucgaaacg gugaaaggcc guagguagcc 110

<210> 88  
<211> 110  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 13

<400> 88  
gggcgacccu gaugagucug gcagacgcua gcgaauggu uccucgaaag gggaaagcgu 60  
uauuaagaaa ccaaaauguc auacgaaacg gugaaaggcc guagguagcc 110

<210> 89  
<211> 110

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ppBRK  
sensitive cis-hammerhead nucleic acid sensor  
molecule construct 14

<400> 89

gggcgacccu gaugagucug gcagacgcua gcgaauggu uccucgaaag gggaaagcgu 60  
uauuaagaaa ccaaaauguc uuacgaaacg gugaaaggcc guagguagcc 110

<210> 90

<211> 131

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-F08  
ERK specific nucleic acid sensor molecule

<400> 90

ggacuucggc gaaagccguu cgaccuacuc agacgcuagc gaauugguuc cucgaaaggg 60  
aaagcguaau uaagaaacca aaaugaguag cuuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

<210> 91

<211> 132

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-H08  
ERK specific nucleic acid sensor molecule

<400> 91

ggacuucggc gaaagccguu cgacccucgu cagacgcuag cgaaugguu ccucgaaagg 60  
gaaagcguaa uuaagaaacc aaaaugacaa gcuuagacag gagguuaggu gcgucaaugc 120  
ugcaaguac ug 132

<210> 92

<211> 132

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-H09  
ERK specific nucleic acid sensor molecule

<400> 92

ggacuucggc gaaagccguu cgaccucgcu cagacgcuag cgaaugguu ccucgaaagg 60  
gaaagcguaa uuaagaaacc aaaaugagcg acuuagacag gagguuaggu gcgucaaugc 120  
ugcaaguac ug 132

<210> 93

<211> 131

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-A10  
ERK specific nucleic acid sensor molecule

<400> 93

ggacuucggc gaaagccguu cgaccaacuc agacgcuagc gaauggguuc cucgaaaggg 60  
aaagcguaau uaagaaacca aaauggguug cuuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

<210> 94

<211> 131

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-F09  
ERK specific nucleic acid sensor molecule

<400> 94

ggacuucggc gaaagccguu cgaccguuuc agacgcuagc gaauggguuc cucgaaaggg 60  
aaagcguaau uaagaaacca aaaugaacag cuuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

<210> 95

<211> 132

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-G08  
ERK specific nucleic acid sensor molecule

<400> 95

ggacuucggc gaaagccguu cgacccuuc cagacgcuag cgaauugguu ccucgaaagg 60  
gaaagcguaa uuaagaaacc aaaaugagaa gcuuagacag gagguuaggu gcgucaaugc 120  
ugcaaguuaac ug 132

<210> 96

<211> 134

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-G02  
ppERK specific nucleic acid sensor molecule

<400> 96

ggacuucggc gaaagccguu cgaccauuug ucagacgcu gcgaaauuggu uccucgaaag 60  
ggaaagcgua auuaagaaac caaaauggca gaucuagac aggagguuag gugcguaau 120  
gcugcaagu acug 134

<210> 97

<211> 134

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-G03  
ppERK specific nucleic acid sensor molecule

<400> 97

ggacuucggc gaaagccguu cgaccuuuag ucagacgcua gcgaauggu uccucgaaag 60  
ggaaagcguu auuaagaaac caaaaugauu agacuuagac aggagguuag gugcguaau 120  
gcugcaaguu acug 134

<210> 98

<211> 131

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-H03  
ppERK specific nucleic acid sensor molecule

<400> 98

ggacuucggc gaaagccguu cgaccgcuc agacgcuagc gaaugguuc cucgaaaggg 60  
aaagcguuau uaagaaacca aaugaucgg cuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

<210> 99

<211> 130

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-H1  
ppERK specific nucleic acid sensor molecule

<400> 99

ggacuucggc gaaagccguu cgaccguugc agacgcuagc gaaugguuc cucgaaaggg 60  
aaagcguuau uaagaaacca aaugcaauc uuagacagga gguuaggugc gucaaugcug 120  
caaguuacug 130

<210> 100

<211> 130

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: CW45-33-B05  
ppERK specific nucleic acid sensor molecule

<400> 100

ggacuucggc gaaagccguu cgaccuuagc agacgcuagc gaaugguuc cucgaaaggg 60  
aaagcguuau uaagaaacca aaugcuaac uuagacagga gguuaggugc gucaaugcug 120  
caaguuacug 130

<210> 101

<211> 98

<212> RNA

<213> Artificial Sequence



<220>

<223> Description of Artificial Sequence: cGMP specific  
hammerhead nucleic acid sensor molecule

<400> 101

ggauguccag ucgcugcaa ugccuuuuu gaccugaug agccugugga aacagacgug 60  
gcacaugacu acgucgaaac ggugaaagcc guaggucu 98

<210> 102

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Capture oligo

<400> 102

tgggcattgc aagcgactgg acatcc 26

<210> 103

<211> 70

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cAMP specific  
nucleic acid sensor molecule

<400> 103

gggaccuga ugagccugug gaaacagacg uggcacauga cuacgucgaa acggugaaag 60  
ccguaggucc 70

<210> 104

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Combined DNA/RNA Molecule: nucleic  
acid sensor molecule ligase

<220>

<223> Description of Artificial Sequence: Capture oligo

<400> 104

cgaaaccagc ttccgtagag ctta 24

<210> 105

<211> 81

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: C.lys.L1.A  
lysozyme sensitive ligase nucleic acid sensor

<400> 105

ggaccucggc gaaagcuaac gucucauggc uaaaauugcca uguugcuaca aaugauauga 60  
cuagagaggu uagcgagagu g 81

<210> 106

<211> 87

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: C.lys.L1.B  
ligase nucleic acid sensor molecule

<400> 106

ggaccucggc gaaagcuaac gucucauggc uaaaauugcca uguugcuaca aaugauauga 60  
cuagagaggu uaggugcgag agcacug 87

<210> 107

<211> 98

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: C.lys.L1.C  
ligase nucleic acid sensor molecule

<400> 107

ggaccucggc gaaagcuaac gucucauggc uaaaauugcca uguugcuaca aaugauauga 60  
cuagagaggu uaaggugcga gccgagaggc ugcacug 98

<210> 108

<211> 119

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (33)..(36)

<223> Wherein n is a or u or t or g or nothing.

<220>

<221> misc\_feature

<222> (70)..(73)

<223> Wherein n is a or u or c or g or nothing.

<400> 108

ugcacugggac uucggcgaaa gccguucgac cagcuaagga ggauuuccga aagcggcuac 60  
gugccgccag cucuuagaca ggagguuagg ugccguccga cugaucucgg aguuaaacg 119

<210> 109

<211> 121

<212> RNA

<213> Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule construct 17

&lt;400&gt; 109

```

ugcacuggac uucggcgaaa gccguucgac cagcuaagg aggauuuccg aaagcggcua 60
cgugccgccca gugucuuga caggagguua ggugccgucc gacugaucuc ggaguuaaac 120
g                                                                 121

```

&lt;210&gt; 110

&lt;211&gt; 119

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule construct 18

&lt;400&gt; 110

```

ugcacuggac uucggcgaaa gccguucgac cagcuaagga ggauuuccga aagcggcuac 60
gugccgccag uucuuagaca ggagguuagg ugccguccga cugaucucgg aguuaaacg 119

```

&lt;210&gt; 111

&lt;211&gt; 119

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule construct 19

&lt;400&gt; 111

```

ugcacuggac uucggcgaaa gccguucgac cagcuaagga ggauuuccga aagcggcuac 60
gugccgccag cucuuagaca ggagguuagg ugccguccga cugaucucgg aguuaaacg 119

```

&lt;210&gt; 112

&lt;211&gt; 121

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule construct 20

&lt;400&gt; 112

```

ugcacuggac uucggcgaaa gccguucgac cagcguaagg aggauuuccg aaagcggcua 60
cgugccgccca uugucuuga caggagguua ggugccgucc gacugaucuc ggaguuaaac 120
g                                                                 121

```

&lt;210&gt; 113

&lt;211&gt; 121

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK dependent  
ligase nucleic acid sensor molecule construct 21

&lt;400&gt; 113

ugcacuggac uucggcgaaa gccguucgac caccuaagg aggauuuccg aaagcggcua 60  
 cgugccgcca cguucuuaga caggagguaa ggugccgucc gacugaucuc ggaguuaaac 120  
 g 121

&lt;210&gt; 114

&lt;211&gt; 119

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK depedent  
 ligase nucleic acid sensor moleucle construct 22

&lt;400&gt; 114

ugcacuggac uucggcgaaa gccguucgac ccgcuaagga ggauuuccga aagcggcuac 60  
 gugccgccag ugcuuagaca ggagguuagg ugccguccga cugaucucgg aguuaaacg 119

&lt;210&gt; 115

&lt;211&gt; 117

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK depedent  
 ligase nucleic acid sensor moleucle construct 23

&lt;400&gt; 115

ugcacuggac uucggcgaaa gccguucgac cgcuaaggag gauuuccgaa agcggcuacg 60  
 ugccgccagu cuuagacagg agguuaggug ccguccgacu gaucucggag uuaaacg 117

&lt;210&gt; 116

&lt;211&gt; 117

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK depedent  
 ligase nucleic acid sensor moleucle construct 24

&lt;400&gt; 116

ugcacuggac uucggcgaaa gccguucgac cgcuaaggag gauuuccgaa agcggcuacg 60  
 ugccgccagc cuuagacagg agguuaggug ccguccgacu gaucucggag uuaaacg 117

&lt;210&gt; 117

&lt;211&gt; 119

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK depedent  
 ligase nucleic acid sensor moleucle construct 26

&lt;400&gt; 117

ugcacuggac uucggcgaaa gccguucgac cccuaagga ggauuuccga aagcggcuac 60  
 gugccgccac gucuuagaca ggagguuagg ugccguccga cugaucucgg aguuaaacg 119

<210> 118  
<211> 118  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: 3-piece ERK  
dependent ligase

<220>  
<221> misc\_feature  
<222> (32)..(35)  
<223> Wherein n is a or u or c or g or nothing.

<220>  
<221> misc\_feature  
<222> (68)..(71)  
<223> Wherein n is a or u or c or g or nothing.

<400> 118  
ugcacuggac uucggcgaaa gccguucgac cnnnaagga ggauuuccga aagcggcuac 60  
ggccgcennn ncuuagacag gagguuaggu gccguccgac ugaucucgga guuaaacg 118

<210> 119  
<211> 95  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: 1-piece ERK  
dependent kinase

<400> 119  
ggacuucggc gaaagccguu cgaccagcua aggaggauuu ccgaaagcgg cuacggccgc 60  
cagcucuuaag acaggagguu aggugcgaga gcacu 95

<210> 120  
<211> 96  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
template

<220>  
<221> misc\_feature  
<222> (1)..(13)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (83)..(96)  
<223> Wherein n is an a or u or c or g or nothing

<400> 120

nnnnnnnnnn nnngucacgc agacgcuagc gaauugguuc cucgaaaggg gaaagcguaa 60  
 uuaagaaacc aaaauguguu acnnnnnnnn nnnnnn 96

<210> 121  
 <211> 139  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ppERK  
 dependent ligase nucleic acid sensor molecule  
 template

<400> 121  
 ggacttcggc gaaagccggt cgaccacgct cagacgctag cgaattgggt cctcgaaagg 60  
 ggaaagcggt attaagaaac caaatgagt gtcttagaca ggaggttagg tgcggctttg 120  
 gtccaaggac atctcgat 139

<210> 122  
 <211> 137  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ppERK  
 dependent ligase nucleic acid sensor molecule  
 construct TK.16.118.L

<400> 122  
 ggacttcggc gaaagccggt cgaccagctc agacgctagc gaattgggtc ctcgaaaggg 60  
 gaaagcggtt ttaagaaacc aaaatgagt cttagacagg aggttaggtg cggctttggt 120  
 ccaaggacat ctcgat 137

<210> 123  
 <211> 139  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ppERK  
 dependent ligase nucleic acid sensor molecule  
 construct TK.16.118.M

<400> 123  
 ggacttcggc gaaagccggt cgaccacggt cagacgctag cgaattgggt cctcgaaagg 60  
 ggaaagcggtt attaagaaac caaatgatt gtcttagaca ggaggttagg tgcggctttg 120  
 gtccaaggac atctcgat 139

<210> 124  
 <211> 139  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ppERK  
 dependent ligase nucleic acid sensor molecule  
 construct TK.16.118.N

<400> 124  
ggacttcggc gaaagccggt cgaccacctt cagacgctag cgaattgggt ctcgaaaagg 60  
ggaaagcggtt attaagaaac caaaatgacg ttcttagaca ggaggtagg tgcggctttg 120  
gtccaaggac atctcgaat 139

<210> 125  
<211> 137  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.O

<400> 125  
ggacttcggc gaaagccggt cgaccagctc agacgctagc gaattgggtc ctcgaaaagg 60  
gaaagcggtt ttaagaaacc aaaatgagct ctagacagg aggttaggtg cggctttggt 120  
ccaaggacat ctcgaat 137

<210> 126  
<211> 137  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.P

<400> 126  
ggacttcggc gaaagccggt cgaccgctc agacgctagc gaattgggtc ctcgaaaagg 60  
gaaagcggtt ttaagaaacc aaaatgagt ctagacagg aggttaggtg cggctttggt 120  
ccaaggacat ctcgaat 137

<210> 127  
<211> 135  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.Q

<400> 127  
ggacttcggc gaaagccggt cgaccgctca gacgctagc aattgggtcc tcgaaaaggg 60  
aaagcggttat taagaaacca aaatgagtct tagacaggag gttagggtgc gctttggtcc 120  
aaggacatct cgaat 135

<210> 128  
<211> 137  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.R

<400> 128  
ggacttcggc gaaagccggtt cgacccggtc agacgctagc gaattgggtc ctcgaaaggg 60  
gaaagcggtta ttaagaaacc aaaatgattg cttagacagg aggttaggtg cggctttggt 120  
ccaaggacat ctggaat 137

<210> 129  
<211> 137  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.S

<400> 129  
ggacttcggc gaaagccggtt cgaccccttc agacgctagc gaattgggtc ctcgaaaggg 60  
gaaagcggtta ttaagaaacc aaaatgacgt cttagacagg aggttaggtg cggctttggt 120  
ccaaggacat ctggaat 137

<210> 130  
<211> 135  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK  
dependent ligase nucleic acid sensor molecule  
construct TK.16.118.T

<400> 130  
ggacttcggc gaaagccggtt cgaccgctca gacgctagcg aattgggttc tcgaaagggg 60  
aaagcggttat taagaaacca aaatgagcct tagacaggag gttagggtcg gctttggtcc 120  
aaggacatct cgaat 135

<210> 131  
<211> 132  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: CW45-33-C08  
BRK specific nucleic acid sensor molecule

<400> 131  
ggacuucggc gaaagccguu cgacccucuc cagacgcuag cgaaauuggu ccucgaaagg 60  
gaaagcguaa uuaagaaacc aaaaugagg gcuuagacag gagguuaggu gcucaaugc 120  
ugcaaguac ug 132

<210> 132  
<211> 131  
<212> RNA  
<213> Artificial Sequence



&lt;220&gt;

<223> Description of Artificial Sequence: CW45-33-C09  
ERK specific nucleic acid sensor molecule

&lt;400&gt; 132

ggacuucggc gaaagccguu cgaccugauc agacgcuagc gaaauugguuc cucgaaaggg 60  
aaagcguuau uaagaaacca aaaugaacag cuuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

&lt;210&gt; 133

&lt;211&gt; 131

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: CW45-33-D09  
ERK specific nucleic acid sensor molecule

&lt;400&gt; 133

ggacuucggc gaaagccguu cgaccuuuuc agacgcuagc gaaauugguuc cucgaaaggg 60  
aaagcguuau uaagaaacca aaaugaaaag cuuagacagg agguuaggug cgucaaugcu 120  
gcaaguacu g 131

&lt;210&gt; 134

&lt;211&gt; 108

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ppERK  
Sensitive nucleic acid sensor

&lt;400&gt; 134

ggcaacctac ggcctttcac cgtttcgtaa cacattttgg tttcttaata acgctttccc 60  
ctttcgagga accaattcgc tagcgtctgc gtgactcatc agggtcgc 108

&lt;210&gt; 135

&lt;211&gt; 90

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: cGMP specific  
nucleic acid sensor molecule

&lt;400&gt; 135

ggauauagc cguagguugc gaaagcgacc cugaugagcc cugcgaugca gaaaggugcu 60  
gacgacacau cgaaacgguu gcgagagcuc 90

&lt;210&gt; 136

&lt;211&gt; 88

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: cCMP specific

## nucleic acid sensor molecule

&lt;400&gt; 136

ggauaaauagc cguagguugc gaaagcgacc cugaugaccu guggaaacag acguggcaca 60  
 ugacuacgug aaacgguagc gagagcuc 88

&lt;210&gt; 137

&lt;211&gt; 88

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: cAMP specific  
 nucleic acid sensor molecule

&lt;400&gt; 137

ggauaaauagc cguagguugc gaaagcgacc cugaugaccu ugcgauagcaa aaaggugcug 60  
 acgacacaug aaacgguagc gagagcuc 88

&lt;210&gt; 138

&lt;211&gt; 71

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: cCMP specific  
 nucleic acid sensor molecule

&lt;400&gt; 138

gggacccuga ugagccuuua gggccaagug uggugaaaga cacacgucga aacggugaaa 60  
 gccguagguc c 71

&lt;210&gt; 139

&lt;211&gt; 70

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: cGMP specific  
 nucleic acid sensor molecule

&lt;400&gt; 139

gggacccuga ugagccuugc gaugcaaaaa ggugcugacg acacaucgaa acggugaaaag 60  
 ccguaggucc 70

&lt;210&gt; 140

&lt;211&gt; 89

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

&lt;400&gt; 140

gggcgaccu gaugagucgg gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
 auacgaaacg gugaaagccg uagguugcc 89

<210> 141  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: BRK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 141  
 gggcgacccu gaugagggag gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 gcgcgaaacg gugaaagccg uagguugcc 89

<210> 142  
 <211> 87  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (70)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (75)..(77)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (79)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (81)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (86)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 142  
 gggcgacccu gaugaggguu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 gccaaacagn gaaannngna nguggnc 87

<210> 143  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 143

gggcgacccu gaugaguacg gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
gaacgaaacg gugaaagccg uagguugcc 89

<210> 144

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (17)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (63)

<223> Wherein n is an a or u or c or g or nothing

<400> 144

gggcgacccu gaugagncau gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
acncgaaacg gugaaagccg uagguugcc 89

<210> 145

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 145

gggcgacccu gaugaguugc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
uaacgaaacg gugaaagccg uagguugcc 89

<210> 146

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 146

gggcgacccu gaugaguguu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
cuacgaaacg gugaaagccg uagguugcc 89

<210> 147

<211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (71)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (83)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (85)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 147  
 gggcgacccu gaugaguaau gcuaaggagg auuuccgaaa gcggcucggg ccgccagua 60  
 ggccaaacgg ngaaagcccg uangnugcc 89

<210> 148  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 148  
 gggcgacccu gaugagcucu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
 uggccgaaac ggugaaagcc guagguugcc 90

<210> 149  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 149  
 gggcgacccu gaugaggcau gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 cggcgaaacg gugaaagccg uagguugcc 89

<210> 150  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 150  
 gggcgacccu gaugagcccc gcuaaggagg auuuccgaaa gcggcuacug guccgccagu 60  
 ugagcgaaac ggugaaagcc guagguugcc 90

<210> 151  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (24)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (56)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (61)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (61)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (63)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (71)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (73)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 151  
 gggcgacccu gaugaguaag gcunaggagg auuuccgaaa gcggnuacug guccgncagu 60  
 nanacgaaac ngngaaagcc guagguugcc 90

<210> 152  
<211> 91  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 152  
gggcgacccu gaugagccau gcuaaggagg auuuccgaaa gcggcuacug guccgccagu 60  
acuacagaaa cgugaaagc cguagguugc c 91

<210> 153  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 153  
gggcgacccu gaugagugac gcuaaggagg auuuccgaaa gcggcuacug guccgccagu 60  
ugcgcgaaac ggugaaagcc guagguugcc 90

<210> 154  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 154  
gggcgacccu gaugagucug gcuaaggagg auuuccgaaa gcggcuacug guccgccagu 60  
uucgcgaaac ggugaaagcc guagguugcc 90

<210> 155  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (17)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (19)

<223> Wherein n is an a or u or c or g or nothing

<400> 155

ggcgacccu gaugagnanc gcuaaggagg auuuccgaaa gcggcuacug guccgccagu 60  
gauacgaaac ggugaaagcc guagguugcc 90

<210> 156

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (4)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (13)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (16)..(17)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (24)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (54)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (60)

<223> Wherein n is an a or u or c or g or nothing

<400> 156

gggngacccu ganganngac gcunaagagg auuuccgaag cgguacuag uccncauugn 60  
accgaaacgg ccuaaagccg gagguugcc 89

<210> 157

<211> 92

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>



<221> misc\_feature

<222> (46)

<223> Wherein n is an a or u or c or g or nothing

<400> 157

gggcgacccu gaugagauac gcuaaggagg auuuccgaaa gccggnuacg guccgacagu 60  
cuagccgaaa cggugaaaagc cguagguugu cc 92

<210> 158

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 158

gggcgacccu gaugaguggc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
caacgaaacg gugaaagccg uagguugcc 89

<210> 159

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 159

gggcgacccu gaugagcugu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
cgacgaaacg gugaaagccg uagguugcc 89

<210> 160

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 160

gggcgacccu gaugaggcug gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
agacgaaacg gugaaagccg uagguugcc 89

<210> 161

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature  
 <222> (59)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 161  
 gggcgacccu gaugagaacu gcuaaggagg auuuccgaaa gcggcuacgg uccgucaguu 60  
 acucgaaacg gugaaagccu guagguugcc 90

<210> 162  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: BRK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 162  
 gggcgacccu gaugagngcu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 cugcgaaacg gugaaagccg uagguugcc 89

<210> 163  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (19)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 163  
 gggcgacccu gaugagnang gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
 augcgaaacg gugaaagccg uagguugcc 89

<210> 164  
 <211> 88  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (19)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (53)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (57)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (60)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (68)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (70)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (75)..(76)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (80)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (82)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (87)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 164  
 gggcgacccu gaugagnunu gcuaaggagg auuuccgaaa gcgcuuncgg ccncacnacn 60  
 ccgaaacngn gaaannccgn anguggnc 88

<210> 165

<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 165  
gggcgacccu gaugaggagu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
auacgaaacg gugaaagccg uagguugcc 89

<210> 166  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (61)  
<223> Wherein n is an a or u or c or g or nothing

<400> 166  
gggcgacccu gaugagaccu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguc 60  
ncacgaaacg gugaaagccg uagguugcc 89

<210> 167  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 167  
gggcgacccu gaugaggcgc gcuaaggagg aaauuccgaaa gcggcuacgg uccgccagug 60  
uaacgaaacg gugaaagccg uagguugcc 89

<210> 168  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (73)  
<223> Wherein n is an a or u or c or g or nothing

<400> 168

gggcgacccu gaugaguaac gcuaaggagg auuuccgaaa gucggguacg guccgccagu 60  
 uuaucgaaac gngaaagcc guagguugcc 90

<210> 169

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (17)

<223> Wherein n is an a or u or c or g or nothing

<400> 169

gggcgacccu gaugagnuac gcuaaggagg auuuccgaaa gcgguacgg uccgccagug 60  
 ucacgaaacg gugaaagccg uagguugcc 89

<210> 170

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (82)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (84)

<223> Wherein n is an a or u or c or g or nothing

<400> 170

gggcgacccu gaugaguugc gcuaaggagg auuuccgaaa gcgguacgg uccgccaguu 60  
 acgcgaaacg gugaaagcgu angnugcc 88

<210> 171

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (18)

<223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (20)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (60)..(61)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 171  
 gggcgacccu gaugagunun gcuaaggagg auuuccgaaa gcggcuacgg uccgccagun 60  
 nuacgaaacg gugaaagccg uagguugcc 89

<210> 172  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (21)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 172  
 gggcgacccu gaugagacug ncuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 ggacgaaacg gugaaagccg uagguugcc 89

<210> 173  
 <211> 88  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 173  
 gggcgacccu gaugaggugc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagac 60  
 gucgaaacg ugaaagccgu agguugcc 88

<210> 174  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 174  
 gggcgacccu gaugaguauu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 uagcgaaacg gugaaagccg uagguugcc 89

<210> 175  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 175  
gggcgacccu gaugaguggc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
aaccgaaacg gugaaagccg uagguugcc 89

<210> 176  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 176  
gggcgacccu gaugagcuca gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
uggcgaaacg gugaaagccg uagguugcc 89

<210> 177  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (60)  
<223> Wherein n is an a or u or c or g or nothing

<400> 177  
gggcgacccu gaugagccun gcuaaggagg auuuccgaaa gcggcuacgg uccgccagun 60  
acugcgaaac ggugaaagcc guagguugcc 90

<210> 178  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 178  
 gggcgacccu gaugagucan gcuaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 uugcgaaacg gugaaagccg uagguugcc 89

<210> 179  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 179  
 gggcgacccu gaugagaacc gcuaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 augcgaaacg gugaaagccg uagguugcc 89

<210> 180  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (19)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 180  
 gggcgacccu gaugagggng gcuaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 acacgaaacg gugaaagccg uagguugcc 89

<210> 181  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (54)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (83)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 181  
 gggcgacccu gaugagugac gcuaggagg auuuccgaaa gcggcuacgg uccnccaguu 60  
 uaacgaaacg gugaaagccg uangguugcc 89



<210> 182  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (19)..(20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (40)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (60)..(61)  
<223> Wherein n is an a or u or c or g or nothing

<400> 182  
gggcgacccu gagaguacnn gcuaaggagg auuuccgaan gcggcuacgg uccgccagun 60  
naacgaaacg gugaaagccg uagguugcc 89

<210> 183  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 183  
gggcgacccu gaugagcuca gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguc 60  
aguacgaaac ggugaaagcc guagguugcc 90

<210> 184  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<400> 184

gggcgacccu gagaggauan gcuaaggagg auuuccgaan gcggcuacgg uccgccagua 60  
caacgaaacg gugaaagccg uagguugcc 89

<210> 185

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 185

gggcgacccu gaugaguagu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
ugacgaaacg gugaaagccg uagguugcc 89

<210> 186

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 186

gggcgacccu gaugagucgu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
gaacgaaacg gugaaagccg uagguugcc 89

<210> 187

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (12)..(13)

<223> Wherein n is an a or u or c or g or nothing

<400> 187

gggcgacccu gnngagaacu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
augcgaaacg gugaaagccg uanguugcc 89

<210> 188

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (13)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (17)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (24)  
<223> Wherein n is an a or u or c or g or nothing

<400> 188  
gggcgacccu gangagnucn gcunaggagg auuuccgaaa gcggcuacgg agccgucagu 60  
auugcgaaac ggcgaaagcc guagguugnc 90

<210> 189  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 189  
gggcgacccu gaugaguaac gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
uugcgaaacg gugaaagccg uagguugcc 89

<210> 190  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 190  
gggcgacccu gaugagcuua gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
aaacgaaacg gugaaagccg uagguugcc 89

<210> 191  
<211> 89  
<212> RNA  
<213> Artificial Sequence

```

<220>
<223> Description of Artificial Sequence: ERK sensitive
        hammerhead nucleic acid sensor molecule

<220>
<221> misc_feature
<222> (17)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (17)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (51)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (72)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (83)
<223> Wherein n is an a or u or c or g or nothing

<400> 191
gggcgacccu gaugagncca gcuaaggagg auuuccgaaa gcggcuacgg nccgccaguu 60
accgaaaacg gngaaagccg uanguugcc                                     89

<210> 192
<211> 89
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: ERK sensitive
        hammerhead nucleic acid sensor molecule

<220>
<221> misc_feature
<222> (17)..(18)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (51)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (72)
<223> Wherein n is an a or u or c or g or nothing

<400> 192
gggcgacccu gaugagncca gcuaaggagg auuuccgaaa gcggcuacgg nccgccaguu 60
accgaaaacg gngaaagccg uagguugcc                                     89

```

<210> 193

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 193

gggcgacccu gaugagcuuc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
agucgaaacg gugaaagccg uagguugcc 89

<210> 194

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (18)..(20)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (61)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (63)

<223> Wherein n is an a or u or c or g or nothing

<400> 194

gggcgacccu gaugagcnnn gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
nuncgaaacg gugaaagccg uagguugccu 90

<210> 195

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 195

gggcgacccu gaugagaccu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
aaucgaaacg gugaaagccg uagguugcc 89

<210> 196

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 196

gggcgacccu gaugagcacu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguc 60  
gcgcgaaacg gugaaagccg uagguugcc 89

<210> 197

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (60)

<223> Wherein n is an a or u or c or g or nothing

<400> 197

gggcgacccu gaugagaucu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagun 60  
acgcgaaacg gugaaagccg uagguugcc 89

<210> 198

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (13)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<400> 198

gggcgacccu gangaguuuu gcuaaggagg auuuccgaan gcggcuacgg uccgccagua 60  
auccgaaacg gugaaagccg uagguugcc 89

<210> 199

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (12)..(13)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<400> 199

gggcgacccu gnnagauua gcuaaggagg auuuccgaan gcggcuacgg uccgccaguu 60  
uagcgaaacg gugaaagccg uagguugcc 89

<210> 200

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (12)..(13)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<400> 200

gggcgacccu gnnagagcgu gcuaaggagg auuuccgaan gcggcuacgg uccgccaguu 60  
agacgaaacg gugaaagccg uagguugcc 89

<210> 201

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<400> 201

gggcgacccu gaugaguacu gcuaaggagg auuuccgaan gcggcuacgg uccgccagua 60  
cuacgaaacg gugaaagccg uagguugcc 89

<210> 202  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 202  
gggcgacccu gaugagcaga gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
uuacgaaacg gugaaagccg uagguugcc 89

<210> 203  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (39)  
<223> Wherein n is an a or u or c or g or nothing

<400> 203  
gggcgacccu gaugagcaua gcuaaggagg auuuccgana gcggcuacgg uccgccaguu 60  
gcacgaaacg gugaaagccg uagguugcc 89

<210> 204  
<211> 81  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (3)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (34)  
<223> Wherein n is an a or u or c or g or nothing

<400> 204  
ugnagagcgu ugcuaaggag gauuuccgaa agcngcuacg uccgccagua acacgaaacg 60  
gugaaagccg uagguugccu u 81

<210> 205  
<211> 89  
<212> RNA  
<213> Artificial Sequence



<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 205

gggcgacccu gaugagcaac gcuaaggaga auuuccgaaa gcggcuacgg uccgccaguu 60  
accgaaaacg gugaaagccg uagguugcc 89

<210> 206

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 206

gggcgacccu gaugagggag gcuaaggagg auuuccgaaa ucggcuacgg uccgccaguc 60  
auacgaaaacg gugaaagccg uagguugcc 89

<210> 207

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (9)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (45)

<223> Wherein n is an a or u or c or g or nothing

<400> 207

gggcgaccnu gaugagcuuc gcuaaggagg auuuccgaaa gcggnuacgg uccgccagug 60  
auucgaaaacg gugaaagccg uagguugcc 89

<210> 208

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (45)

<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (73)  
<223> Wherein n is an a or u or c or g or nothing

<400> 208  
gggcgacccu gaugagaccg gcuaaggagg auuuccgaaa gcggnuacgg uccgccaguc 60  
uaucgaaacg gunaaagccg uagguugcc 89

<210> 209  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (9)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (45)  
<223> Wherein n is an a or u or c or g or nothing

<400> 209  
gggcgaccnu gaugaguau gcuaaggagg auuuccgaaa gcggnuacgg uccgccagug 60  
aaucgaaacg gugaaagccg uagguugcc 89

<210> 210  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (9)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (45)  
<223> Wherein n is an a or u or c or g or nothing

<400> 210  
gggcgaccnu gaugaguau gcuaaggagg auuuccgaaa gcggnuacgg uccgccaguc 60  
agacgaaacg gugaaagccg uagguugcc 89

<210> 211  
<211> 89

<212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (18)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (63)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 211  
 gggcgaccnu gaugagungc gcuaaggagg auuuccguua gcggcuacgg uccgccaguu 60  
 auncgaaacg gugaaagccg uagguugcc 89

<210> 212  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 212  
 gggcgaccnu gaugagcaac gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
 ccacgaaacg gugaaagccg uagguugcc 89

<210> 213  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK Sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (60)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (63)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 213

gggcgacccu gaugagucgc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagun 60  
 aancgaaacg gugaaagccg uagguugcc 89

<210> 214  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 214  
 gggcgacccu gaugagucac gcuaaggagg auuuccgaaa gcggnuacgg uccgccaguu 60  
 acacgaaacg gugaaagccg uagguugcc 89

<210> 215  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (5)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (27)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (30)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (38)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (41)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (53)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (73)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (86)  
<223> wherein n is an a or u or c or g or nothing

<400> 215  
gggcnaccuc ugaguagcag gcuaagnagn auuuccgnaa ncggnuacgg ucngccagua 60  
ugacgaaacg gunaaagccg uaggungcc 89

<210> 216  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (45)..(46)  
<223> Wherein n is an a or u or c or g or nothing

<400> 216  
gggcgacccu gaugagcugu gcuaaggagg auuuccgaaa gcggnnacgg uccgccaguu 60  
ucccgaaacg gugaaagccg uagguugcc 89

<210> 217  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (5)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (19)..(20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature

<222> (27)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (37)..(38)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (41)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (60)..(63)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (66)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (73)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 217  
 gggcnaccuu gaugagcann gcuaagnagu auuuccnnaa ncggauacgg uccgccagun 60  
 nnncgnaacg gunaaagccg uagguugcc 89  
  
 <210> 218  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <400> 218  
 gggcgacccu gaugagcaua gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
 cgacgaaacg gugaaagccg uagguugcc 89  
  
 <210> 219  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (45)  
<223> Wherein n is an a or u or c or g or nothing

<400> 219  
gggcgaccnu gaugagcucu gcuaaggagg auuuccgaaa gcggnuacgg uccgccaguu 60  
caacgaaacg gugaaagccg uagguugcc 89

<210> 220  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 220  
gggcgaccuu gaugagacca gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
auucgaaacg gugaaagccg uagguugcc 89

<210> 221  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (9)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (18)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (45)  
<223> Wherein n is an a or u or c or g or nothing

<400> 221  
gggcgaccnu gaugagcnug gcuaaggagg auuuccgaaa gcggnuacgg uccgccagug 60  
uuacgaaacg gugaaagccg uagguugcc 89

<210> 222  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 222  
 gggcgaccnu gaugagucuc gcuaaggagg auuuccgaaa gcggnuacgg uccgccagua 60  
 augcgaaacg gugaaagccg uagguugcc 89

<210> 223  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (5)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (22)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (41)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (73)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 223  
 gggcnaccuu gaugaggugg gnuaaggagu auuuccgaaa ncggauacgg uccgccagua 60  
 acccgaaacg gunaaagccg uagguugcc 89

<210> 224  
 <211> 88  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature



<222> (4)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (37)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (58)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (60)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 224  
 gggngacccu gaugaguagg gcuaaggagg auuuccnaaa gcggcuacgg uccgccanan 60  
 gucgaaacgg ugaaagccgu agguugcc 88

<210> 225  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (4)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (30)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (41)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (53)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 225  
 gggngacuu gaugaguagc gcuaaggagn auuuccgaaa ncggnuacgg ucngccaguu 60  
 gagcgcaacg gugaaagccg uagguugcc 89

<210> 226  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 226  
 gggcgacccu gaugagcucu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 ggucgaaacg gugaaagccg uagguugcc 89

<210> 227  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 227  
 gggcgacccu gaugagggcu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 aaacgaaacg gugaaagccg uagguugcc 89

<210> 228  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (4)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (30)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (37)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (46)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 228  
 gggngacccu gaugagccau gcuaacgagn auuucunaaa gcggcnacgg uccgccaguu 60  
 ucgcgaaacg gugaaagccg uagguugcc 89

<210> 229  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (9)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (27)  
<223> Wherein n is an a or u or c or g or nothing

<400> 229  
gggcgaccnu gaugagucac gcuaagnagg auuuccgaaa gcggcuacgg uccgccaguc 60  
ucacgaaacg gugaaagccg uagguugcc 89

<210> 230  
<211> 90  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (50)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (53)..(54)  
<223> Wherein n is an a or u or c or g or nothing

<400> 230  
gggcgacccu gaugagacuu gguaaggagg auuuccgaaa ncgnnuacgg uccgccaugu 60  
uuagcgaaac ggugaaagcc guagguugcc 90

<210> 231  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature

<222> (9)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (45)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 231  
 gggcgaccnu gaugagnuuu gcuaaggagg auuuccgaaa gcggnuacgg uccgccaguu 60  
 uaucgaaacg gugaaagccg uagguugcc 89  
  
 <210> 232  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (45)..(46)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 232  
 gggcgaccnu gaugagcuac gcuaaggagg auuuccgaaa gcggnnacgg uccgccaguu 60  
 ccacgaaacg gugaaagccg uagguugcc 89  
  
 <210> 233  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (17)..(18)  
 <223> Wherein n is an a or u or c or g or nothing

```

<220>
<221> misc_feature
<222> (20)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (22)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (30)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (46)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (53)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (60)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (62)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature
<222> (66)
<223> Wherein n is an a or u or c or g or nothing

<400> 233
gggcgaccnu gaugagnncn gnuaaggagn auuuccgaaa gcgganacgg ucngccagun 60
anccgnaacg gugaaagccg uagguugcc 89

<210> 234
<211> 89
<212> RNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: ERK sensitive
        hammerhead nucleic acid sensor molecule

<220>
<221> misc_feature
<222> (9)
<223> Wherein n is an a or u or c or g or nothing

<220>
<221> misc_feature

```

<222> (45)

<223> Wherein n is an a or u or c or g or nothing

<400> 234

gggcgaccnu gaugaggaau gcuaaggagg auuuccgaaa gcggnuacgg uccgccagua 60  
cagcgaaacg gugaaagccg uagguugcc 89

<210> 235

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (9)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (45)..(46)

<223> Wherein n is an a or u or c or g or nothing

<400> 235

gggcgaccnu gaugaguucu gcuaaggagg auuuccgaaa gcggnnacgg uccgccagua 60  
aagcgaaacg gugaaagccg uagguugcc 89

<210> 236

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (45)

<223> Wherein n is an a or u or c or g or nothing

<400> 236

gggcgacccu gaugaguucu gcuaaggagg auuuccgaaa gcggnuacgg uccgccagua 60  
uuacgaaacg gugaaagccg uagguugcc 89

<210> 237

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature  
 <222> (9)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (20)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (62)..(63)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 237  
 gggcgaccnu gaugagnaun gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 anncgaaacg gugaaagccg uagguugcc 89

<210> 238  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 238  
 gggcgaccnu gaugaggaau gcuaaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 aagcgaaacg gugaaagccg uagguugcc 89

<210> 239  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 239  
 gggcgaccnu gaugagcagu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 gagcgaaacg gugaaagccg uagguugcc 89

<210> 240  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (17)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (63)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 240  
 gggcgacccu gaugagnuau gcuaaggagg auuuccgaaa gcggcuacgg uccgccagun 60  
 agncgaaacg gugaaagccg uagguugcc 89

<210> 241  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 241  
 gggcgacccu gaugagcacu gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 agacgaaacg gugaaagccg uagguugcc 89

<210> 242  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (34)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 242  
 gggcgacccu gaugagcuca gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 ucgcgaaacg gugaaagccg uagguugcc 89

<210> 243  
 <211> 89  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (46)  
 <223> Wherein n is an a or u or c or g or nothing



<400> 243  
gggcgacccu gaugaggcuu gcuaaggagg auuuccgaaa gcggcnacgg uccgccaguc 60  
cgucgaaacg gugaaagccg uagguugcc 89

<210> 244  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 244  
gggcgacccu gaugagcucu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
cggcgaaacg gugaaagccg uagguugcc 89

<210> 245  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 245  
gggcgacccu gaugaggcaa gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
caacgaaacg gugaaagccg uagguugcc 89

<210> 246  
<211> 82  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 246  
gggcgacccu gaugagcguc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
ucucgaaacg gugaaagccg ua 82

<210> 247  
<211> 85  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 247  
gggcgacccu gaugagcuuc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
aaacgaaacg gugaaagccg uaggu 85

<210> 248  
 <211> 75  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
           hammerhead nucleic acid sensor molecule  
  
 <400> 248  
 gggcgacccu gaugaguggc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 uaacgaaacg gugaa 75  
  
 <210> 249  
 <211> 85  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
           hammerhead nucleic acid sensor molecule  
  
 <400> 249  
 gggcgacccu gaugagacuc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
 guacgaaacg gugaaagccg uaggu 85  
  
 <210> 250  
 <211> 91  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
           hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (35)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 250  
 gggcgacccu gaugagauuu gcuaaggagg auuunccgaa agcggcucac ggugccgcca 60  
 guugagcgaa acggugaaaag ccuagguugc c 91  
  
 <210> 251  
 <211> 88  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
           hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (18)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 251  
gggcgacccu gaugagcnuc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
auacgaaacg gugaaagccg uagguugc 88

<210> 252  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (19)..(20)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (62)  
<223> Wherein n is an a or u or c or g or nothing

<400> 252  
gggcgacccu gaugagcunn gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
cnacgaaacg gugaaagccg uagguugc 89

<210> 253  
<211> 85  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 253  
gggcgacccu gaugaguucg gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
aagcgaaacg gugaaagccg uaggu 85

<210> 254  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 254  
gggcgacccu gaugagauga gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
aaccgaaacg gugaaagccg uagguugc 89

<210> 255  
<211> 77  
<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (40)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (44)

<223> Wherein n is an a or u or c or g or nothing

<400> 255

gggcgacccu gaygagaau cgcuaaggagg auuuccgaan gcgncuacgg uccgccaguu 60  
ccccgaaacg gugaaag 77

<210> 256

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: BRK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (62)

<223> Wherein n is an a or u or c or g or nothing

<400> 256

gggcgacccu gaugagucac gcuaaggagg auuuccgaaa gcgguacgg uccgccaguu 60  
gncgcgaaac ggugaaagcc guagguugcc 90

<210> 257

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: BRK sensitive  
hammerhead nucleic acid sensor molecule

<400> 257

gggcgacccu gaugagcauc gcuaaggagg auuuccgaaa gcgguacgg uccgccaguu 60  
aagcgaaacg gugaaagccg uagguugcc 89

<210> 258

<211> 84

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: BRK sensitive

## hammerhead nucleic acid sensor molecule

&lt;400&gt; 258

```

gggcgacccu gaugagguca gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60
cagcgaaaacg gugaaagccg uagg                                     84

```

&lt;210&gt; 259

&lt;211&gt; 79

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

&lt;400&gt; 259

```

gggcgacccu gaugagaagc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguc 60
uagcgaaaacg gugaaagcc                                     79

```

&lt;210&gt; 260

&lt;211&gt; 90

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

&lt;400&gt; 260

```

gggcgacccu gaugaggcuu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60
gaucgaaaacg gugaaagccg ugagguugcc                               90

```

&lt;210&gt; 261

&lt;211&gt; 91

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

&lt;400&gt; 261

```

gggcgacccu gaugaggcug gcuaaggagg auuuccgaaa gcggcucacg guccgccagu 60
ccuaacgaaa cggugaaagc cguagguugc c                               91

```

&lt;210&gt; 262

&lt;211&gt; 88

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

&lt;400&gt; 262

```

gggcgacccu gaugagccuu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60
agucgaaaacg gugaaagccg uagguugc                                     88

```

<210> 263  
 <211> 67  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (61)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 263  
 gggcgacccu gaugagucac gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 ncgcgaa 67

<210> 264  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 264  
 gggcgacccu gaugagaacu gcuaaggagg auuuccgaaa agcgguacgg guccgccagu 60  
 ugcgcgaaac ggugaaagcc guagguugcc 90

<210> 265  
 <211> 84  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 265  
 gggcgacccu gaugagcgac gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
 gggcgaaaac gugaaagccg uagg 84

<210> 266  
 <211> 85  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<400> 266  
 gggcgacccu gaugaguaac gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguc 60  
 agacgaaaac gugaaagccg uaggu 85

<210> 267

<211> 85

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 267

gggcgacccu gaugaggacu gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
acucgaaacg gugaaagccg uaggu 85

<210> 268

<211> 89

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 268

gggcgacccu gaugagccgg gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
gagcgaaacg gugaaagccg uagguugcc 89

<210> 269

<211> 41

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (19)

<223> Wherein n is an a or u or c or g or nothing

<400> 269

gggcgacccu gaugagcana gcuaaggagg auuuccgaaa g 41

<210> 270

<211> 85

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 270

gggcgacccu gaugaguugc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
uagcgaaacg gugaaagccg uaggu 85

<210> 271  
<211> 85  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 271  
gggcgacccu gaugagcagc gcuaaggagg auuuccgaaa gcggcuaggu ccgccagucu 60  
ucgaaacggg gaaagccgua gguug 85

<210> 272  
<211> 75  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 272  
gggcgacccu gaugaguaag gcuaaggagg auuuccgaaa gcggcuacgg uccgccagug 60  
uugcgaaacg gugaa 75

<210> 273  
<211> 89  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 273  
gggcgacccu gaugagcacc gcuaaggagg auuuccgaaa gcggcuacgg uccgccaguu 60  
acgcgaaacg gugaaagccg uagguugcc 89

<210> 274  
<211> 82  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 274  
gggcgacccu gaugaggcau gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
aggcgaaacg gugaaagccg ua 82

<210> 275  
<211> 77  
<212> RNA  
<213> Artificial Sequence



<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 275

gggcgacccu gaugaguua gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
gaccgaaacg gugaaag 77

<210> 276

<211> 77

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<400> 276

gggcgacccu gaugagaagc gcuaaggagg auuuccgaaa gcggcuacgg uccgccagua 60  
augcgaaacg gugaaag 77

<210> 277

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (46)

<223> Wherein n is an a or u or c or g or nothing

<400> 277

gggugaccuc ugauaguguu gcuaaggagg gauuuccgaa agcggnuacg guccgccagu 60  
agagcgaaac ggugaaagcc guagguugcc 90

<210> 278

<211> 90

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>

<221> misc\_feature

<222> (9)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (19)

<223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (46)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 278  
 gggcgaccnu' gaugagcang cgcuaaggag gauuuccgaa agcggnuacg guccgccagu 60  
 gacucgaaac ggugaaagcc guagguugcc 90

<210> 279  
 <211> 80  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (33)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (70)..(71)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (76)  
 <223> Wherein n is an a or u or c or g or nothing

<400> 279  
 gggcgacccu gauaguagcg cuaaguaagg aunuuccgaa agcgcgcuca cgguccgcca 60  
 gucucacgan ncggunaaag 80

<210> 280  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule

<220>  
 <221> misc\_feature  
 <222> (4)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (20)  
 <223> Wherein n is an a or u or c or g or nothing

<220>  
 <221> misc\_feature

<222> (52)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (65)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 280  
 gggngacccu gaugaguaun agcuaaggag gauuuccgaa agcgguacg gnccgccagu 60  
 aaugngaaac ggugaaagcc guagguugcc 90  
  
 <210> 281  
 <211> 90  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <220>  
 <221> misc\_feature  
 <222> (19)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (23)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <220>  
 <221> misc\_feature  
 <222> (46)  
 <223> Wherein n is an a or u or c or g or nothing  
  
 <400> 281  
 gggcgaccuu gaugaagcnc agnuaaggag gauuuccgaa agcggnuacg gucagccagu 60  
 ggcacgaaac ggugaaagcc guagguugcc 90  
  
 <210> 282  
 <211> 88  
 <212> RNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 hammerhead nucleic acid sensor molecule  
  
 <400> 282  
 gggcgacccu gaugagcaug cuaaggagga uuuccgaaag cgguacggg ccgccaguuu 60  
 gucgaaacgg ugaaagccgu agguugcc 88  
  
 <210> 283  
 <211> 87  
 <212> RNA  
 <213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ERK sensitive  
hammerhead nucleic acid sensor molecule

<220>  
<221> misc\_feature  
<222> (16)..(17)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (22)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (40)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (56)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (60)..(61)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (72)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (76)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (81)  
<223> Wherein n is an a or u or c or g or nothing

<220>  
<221> misc\_feature  
<222> (86)  
<223> Wherein n is an a or u or c or g or nothing

<400> 283  
gggcgacccu gauganngcu gnuaggagga uuuccgaaan ugcucuggc ccgcanuuan 60  
nccgaacggg cnaaancgaa nggugnc 87

<210> 284  
<211> 65  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: BRK sensitive

## hammerhead nucleic acid sensor molecule

&lt;400&gt; 284

```

gggcgacccu gaugaguagg cuaaggagga uuucgaaagc ggcuaagguc cgccagugug 60
ccgaa                                         65

```

&lt;210&gt; 285

&lt;211&gt; 88

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: BRK sensitive  
hammerhead nucleic acid sensor molecule

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (18)

&lt;223&gt; Wherein n is a or u or c or g or nothing.

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (25)

&lt;223&gt; Wherein n is a or u or c or g or nothing.

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (50)

&lt;223&gt; Wherein n is a or u or c or g or nothing.

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (60)

&lt;223&gt; Wherein n is a or u or c or g or nothing.

&lt;400&gt; 285

```

gggcgacccu gcugagcnac gcuaaggagga uuuccgaaag cggcuacggn ccgccagucn 60
gacgaaacgg ugaaagccgu agguugcc                                         88

```

&lt;210&gt; 286

&lt;211&gt; 87

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Construct 1-14

&lt;400&gt; 286

```

gggcgacccu gaugaggguc gcuaaggagg auuuccgaaa gcggcacggu ccgccagacg 60
ucgaaacggu gaaagccgua gguugcc                                         87

```

&lt;210&gt; 287

&lt;211&gt; 87

&lt;212&gt; RNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Construct 1-13

<400> 287

gggcgacccu gaugagucuu cuaaggagga uuuccgaaag cggcacgguc cgccaguacg 60  
ucgaaacggg gaaagccgua gguugcc 87

<210> 288

<211> 86

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 1-2

<400> 288

gggcgacccu gaugagccuu cuaaggagga uuuccgaaag cggcacgguc cgccagacgu 60  
cgaaacggug aaagccguag guugcc 86

<210> 289

<211> 87

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 1-6

<400> 289

gggcgacccu gaugagucuu gcuaaggagg auuuccgaaa gcggcacggg ccgccagacg 60  
ucgaaacggg gaaagccgua gguugcc 87

<210> 290

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-7

<400> 290

gggcgacccu gaugagccuu gcuaaggagg auuuccgaaa gcggcacggg ccgccaguca 60  
gucgaaacgg ugaaagccgu agguugcc 88

<210> 291

<211> 87

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-2

<400> 291

gggcgacccu gaugagccuc gcuaaggagg auuuccgaaa gcggcacggg ccgccagcua 60  
gcgaaacggg gaaagccgua gguugcc 87

<210> 292

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-3

<400> 292

gggcgacccu gaugagaacu gcuaaggagg auuuccgaaa gcggcacggu ccgccaguuu 60  
aacgaaacgg ugaaagccgu agguugcc 88

<210> 293

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-13

<400> 293

gggcgacccu gaugaguaaa gcuaaggagg auuuccgaaa gcggcacggu ccgccagucu 60  
ggcgaaacgg ugaaagccgu agguugcc 88

<210> 294

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-14

<400> 294

gggcgacccu gaugaguuuu gcuaaggagg auuuccgaaa gcggcacggu ccgccaguga 60  
uccgaaacgg ugaaagccgu agguugcc 88

<210> 295

<211> 88

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 2-20

<400> 295

gggcgacccu gaugaggcga gcuaaggagg auuuccgaaa gcggcacggu ccgccaguuu 60  
aacgaaacgg ugaaagccgu agguugcc 88

<210> 296

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Combined DNA/RNA Molecule:  
Oligonucleotide substrate

<220>

<223> Description of Artificial Sequence: 5' primer

<400> 296  
tctaatacga ctactatag gacctcggcg aaagc 35

<210> 297  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: 3' Primer A

<400> 297  
actctcgcta acctctctag tcata 25

<210> 298  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: 3' Primer B

<400> 298  
agtgcctctcg cacctaacct ctctagt 27

<210> 299  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: 3' Primer C

<400> 299  
agtgcgagcc tctcgctcg cacctaacct ctctagt 37

<210> 300  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer #1 from  
first set

<400> 300  
tctaatacga ctactatag gacctcggcg aaagc 35

<210> 301  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer #2 from  
first set



<400> 301  
agtgtctctcg cacctaacct ctctagt 27

<210> 302  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer 1 from  
second set

<400> 302  
gttgctacaa atgatatgac 20

<210> 303  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer #2 from  
second set

<400> 303  
atggcaattt agccatgaga 20

<210> 304  
<211> 106  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Ligase RNA  
sequence

<400> 304  
ggaccucggc gaaagcuaac gucucauggc uaaaauugcca uguugcuaca aaugauauga 60  
cuagagaggu uaggugcauc uucaugucca gucgcuugca augccc 106

<210> 305  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Reverse  
transcriptase primer

<400> 305  
gggcattgca agcgactgga cat 23

<210> 306  
<211> 28  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide substrate

<400> 306

actgaacctg accgtacaaa gatgcacu

28

<210> 307

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: TaqMan forward  
primer

<400> 307

actgaacctg accgtacaaa ga

22

<210> 308

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: TaqMan reverse  
primer

<400> 308

tttgtagcaa catggcaatt ta

22

<210> 309

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: TaqMan probe  
primer

<400> 309

cggcgaaagc taacgtctca tgg

23

<210> 310

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Forward primer

<400> 310

actgaacctg accgtacaaa ga

22

<210> 311  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Reverse primer

<400> 311  
tttgtagcaa catggcaatt ta 22

<210> 312  
<211> 95  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Lysozyme  
dependent nucleic acid sensor molecule

<400> 312  
ggaccucggc gaaagcuaac gucucauggc uaaauugcca uguugcuaca aaugauauga 60  
cuagagaggu uaggugccuc gugaugucca gucgc 95

<210> 313  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Effector oligo

<400> 313  
gcgactggac atcacgag 18

<210> 314  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Combined DNA/RNA Molecule:  
Substrate

<220>  
<223> Description of Artificial Sequence:  
Oligonucleotide substrate

<400> 314  
gtacgatgcy atgctagcga ttgttgugca cu 32

<210> 315  
<211> 42  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Forward primer

<400> 315

taatacgact cactatagga ctccggcgaa agccgttcga cc

42

<210> 316

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer  
spanning the 3' proximal portion of the L1 ligase  
domain

<400> 316

attcgagatg tccttgacc aaagccgcac ctaacctct gtctaag

47

<210> 317

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Forward primer

<400> 317

gcgaccttac gatcagatga c

21

<210> 318

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Reverse primer

<400> 318

ccgcacctaa cctcctgtct aa

22

<210> 319

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ERK ligase  
probe primer

<400> 319

aaggaggatt tccgaaagcg gctacg

26

<210> 320

<211> 26

<212> DNA

<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: ppERK ligase  
probe primer

<400> 320  
cgctagcgaa ttggttcctc gaaagg

26

<210> 321  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Combined DNA/RNA Molecule:  
Substrate

<220>  
<223> Description of Artificial Sequence: Substrate 3

<400> 321  
catgcgacct tacgatacaga tgacctugca cu

32

<210> 322  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Substrate 3  
specific 5' PCR primer

<400> 322  
catgcgacct tacgatacaga t

21

<210> 323  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Regeneration  
5' PCR primer

<400> 323  
tttaatacga ctactatag gacttcggcg aaagc

35

<210> 324  
<211> 109  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Library (DNA)  
(random regions 3-5 nucleotides in length)

<220>

<221> misc\_feature  
 <222> (26)..(30)  
 <223> Wherein n is an a or t or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (64)..(68)  
 <223> Wherein n is an a or t or c or g or nothing

<400> 324  
 ggacttcggc gaaagccgtt cgaccnnnnn aaggaggatt tccgaaagcg gctacggtcc 60  
 gccnnnnnct tagacaggag gttaggtgcg taggtaaccg atagtccg 109

<210> 325  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Effector  
 primer

<400> 325  
 cggaactatc gggtacctac 20

<210> 326  
 <211> 135  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ppERK  
 selection-specific oligonucleotides

<220>  
 <221> misc\_feature  
 <222> (26)..(31)  
 <223> Wherein n is an a or t or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (89)..(94)  
 <223> Wherein n is an a or t or c or g or nothing

<400> 326  
 ggacttcggc gaaagccgtt cgaccnnnnn ncagacgcta gcgaattggt tcctcgaaag 60  
 gggaaagcgt tattaagaaa ccaaaatggn nnnncttaga caggagggtta ggtgcgtcaa 120  
 tgctgcaagt tactg 135

<210> 327  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Effector  
 primer

<400> 327  
cagtaacttg cagcattgac

20

<210> 328  
<211> 101  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: bFGF selection  
specific oligonucleotide

<220>  
<221> misc\_feature  
<222> (26)..(32)  
<223> Wherein n is an a or t or c or g or nothing

<220>  
<221> misc\_feature  
<222> (54)..(60)  
<223> Wherein n is an a or t or c or g or nothing

<400> 328  
ggacttcggc gaaagccgtt cgaccnnnnn nngcaacgct acagacaagt gcannnnnnn 60  
cttagacagg aggttaggtg cccgagttgt tcgaacgaga c 101

<210> 329  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Effector  
primer

<400> 329  
gtctcggtcg aacaactcgg

20

<210> 330  
<211> 92  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Thrombin  
selection specific oligonucleotide

<220>  
<221> misc\_feature  
<222> (26)..(31)  
<223> Wherein n is an a or t or c or g or nothing

<220>  
<221> misc\_feature  
<222> (47)..(51)  
<223> Wherein n is an a or t or c or g or nothing

<400> 330

ggacttcggc gaaagccgtt cgaccnnnnn nategaagtt agtaggnnnn ncttagacag 60  
gaggtaggt gcgtcaatcg attgcagatc cg 92

<210> 331  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Effector  
primer

<400> 331  
cggatctgca atcgattgac 20

<210> 332  
<211> 77  
<212> RNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Nucleic acid  
sensor molecule with effector domain deleted

<400> 332  
ggaccucggc gaaagcuaac gucauggc uaaaugcca uguugcuaca aaugauuga 60  
cuagagaggu uaggugc 77

<210> 333  
<211> 83  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: MK.08.92.A

<400> 333  
gcttgcaagc ccttagaccc tgatgagcct tgcgatgcaa aaaggtgctg acgacacatc 60  
gaaacggtga aagccgtagg tct 83

<210> 334  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: MK.08.66.B

<400> 334  
agacctacgg ctttcaccgt ttcg 24

<210> 335  
<211> 57  
<212> DNA  
<213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: MK.08.130.B

&lt;400&gt; 335

atacgactca ctataggatg tccagtcgct tgcaatgcc ttttagacct tgatgag 57

&lt;210&gt; 336

&lt;211&gt; 82

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
LYS.LIG.TMPLT.L1-11.2

&lt;400&gt; 336

ctataggact tcggcgaaag ctaacgtctc atggctaaat tgccatgttg ctacaaatga 60  
tatgactaga gaggttaggt gc 82

&lt;210&gt; 337

&lt;211&gt; 92

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
FMN.LIG.TMPLT.L1-R7C13

&lt;400&gt; 337

ctataggact tcggtccagt gctcgtgcac taggccgttc gaccttcagg atatgcttcg 60  
gcagaaggga acttagacag gaggttaggt gc 92

&lt;210&gt; 338

&lt;211&gt; 101

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence:  
THBO.LIG.TMPLT.L1-D1

&lt;400&gt; 338

ctataggact tcggtccagt gctcgtgcac taggccgttc gacctgata ccagcatcgt 60  
cttgatgccc ttggcagcat cttagacagg aggttaggtg c 101

&lt;210&gt; 339

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: TK.16.32.A

&lt;400&gt; 339

ttctaatacg actcactata ggacttc

27

<210> 340  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: TK.16.32.B

<400> 340  
attcgagatg tccttggacc aaagccgcac ctaacctc

38

<210> 341  
<211> 48  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Combined DNA/RNA Molecule:  
Substrate

<220>  
<223> Description of Artificial Sequence: TK.16.32.15NT

<400> 341  
attcgagatg tccttggacc aaagcctcca tcgtgcgcac ctaacctc

48

<210> 342  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Combined DNA/RNA Molecule:  
Substrate

<220>  
<223> Description of Artificial Sequence: TK.04.82.A

<400> 342  
catgcgacct tacgatacaga tgacctugca cu

32

<210> 343  
<211> 15  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: MK.08.125A

<400> 343  
tccatcgtgc gcacu

15

<210> 344  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: TK16.32.A

<400> 344

ttctaatacgt actcactata ggacttc

27

<210> 345

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: MK.08.125B

<400> 345

tccatcgtgc gcacctaacc tc

22

<210> 346

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: MK.08.125A

<400> 346

tccatcgtgc gcacu

15

<210> 347

<211> 55

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Combined DNA/RNA Molecule:  
Oligonucleotide substrate

<220>

<223> Description of Artificial Sequence: 2-piece ligase  
template

<220>

<221> misc\_feature

<222> (26)..(33)

<223> Wherein n is an a or u or c or g or nothing

<400> 347

ggacuucggc gaaagccguu cgaccnnnnn nnnccuagac aggagguuag gugcg

55

<210> 348

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide substrate

<400> 348  
tccatcgtgc gcacu

15

<210> 349  
<211> 99  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 29

<400> 349  
ggacuucggc gaaagccguu cgaccacgcu aaggaggauu uccgaaagcg gcuacgugcc 60  
gccagugucu uagacaggag guuaggugcg cacgaugga 99

<210> 350  
<211> 6  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide substrate

<400> 350  
ugcacu

6

<210> 351  
<211> 97  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Construct 30

<400> 351  
ggacuucggc gaaagccguu cgaccagcua aggaggauu ccgaaagcgg cuacgugccg 60  
ccagcucua gacaggaggu uaggugcgca cgaugga 97

<210> 352  
<211> 138  
<212> RNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ppERK  
dependent ligase template

<220>

<221> misc\_feature

<222> (27)..(30)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (87)..(90)

<223> Wherein n is an a or u or c or g or nothing

<400> 352

```
ggacuucggc gaaagccguu cgaccannnn cagacgcuag cgaauugguu ccucgaaagg 60
gaaagcguua uuaagaaacc aaaaugnnnn ucuuagacag gagguuaggu gccguccgac 120
ugaucucgga guuaaacg                                     138
```

<210> 353

<211> 107

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ppERK  
sensitive hammerhead template

<220>

<221> misc\_feature

<222> (17)..(21)

<223> Wherein n is an a or u or c or g or nothing

<220>

<221> misc\_feature

<222> (76)..(80)

<223> Wherein n is an a or u or c or g or nothing

<400> 353

```
gggcgacccu gaugagnnnn nagacgcuag cgaauugguu ccucgaaagg ggaaagcguu 60
auuaagaaac caaaannnnn cgaaacggug aaagccguag guugccc          107
```

<210> 354

<211> 134

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ppERK nucleic  
acid sensor molecule template

<220>

<221> misc\_feature

<222> (25)..(31)

<223> Wherein n is an a or t or c or g or nothing

<220>

<221> misc\_feature

<222> (88)..(93)

<223> Wherein n is an a or t or c or g or nothing

<400> 354

```
ggacttcggc gaaagccggt cgaccnnnnn ncagacgcta gcgaattggt tcctcgaaag 60
ggaaagcggt attaagaaac caaaatgnnn nnnccttagac aggaggtag gtgcgtcaat 120
gctgcaagtt actg                                     134
```

<210> 355

<211> 109

<212> DNA

<213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: ERK sensitive  
 nucleic acid sensor molecule template

<220>  
 <221> misc\_feature  
 <222> (26)..(30)  
 <223> Wherein n is an a or t or c or g or nothing

<220>  
 <221> misc\_feature  
 <222> (64)..(68)  
 <223> Wherein n is an a or t or c or g or nothing

<400> 355  
 ggacttcggc gaaagccgtt cgaccnnnnn aaggaggatt tccgaaagcg gctacggtcc 60  
 gccnnnnnct tagacaggag gttaggtgcg taggtaaccg atagttccg 109

<210> 356  
 <211> 131  
 <212> RNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: CW45-33-A08  
 ERK sensitive nucleic acid sensor molecule

<220>  
 <223> Description of Combined DNA/RNA Molecule:  
 Oligonucleotide substrate

<400> 356  
 ggacuucggc gaaagccguu cgaccucuc agacgcuagc gaaugguuc cucgaaaggg 60  
 aaagcguuau uaagaaacca aaagagagg cuagacagg agguuaggug cguaaugcu 120  
 gcaaguacu g 131

<210> 357  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:  
 Oligonucleotide substrate

<400> 357  
 acgtagcata gcatcgatag ctgttgugca cu 32

<210> 358  
 <211> 265  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Thymidylate  
 synthase ("td") intron

&lt;400&gt; 358

taattgaggc ctgagtataa ggtgacttat acttgtaatc tatctaaacg gggaacctct 60  
 ctagtagaca atccccgtgct aaattgtagg actgcccggg ttctacataa atgcctaacg 120  
 actatccctt tggggagtag ggtcaagtga ctcgaaacga tagacaactt gctttaacaa 180  
 gttggagata tagtctgctc tgcattggtga catgcagctg gatataattc cggggtaaga 240  
 ttaacgacct tatctgaaca taatg 265

&lt;210&gt; 359

&lt;211&gt; 276

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Theophylline dependent td group I intron

&lt;400&gt; 359

taattgaggc ctgagtataa ggtgacttat acttgtaatc tatctaaacg gggaacctct 60  
 ctagtagaca atccccgtgct aaattgatac cagcatcgtc ttgatgccct tggcagcata 120  
 aatgcctaac gactatccct ttggggagta gggcaagtgc actcgaaacg atagacaact 180  
 tgctttaaca agttggagat atagtctgct ctgcatggtg acatgcagct ggatataatt 240  
 ccggggtaag attaacgacc ttatctgaac ataatt 276

&lt;210&gt; 360

&lt;211&gt; 10

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Exon 1, the 5'-exon

&lt;400&gt; 360

tttcttgggt

10

&lt;210&gt; 361

&lt;211&gt; 10

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Exon 2, the 3'-exon

&lt;400&gt; 361

ctaccgttta

10

&lt;210&gt; 362

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: PCR primer

&lt;400&gt; 362

agtgtctctg cacctaacct cctgtct

27

<210> 363  
<211> 16  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 363  
ggacttcggc gaaagc

16

<210> 364  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 364  
agtgtctctg cacctaacct cctgtct

27

<210> 365  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 365  
gctacgggcc gccagttctt

20

<210> 366  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 366  
cgctttcggc aatcctcctt

20

<210> 367  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 367  
gctacgggcc gccaggggct

20



<210> 368  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 368  
cgctttcgga aatcctcctt

20

<210> 369  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 369  
gctacgggcc gccaaaagct

20

<210> 370  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 370  
cgctttcgga aatcctcctt

20

<210> 371  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 371  
aaggggaaag cggtattaag

20

<210> 372  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: PCR primer

<400> 372  
tcgaggaacc aatcgctag

20

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number  
WO 03/014375 A3

(51) International Patent Classification<sup>7</sup>: C12Q 1/68,  
C07H 21/02, 21/04

(21) International Application Number: PCT/US02/25319

(22) International Filing Date: 9 August 2002 (09.08.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/311,378	9 August 2001 (09.08.2001)	US
60/313,932	21 August 2001 (21.08.2001)	US
09/952,680	13 September 2001 (13.09.2001)	US
60/338,186	13 November 2001 (13.11.2001)	US
60/349,959	18 January 2002 (18.01.2002)	US
60/364,486	13 March 2002 (13.03.2002)	US
60/367,991	25 March 2002 (25.03.2002)	US
60/369,887	4 April 2002 (04.04.2002)	US
60/376,744	1 May 2002 (01.05.2002)	US
60/385,097	31 May 2002 (31.05.2002)	US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US	60/311,378 (CIP)
Filed on	9 August 2001 (09.08.2001)
US	60/313,932 (CIP)
Filed on	21 August 2001 (21.08.2001)
US	60/338,186 (CIP)
Filed on	13 November 2001 (13.11.2001)
US	60/349,959 (CIP)
Filed on	18 January 2002 (18.01.2002)
US	60/364,486 (CIP)
Filed on	13 March 2002 (13.03.2002)
US	60/367,991 (CIP)
Filed on	25 March 2002 (25.03.2002)
US	60/369,887 (CIP)
Filed on	4 April 2002 (04.04.2002)
US	60/376,744 (CIP)
Filed on	1 May 2002 (01.05.2002)
US	60/385,097 (CIP)
Filed on	31 May 2002 (31.05.2002)
US	09/952,680 (CIP)
Filed on	13 September 2001 (13.09.2001)

(72) Inventors; and

(75) Inventors/Applicants (for US only): STANTON, Martin [US/US]; 64 Harvard Road, Stow, MA 01775 (US). EPSTEIN, David [US/US]; 56 Pine Street, Belmont, MA 02478 (US). HAMAGUCHI, Nobuko [JP/US]; 7 Dartmouth Drive, Framingham, MA 01701 (US). KURZ, Markus [DE/US]; 44 Pine Street, Newton, MA 02465 (US). KEEFE, Tony [GB/US]; 97 1/2 Thorndike Street, #1, Cambridge, MA 02139 (US). WILSON, Charles [US/US]; 229 Lexington Road, Concord, MA 01742 (US). GRATE, Dilara [US/US]; 16 Wheelock Terrace, Waltham, MA 02453 (US). MARSHALL, Kristin, A. [US/US]; 15 College Avenue, Arlington, MA 02472 (US). MCCAULEY, Thomas [US/US]; 7 Spencer Avenue, Somerville, MA 02144 (US). KURZ, Jeffrey [US/US]; 431 Broadway, Apartment 23, Somerville, MA 02145 (US).

(74) Agent: ELRIFI, Ivor, R.; Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
16 October 2003

(71) Applicant (for all designated States except US): AR-  
CHEMIX CORPORATION [US/US]; 20 Hampden  
Street, Boston, MA 02111 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACID SENSOR MOLECULES AND METHODS OF USING SAME

(57) Abstract: Methods for engineering a nucleic acid sensor molecule are provided. Biosensors comprise a plurality of nucleic acid sensor molecules labeled with a first signaling moiety and a second signaling moiety. The nucleic acid sensor molecules recognizes target molecules which do not naturally bind to DNA. Binding of a target molecule to the sensor molecules triggers a change in the proximity of the signaling moieties which leads to a change in the optical properties of the nucleic acid sensor molecules on the biosensor. Reagents and systems for performing the method are also provided. The method is useful in diagnostic applications and drug optimization.

WO 03/014375 A3

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/25319

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12Q1/68; C07H 21/02, 21/04

US CL : 435/6; 536/23.1, 24.3

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6; 536/23.1, 24.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Continuation Sheet

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,589,332 A (SHIH et al) 31 December 1996, see at least for example Figure 1C and Column 5 at the paragraph beginning at about line 45.	1-2, 8-11, 37-40, 44-45, 59, and 61-62  3-7, 41-43, 46-58, and 63-71
X --- Y	WO96/27026 A1 (INTELLIGENE LTD.) 06 September 1996, see entire document	1-11, 37-40, 44, 59, 61-62  41-43, 45-58, 60, and 63-71
A	US 6,201,113 B1 (TODD et al) 13 March 2001, see the entire document.	1-11 and 37-71

☒ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Δ" document member of the same patent family

Date of the actual completion of the international search

01 May 2003 (01.05.2003)

Date of mailing of the international search report

18 JUL 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Ethan Whisenant, Ph.D.

Telephone No. (703) 308-0196

*Janice Ford*  
*for*

# INTERNATIONAL SEARCH REPORT

PCT/US02/25319

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO99/47704 A1 (JENNE et al) 23 September 1999, see the entire document.	1-11 and 37-71
A	HASELOFF et al. Simple RNA enzymes with new and highly specific endoribonuclease activities. Nature. 18 August 1988, Vol. 334, pages 585-591, see the entire document.	1-11 and 37-71
A	WO99/50277 A1 (UNIVERSITY OF UTAH RESEARCH FOUNDATION) 07 October 1999, see the entire document.	1-11 and 37-71
A	TUSCHL et al. A three-dimensional model for the hammerhead ribozyme based on fluorescence measurements. Science. 04 November 1994, Vol. 266, pages 785-789, see the entire document.	1-11 and 37-71

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/25319

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:  
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-11 and 37-71

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

**BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING****Lack of Unity**

**1.** This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

**Group 1, Claim(s) 1-11 and 37-71** drawn to a nucleic acid sensor molecule which molecule comprises a catalytic domain which catalytic domain comprises a *cis*-endonucleolytic ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 2, Claim(s) 1-11 and 37-69** drawn to a nucleic acid sensor molecule which molecule comprises a catalytic domain which catalytic domain comprises a *trans*-endonucleolytic ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 3, Claim(s) 1-8, 12-14, 37-73, 76, 78-86, 93 94 and 97,** drawn to a nucleic acid sensor molecule which is a self-ligating 1-piece *cis*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 4, Claim(s) 1-8, 12-14, 37-73, 94, and 97,** drawn to a nucleic acid sensor molecule which is a self-ligating 1-piece *trans*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 5, Claim(s) 1-8, 12-14, 37-73, 95, and 98,** drawn to a nucleic acid sensor molecule which is a self-ligating 2-piece *cis*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 6, Claim(s) 1-8, 12-14, 37-73, 95, and 98,** drawn to a nucleic acid sensor molecule which is a self-ligating 2-piece *trans*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 7, Claim(s) 1-8, 12-14, 37-73, 96, and 99,** drawn to a nucleic acid sensor molecule which is a self-ligating 3-piece *cis*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 8, Claim(s) 1-8, 12-14, 37-69, 96 and 99,** drawn to a nucleic acid sensor molecule which is a self-ligating 3-piece *trans*-ligase ribozyme and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 9, Claim(s) 1-8, 15-16, 18, 21, 23, 37-74, 78-88, 89 and 90,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a protein kinase and wherein said protein kinase is ERK1 or ERK2 and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

## INTERNATIONAL SEARCH REPORT

PCT/US02/25319

**Group 10, Claim(s) 1-8, 15-16, 18, 29 and 37-73,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a protein kinase and wherein said protein kinase is RAF kinase and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 11, Claim(s) 1-8, 15-16, 18, 32 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a GTP binding protein and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 12, Claim(s) 1-8, 15-16, 18, 31 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a phosphatase and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 13, Claim(s) 1-8, 15-16, 18, 31, 34 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a cytokine and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 14, Claim(s) 1-8, 15-16, 30, and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is a Ras protein and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 15, Claim(s) 1-8, 15-16, 18, 31, 33 and 37-69** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a protein, wherein said protein is GPCR and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 16, Claim(s) 1-8, 15-17, and 37-69** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes post-translationally modified form of a protein wherein the post-translational modification is selected from a defined group which includes prenylation, glycosylation, methionine removal, N-acetylation, acylation acylation of cysteines, myristoylation, alkylation, ubiquitinylation, prolyl-4-hydroxylation, carboxylation of glutamyl residues, advanced glycosylation, deamination of glutamine and asparagine, addition of glycosphosphatidylinositol, disulfide bond formation, hydroxylation and lipidation; and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 17, Claim(s) 1-8, 15-17, 19-20, 22, 37-69, 75, 78-86, and 91,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a post-translationally modified form of a protein wherein the post-translational modified protein is phosphorylation and wherein said phosphorylation is monophosphorylation or diphosphorylated and wherein said post-translationally modified diphosphorylated protein is pp ERK and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 18, Claim(s) 1-8, 15, 34 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes peptides and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 19, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes nucleic acids and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 20, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes oligosaccharides and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 21, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes nucleotides and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 22, Claim(s) 1-8, 15 and 36-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes metabolites and/or cellular metabolites and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 23, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes drugs and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 24, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes toxins and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 25, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes biohazards and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 26, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes ions and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 27, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes carbohydrates and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 28, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes polysaccharides and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 29, Claim(s) 1-8, 15, and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes hormones and



# INTERNATIONAL SEARCH REPORT

PCT/US02/25319

composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 30, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes receptors and wherein said receptor is a GPCR and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 31, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes antigens and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 32, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes antibodies and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 33, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a virus and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 34, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a co-factor and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 35, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a dye and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 36, Claim(s) 1-8, 15 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a nutrient and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 37, Claim(s) 1-8, 15, 35 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a growth factor and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 38, Claim(s) 1-8, 24-28 and 37-69,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes a component, product, or associated protein/product of a MAP kinase pathway and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

**Group 39, Claim(s) 1-8, 37-69, 77-86, and 92,** drawn to a nucleic acid sensor molecule which comprises a target modulation domain which recognizes a target selected from a defined group which includes cCMP, cAMP,

# INTERNATIONAL SEARCH REPORT

PCT/US02/25319

or cGMP pathway and composition(s) thereof, as well as, a method of using the nucleic acid sensor molecule defined above in this paragraph.

2. The inventions listed as Groups 1-39 do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical feature(s).

The claims as drawn are related to each other because of the product, i.e. the nucleic acid sensor molecule. However, since the the nucleic acid sensor molecule as claimed (Claim 1) was known - see, for example, Shih et al. US 5,589,332 (31 DEC 1996), see at least Figure 1a of Shih et al.- the claims are no longer linked by a special technical feature, because, by definition, the special technical feature must distinguish over the prior art. Without the special technical feature the claims lack unity.

3. In addition, if the applicant elects to (i.e. pays to) have any of Groups 3-9, 17 or 39 searched the applicant will also be required to elect a single nucleotide sequence for searching from each of Claims 89-93 and 97-99.

## Continuation of B. FIELDS SEARCHED Item 3:

USPATFULL via East, Medline CAplus

search terms: probe?, cis-endonucleolytic adj ribozyme?, ribozyme?, FRET